

Euclid's Elements — Book I

Proposition 19

Formal Reconstruction — Technical Documentation

Jurek Róžański
Military University of Technology, Warsaw, Poland
`jerzy.rozanski@wat.edu.pl`

CGJTEAMLAB

The greater angle subtends the greater side

1 Introduction

Euclid's Proposition I.19 is the converse of Proposition I.18. In a nondegenerate triangle, the greater angle is opposite the greater side.

For a triangle ABC , if

$$\angle ACB < \angle ABC,$$

then

$$AB < AC.$$

The classical proof is indirect. One compares the two opposite sides AB and AC . Exactly one of the following three possibilities must occur:

$$AB \cong AC, \quad AB < AC, \quad AC < AB.$$

The first case contradicts Proposition I.5, because equal sides would imply congruent base angles. The third case contradicts Proposition I.18, because the greater side AB would force the opposite angle at C to be greater than the angle at B . Hence only

$$AB < AC$$

remains.

The current formal reconstruction makes this trichotomy explicit through the reusable general theorem

`hilbert_segment_trichotomy.`

It also uses the general irreflexivity theorem for strict angle comparison:

`hilbert_angleLess_irrefl.`

These two results are reusable structural components of the Hilbert layer rather than local devices specific to Proposition I.19.

2 The Classical Configuration

Let ABC be a nondegenerate triangle and suppose

$$\angle ACB < \angle ABC.$$




Figure 1: Classical and formal geometric configuration for Proposition I.19. The strict angle hypothesis $\angle ACB < \angle ABC$ and the target $AB < AC$ are stated in the text; they are not encoded by the apparent sizes of the drawn angles or sides.

The theorem asserts that the side opposite the larger angle must be larger:

$$AB < AC.$$

No new auxiliary point is required. The proof is entirely comparative: it uses the trichotomy of the two sides and eliminates the two alternatives incompatible with the assumed angle inequality.

3 Diagrammatic Audit

The diagrammatic audit retains exactly one geometric figure. Proposition I.19 itself constructs no new point, line, extension, intersection, betweenness relation, or auxiliary triangle. The three alternatives produced by segment trichotomy,

$$AB \cong AC, \quad AB < AC, \quad AC < AB,$$

are logical alternatives for the same fixed triangle ABC ; they are not three geometric states and therefore do not receive three panels.

The reusable theorem `hilbert_segment_trichotomy` has an internal construction involving a copied endpoint on a ray. That geometry belongs to the generic Hilbert comparison theorem, not to the proposition-specific proof of I.19, so it is documented textually rather than duplicated here. Likewise, `hilbert_angleLess_irrefl` is used only as a contradiction endpoint and introduces no new proposition-specific geometry.

Figure 19.1 therefore records only the common triangle. Neither the strict angle inequality nor the desired strict side inequality is inferred from the visual proportions of the drawing.

4 Classical Proof

Assume

$$\angle ACB < \angle ABC.$$

Compare the two sides AB and AC .

If

$$AB = AC,$$

then triangle ABC is isosceles. Proposition I.5 gives

$$\angle ABC = \angle ACB,$$

contradicting the hypothesis that

$$\angle ACB < \angle ABC.$$

If instead

$$AC < AB,$$

then Proposition I.18, applied to triangle ACB , yields

$$\angle ABC < \angle ACB,$$

which again contradicts the original hypothesis.

The only remaining possibility is

$$AB < AC.$$

Thus the greater angle is subtended by the greater side.

The classical dependency pattern is therefore

$$\begin{array}{c}
 \angle ACB < \angle ABC \\
 \downarrow \\
 AB \cong AC \vee AB < AC \vee AC < AB \\
 \downarrow \\
 \begin{array}{ccc}
 AB \cong AC & AB < AC & AC < AB \\
 \downarrow & & \downarrow \\
 \text{I.5 contradiction} & \text{desired case} & \text{I.18 contradiction}
 \end{array}
 \end{array}$$

5 Segment Trichotomy

The key reusable structural theorem called by the formal proof is

$$\text{hilbert_segment_trichotomy.}$$

Its conclusion has the form

$$AB \cong CD \vee AB < CD \vee CD < AB.$$

The proof does not introduce a new axiom. It is derived from segment construction and the order theory of collinear points.

A copy of AB is laid off on the ray CD , producing a point X such that

$$CX \cong AB.$$

If $X = D$, then

$$AB \cong CD.$$

Otherwise, C, D, X are distinct collinear points. The Hilbert betweenness trichotomy gives three possibilities. The case

$$D - C - X$$

is incompatible with the fact that X lies on the ray CD . The remaining two orders give respectively

$$CD < CX$$

and

$$CX < CD.$$

Transporting these inequalities through

$$CX \cong AB$$

yields the required segment trichotomy.

Thus the theorem packages a general completeness property of strict segment comparison.

6 Irreflexivity of Strict Angle Comparison

The second structural theorem used in I.19 is

$$\text{hilbert_angleLess_irrefl.}$$

It states

$$\neg(\alpha < \alpha).$$

In the synthetic definition of `HilbertAngleLess`, the relation

$$\alpha < \beta$$

means that a congruent copy of α occurs as a proper interior subangle of β .

Therefore

$$\alpha < \alpha$$

would imply that an angle is congruent to one of its own proper interior subangles. This is excluded by the previously proved theorem

$$\text{hilbert_interior_subangle_not_congruent_whole.}$$

The irreflexivity result provides exactly the contradiction mechanism needed in both non-target branches of Proposition I.19.

7 Proof Architecture of the Reconstruction

The Lean proof divides naturally into three stages.

7.1 Nondegeneracy of the Compared Side

From the noncollinearity hypothesis

$$\neg \text{Collinear}(A, B, C)$$

the proof first derives

$$A \neq C.$$

This is the nondegeneracy condition required by the comparison theorem `hilbert_segment_trichotomy`.

7.2 Apply Segment Trichotomy

The theorem

$$\text{hilbert_segment_trichotomy}$$

is applied to the segments AB and AC .

This produces three branches:

$$AB \cong AC, \quad AB < AC, \quad AC < AB.$$

The middle branch is exactly the desired conclusion.

7.3 Eliminate the Two Impossible Branches

In the congruent-side branch, Proposition I.5 gives

$$\angle ABC \cong \angle ACB.$$

The assumed strict inequality

$$\angle ACB < \angle ABC$$

is transported across that congruence, producing

$$\angle ACB < \angle ACB.$$

This contradicts `hilbert_angleLess_irrefl`.

In the branch

$$AC < AB,$$

Proposition I.18 applied to triangle ACB gives

$$\angle ABC < \angle ACB.$$

Combining this with the original inequality by transitivity yields

$$\angle ACB < \angle ACB,$$

again contradicting irreflexivity.

Only

$$AB < AC$$

remains.

8 Lean Formalization

The current production module imports `CGJteamLab.Proposition18`. The public theorem is:

```
theorem euclid_proposition_19
  [HilbertIncidence Geo]
  [HilbertCongruence Geo]
  (A B C : Geo.Point)
  (hABC : not PrimCollinear Geo A B C)
  (hAngle :
    HilbertAngleLess Geo A C B A B C) :
  HilbertSegmentLess Geo A B A C := by
```

The proof first derives the nondegeneracy of AC :

```
have hAC : A != C :=
  hilbert_noncollinear_ne_first
  Geo A C B
  (by
    intro h
    exact hABC
    (PrimCollinearRotate Geo A C B h))
```

Then segment trichotomy is applied:

```
rcases
  hilbert_segment_trichotomy
  Geo
  A B
  A C
  hAC with
  hEq | hABltAC | hACltAB
```

The middle branch closes immediately:

```
exact hABltAC
```

For the congruent-side branch, the generic isosceles-base-angle theorem (the formal counterpart of the classical I.5 step) gives:

```
have hIso :
  Geo.AngleCongruent A B C A C B :=
  hilbert_isosceles_base_angles
```

```

Geo
A B C
hABC
hEq

```

The strict inequality is transported to the congruent target angle:

```

have hSame :
  HilbertAngleLess Geo A C B A C B :=
  hilbert_angleLess_transport_right
  Geo
  A C B
  A B C
  A C B
  hAngle
  hACB
  hIso

```

This contradicts irreflexivity:

```

exact
  False.elim
    ((hilbert_angleLess_irrefl
      Geo A C B)
     hSame)

```

In the final branch, Proposition I.18 gives the opposite strict inequality:

```

have hOpp :
  HilbertAngleLess Geo A B C A C B :=
  euclid_proposition_18
  Geo
  A C B
  hACB
  hACltAB

```

Transitivity then creates an impossible strict self-comparison:

```

have hCycle :
  HilbertAngleLess Geo A C B A C B :=
  hilbert_angleLess_trans
  Geo
  A C B
  A B C
  A C B
  hAngle
  hOpp

```

and finally:

```

exact
  False.elim
    ((hilbert_angleLess_irrefl
      Geo A C B)
     hCycle)

```

9 Relationship with Earlier Propositions

9.1 Proposition I.5 and the formal isosceles theorem

Classically, I.5 eliminates the equality case: if

$$AB \cong AC,$$

then the base angles are congruent. In the current Lean source, however, `euclid_proposition_5` is not called directly. The proof uses the reusable theorem

`hilbert_isosceles_base_angles`

to obtain

$$\angle ABC \cong \angle ACB.$$

Thus I.5 belongs to the classical dependency DAG, while `hilbert_isosceles_base_angles` is the direct formal dependency for this branch.

9.2 Proposition I.18

I.18 is used to eliminate the reversed side inequality. If

$$AC < AB,$$

then applying I.18 to triangle ACB gives

$$\angle ABC < \angle ACB.$$

Together with the hypothesis

$$\angle ACB < \angle ABC,$$

this produces a strict cycle and therefore a contradiction.

9.3 Logical Relation between I.18 and I.19

The two propositions form a converse pair:

$$AB < AC \implies \angle ACB < \angle ABC$$

for I.18, and

$$\angle ACB < \angle ABC \implies AB < AC$$

for I.19.

Together they establish the equivalence between ordering of opposite sides and ordering of opposite angles in a nondegenerate triangle.

10 Hilbert Components Used

10.1 `hilbert_segment_trichotomy`

This theorem provides the complete three-way comparison of two segments and is the main reusable structural component called by I.19.

10.2 hilbert_angleLess_irrefl

This theorem excludes strict self-comparison of an angle. It is used in both contradictory branches of the proof.

10.3 hilbert_isosceles_base_angles

This is the formal version of Proposition I.5 needed to eliminate the equality case.

10.4 hilbert_angleLess_transport_right

This theorem transports a strict angle comparison across congruence of the containing angle. It converts the equality branch into an impossible self-comparison.

10.5 hilbert_angleLess_trans


Transitivity combines the original strict inequality with the opposite inequality supplied by I.18.

10.6 Direct proposition-level dependency

The only imported Book-I proposition used directly by `euclid_proposition_19` is `euclid_proposition_18`, in the branch $AC < AB$. The equality branch is closed by generic Hilbert infrastructure rather than a direct call to `euclid_proposition_5`.

11 Dependency Summary

The formal dependency structure is



No local axiom declaration occurs in the current `Proposition19.lean`; this diagrammatic audit did not run a fresh axiomatic dependency check.

12 Conclusion

Proposition I.19 is formally shorter than I.18, but it exposes an important structural issue: the proof requires a complete comparison principle for segments.

Once the theorem `hilbert_segment_trichotomy` is available, the Euclidean argument becomes exactly the expected three-case proof. Equal sides are ruled out by I.5, the reversed inequality is ruled out by I.18, and the remaining case is the desired conclusion.

The proof also clarifies the role of strict angle order. Instead of introducing an explicit asymmetry theorem, the development uses the more fundamental statement `hilbert_angleLess_irrefl` together with transitivity. This keeps the order theory modular and avoids unnecessary duplication.

Thus the current I.19 proof relies on two reusable Hilbert-layer principles: completeness of segment comparison and irreflexivity of strict angle comparison. The Euclidean theorem itself becomes a concise application of those general principles.

The preserved Book1R document recorded a clean Lean development with no `sorry`. In the present diagrammatic audit the production Lean file was read but not rebuilt, and no fresh `#print axioms` check was run.