

Non-backtracking Random Walks

Cory Glover

Brigham Young University
Department of Mathematics

Joint work with
Mark Kempton, Brigham Young University
Tyler Jones, Brigham Young University

SCR 2020
Provo, UT
February 29, 2020

Is the mixing rate of a non-backtracking random walk faster than that of a simple random walk on a graph?

Definition

Let P be the transition probability matrix of a graph G and let π be the stationary distribution of a random walk on G . Then the **mixing rate** of a random walk on a graph G is defined pointwise by

$$\rho = \limsup_{t \rightarrow \infty} \max_{u,v} |P^t(u, v) - \pi(v)|^{1/t}.$$

Mixing Rate and Eigenvalues

Theorem (Lovasz 1993)

Let G be a connected non-bipartite graph with transition probability matrix P . Let $\mu_1 \geq \mu_2 \geq \dots \geq \mu_n$ be the eigenvalues of P . Then the mixing rate of G is $\rho = \max\{|\mu_2|, |\mu_n|\}$.

Hashimoto Matrix

Definition

The **Hashimoto Matrix** of a graph G is

$$B((u, v), (x, y)) = \begin{cases} 1 & v = x \text{ and } u \neq y \\ 0 & \text{otherwise} \end{cases} \quad (1)$$

Ihara's Theorem

Theorem (Ihara)

Given a graph G with n vertices and m edges, define B to be the Hashimoto matrix of G . Let A be the adjacency matrix of G and let D be the diagonal degree matrix. Then

$$\det(I - \mu B) = (1 - \mu^2)^{m-n} \det(I - \mu A + \mu^2(D - I)).$$

Thus, the eigenvalues of B are μ .

The Matrix K

From Krzakala et. al. (2013), the matrix K of a graph G

$$K = \begin{pmatrix} A & D - I \\ -I & \mathbf{0} \end{pmatrix}$$

is a invariant subspace of B .

The Matrix K

Define the following matrices:

$$S((u, v), x) = \begin{cases} 1 & v = x \\ 0 & \text{otherwise} \end{cases}$$

$$T(x, (u, v)) = \begin{cases} 1 & u = x \\ 0 & \text{otherwise} \end{cases}.$$

Then

$$B \begin{pmatrix} S & T^T \end{pmatrix} v = \begin{pmatrix} S & T^T \end{pmatrix} K v.$$

If v is an eigenvector of K with eigenvalue μ , then

$$B \begin{pmatrix} S & T^T \end{pmatrix} v = \mu \begin{pmatrix} S & T^T \end{pmatrix} v.$$

Eigenvalues of K

If $(x \ y)^T$ is an eigenvector of K ,

$$\begin{pmatrix} A & D - I \\ -I & \mathbf{0} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \mu \begin{pmatrix} x \\ y \end{pmatrix},$$

then $x = -\mu y$. This means that $\mu^2 y - \mu A y + (D - I)y = \mathbf{0}$ for all $\mu \in \sigma(K)$.

Ihara's Theorem

Theorem (Ihara, 1966)

Given a graph G and the associated Hasimoto matrix B , adjacency matrix A , and diagonal degree matrix D , the following is true:

$$\det(I - \mu B) = (1 - u^2)^{m-n} \det(I - uA - u^2(D - I)).$$

So the eigenvalues of B are the $\frac{1}{u}$.

Relationship Between A and K

Let \mathbf{x} be the eigenvector associated with λ_2 (the second largest eigenvalue) of A . Let μ_2 be the second largest eigenvalue of K such that $K \begin{pmatrix} -\mu_2 \mathbf{y} & \mathbf{y} \end{pmatrix}^T = \mu_2 \begin{pmatrix} -\mu_2 \mathbf{y} & \mathbf{y} \end{pmatrix}^T$. Scale \mathbf{x} such that $\mathbf{x}^T \mathbf{y} = 1$.

Then,

$$\begin{aligned}\mu_2^2 \mathbf{x}^T \mathbf{y} - \mu_2 \mathbf{x}^T A \mathbf{y} + \mathbf{x}^T (D - I) \mathbf{y} &= 0, \\ \mu_2^2 - \mu_2 \lambda_2 + \mathbf{x}^T (D - I) \mathbf{y} &= 0.\end{aligned}$$

Solving for μ_2 ,

$$\mu_2 = \frac{\lambda_2 \pm \sqrt{\lambda_2^2 - 4 \mathbf{x}^T (D - I) \mathbf{y}}}{2}.$$

Questions That Arise

$$\mu = \frac{\lambda \pm \sqrt{\lambda^2 - 4\mathbf{x}^T(D - I)\mathbf{y}}}{2}$$

- How are μ and λ related?
- Is it $+$ or $-$?
- Can we get all eigenvalues μ ?
- How to convert μ to ρ ?

Relating μ and λ

Assume that $\lambda \geq 2\sqrt{\mathbf{x}^T(D-I)\mathbf{y}}$:

$$\mu = \frac{\lambda + \sqrt{\lambda^2 - 4\mathbf{x}^T(D-I)\mathbf{y}}}{2} \leq \frac{\lambda + \lambda}{2} = \lambda$$

$$\mu = \frac{\lambda - \sqrt{\lambda^2 - 4\mathbf{x}^T(D-I)\mathbf{y}}}{2} \leq \frac{\lambda}{2}.$$

In either case $\mu \leq \lambda$.

INSERT PICTURE OF BIPARTITE GRAPH HERE

Theorem

The following are equivalent for a graph G :

- 1 G is bipartite,
- 2 The spectrum of B is symmetric,
- 3 The spectrum of K is symmetric,
- 4 -1 is an eigenvalue of K .

Bipartite $\Rightarrow \sigma(B)$ Is Symmetric

If G is bipartite, then

$$B = \begin{pmatrix} 0 & B_2 \\ B_1 & 0 \end{pmatrix}.$$

Let $\lambda \in \sigma(B)$. Then,

$$B \begin{pmatrix} x & y \end{pmatrix}^T = \lambda \begin{pmatrix} x & y \end{pmatrix}^T \quad (2)$$

$$\begin{pmatrix} B_2 y & B_1 x \end{pmatrix}^T = \lambda \begin{pmatrix} x & y \end{pmatrix}^T. \quad (3)$$

This means that

$$B \begin{pmatrix} x & -y \end{pmatrix}^T = \begin{pmatrix} -B_2 y & B_1 x \end{pmatrix}^T \quad (4)$$

$$= \begin{pmatrix} -\lambda x & \lambda y \end{pmatrix}^T \quad (5)$$

$$= -\lambda \begin{pmatrix} x & -y \end{pmatrix}^T. \quad (6)$$

So $-\lambda \in \sigma(B)$.

$\sigma(B)$ Is Symmetric $\Rightarrow \sigma(K)$ Is Symmetric

Theorem (Ihara, 1966)

Given a graph G and the associated Hasimoto matrix B , adjacency matrix A , and diagonal degree matrix D , the following is true:

$$\det(I - \mu B) = (1 - u^2)^{m-n} \det(I - uA - u^2(D - I)).$$

$\sigma(K)$ is symmetric $\Rightarrow \sigma(B)$ is symmetric.

Theorem (Ihara, 1966)

Given a graph G and the associated Hasimoto matrix B , adjacency matrix A , and diagonal degree matrix D , the following is true:

$$\det(I - \mu B) = (1 - u^2)^{m-n} \det(I - uA - u^2(D - I)).$$

$\sigma(B)$ is symmetric $\Rightarrow G$ is bipartite

Consider B as the adjacency matrix of its own graph.

Paths: Cospectrality and Fractional Cospectrality

When can two vertices of a path be cospectral or fractionally cospectral?

Paths: Cospectrality and Fractional Cospectrality

When can two vertices of a path be cospectral or fractionally cospectral?

Lemma

Let u, v be vertices of a path on n vertices P_n

- u and v are cospectral if and only if $\{u, v\} = \{i, n - i\}$.*


Paths: Cospectrality and Fractional Cospectrality

When can two vertices of a path be cospectral or fractionally cospectral?

Lemma

Let u, v be vertices of a path on n vertices P_n

- u and v are cospectral if and only if $\{u, v\} = \{i, n - i\}$.*
- u and v are fractionally cospectral (and not cospectral) if and only if $n = 5d - 1$ for some positive integer d , and $\{u, v\} = \{d, 3d\}$ or (by symmetry) $\{u, v\} = \{2d, 4d\}$.*



Fractional Cospectrality in Paths

We prove this by counting walks in paths

$$A^{2d+2j}(u, u) = \binom{2d+2j}{d+j} - \binom{2d+2j}{j}$$

$$A^{2d+2j}(v, v) = \binom{2d+2j}{d+j}$$

$$A^{2d+2j}(u, v) = \binom{2d+2j}{j}.$$

Paths: Eigenvalue Condition

The eigenvalues of P_{5d-1} are

$$\lambda_j = 2 \cos \frac{\pi j}{5d}, \quad j = 1, \dots, 5d - 1$$

Paths: Eigenvalue Condition

The eigenvalues of P_{5d-1} are

$$\lambda_j = 2 \cos \frac{\pi j}{5d}, \quad j = 1, \dots, 5d - 1$$

Lemma

Let m be an odd integer, $0 \leq a < k$ integers. Then

$$\sum_{i=0}^{m-1} (-1)^i \cos \left(\frac{(a + ik)\pi}{km} \right) = 0.$$

Paths: Eigenvalue Condition

The eigenvalues of P_{5d-1} are

$$\lambda_j = 2 \cos \frac{\pi j}{5d}, \quad j = 1, \dots, 5d - 1$$

Lemma

Let m be an odd integer, $0 \leq a < k$ integers. Then

$$\sum_{i=0}^{m-1} (-1)^i \cos \left(\frac{(a + ik)\pi}{km} \right) = 0.$$

Eigenvalue condition satisfied if and only if $d = 2^k$ for some k .

Theorem

Vertices u and v of P_n exhibit PGFR if and only if we are in one of the following cases:

- *There is PGST from u to v (characterized previously).*
- *$n = 5 \cdot 2^k - 1$ and $\{u, v\} = \{2^k, 3 \cdot 2^k\}$ or $\{2 \cdot 2^k, 4 \cdot 2^k\}$.*

- What other graphs have fractionally cospectral pairs of nodes?
Which of these exhibit FR/PGFR?

Future Work

- What other graphs have fractionally cospectral pairs of nodes?
Which of these exhibit FR/PGFR?
- Weighted paths?

- What other graphs have fractionally cospectral pairs of nodes?
Which of these exhibit FR/PGFR?
- Weighted paths?
- FR or PGFR among more than 2 nodes in paths?

Thank You!