

Non-backtracking Random Walks

Cory Glover

Brigham Young University
Department of Mathematics

Joint work with
Mark Kempton, Brigham Young University
Tyler Jones, Brigham Young University


SCR 2020
Provo, UT
February 29, 2020

Problem

- When the Pagerank vector of a non-backtracking random walk give the same values as a simple random walk?
- Can the Pagerank vector of a non-backtracking random walk be calculated faster than a simple random walk?

Definition (Hashimoto Matrix)

$$B((u, v), (x, y)) = \begin{cases} 1 & v = x, u \neq y \\ 0 & \text{otherwise} \end{cases}$$



Theorem (Arrigo et. al., 2018)

Let A be the adjacency matrix of an undirected k -regular graph, with $k \geq 2$. For any $\epsilon \in [0, 1]$, define

$$\hat{P} = \epsilon \hat{D}^{-1} B + \frac{1 - \epsilon}{n} \mathbf{1} \mathbf{v}^T$$

and $P = \epsilon D^{-1} A + \frac{1 - \epsilon}{n} \mathbf{1} \mathbf{1}^T$, where $\mathbf{v} = T^T D^{-1} \mathbf{1}$. If $\hat{P}^T \hat{y} = \hat{y}$ and $P^T x = x$, with $\|x\|_1 = \|\hat{y}\|_1 = 1$, then

$$T\hat{y} = x.$$

Theorem

Let A be the adjacency matrix of a bipartite biregular graph, with $d_1, d_2 \geq 2$. For any $\epsilon \in [0, 1]$, define $\hat{P} = \epsilon \hat{D}^{-1} B + \frac{1-\epsilon}{n} \mathbf{1} \mathbf{v}^T$ and $P = \epsilon D^{-1} A + \frac{1-\epsilon}{n} \mathbf{1} \mathbf{1}^T$, where $\mathbf{v} = T^T D^{-1} \mathbf{1}$. If $\hat{P}^T \hat{y} = \hat{y}$ and $P^T x = x$, with $\|x\|_1 = \|\hat{y}\|_1 = 1$, then

$$T\hat{y} = x.$$

Estimating Pagerank vs Nonbacktracking Pagerank

Conjecture

In most cases, NBRW mix faster than simple random walks.

Alon et. al. (2007) showed that in most cases, NBRW mix faster on k -regular graphs, with $k \geq 2$.

Theorem (Lovasz 1993)

Let G be a connected non-bipartite graph with transition probability matrix P . Let $\mu_1 \geq \mu_2 \geq \dots \geq \mu_n$ be the eigenvalues of P . Then the mixing rate of G is $\rho = \max\{|\lambda_2|, |\lambda_n|\}$.

Ihara's Theorem

Theorem (Ihara)

Given a graph G with n vertices and m edges, define B to be the Hashimoto matrix of G . Let A be the adjacency matrix of G and let D be the diagonal degree matrix. Then

$$\det(I - uB) = (1 - u^2)^{m-n} \det(I - uA + u^2(D - I)).$$

Thus, the eigenvalues of B are $\mu = \frac{1}{u}$.

Comparing Eigenvalue

From Krzakala et. al. (2013), the matrix K of a graph G

$$K = \begin{pmatrix} A & D - I \\ -I & \mathbf{0} \end{pmatrix}$$

is a invariant subspace of B .

Eigenvalues of K

$$\begin{pmatrix} A & D - I \\ -I & \mathbf{0} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \mu \begin{pmatrix} x \\ y \end{pmatrix},$$

$$x = -\mu y$$

$$\mu^2 y - \mu A y + (D - I)y = \mathbf{0}$$

for all $\mu \in \sigma(K)$.

Relationship Between A and K

Let \mathbf{x} be the eigenvector associated with $\lambda \in \sigma(A)$.

Then,

$$\begin{aligned}\mu^2 \mathbf{x}^T \mathbf{y} - \mu \mathbf{x}^T A \mathbf{y} + \mathbf{x}^T (D - I) \mathbf{y} &= 0, \\ \mu^2 - \mu \lambda + \mathbf{x}^T (D - I) \mathbf{y} &= 0.\end{aligned}$$

Solving for μ ,

$$\mu = \frac{\lambda \pm \sqrt{\lambda^2 - 4 \mathbf{x}^T (D - I) \mathbf{y}}}{2}.$$


Open Questions

$$\mu = \frac{\lambda \pm \sqrt{\lambda^2 - 4\mathbf{x}^T(D - I)\mathbf{y}}}{2}$$

- How are μ and λ related?
- Is it $+$ or $-$?
- Can we get all eigenvalues μ ?
- How to convert μ to eigenvalue of P ?


Current Pagerank Algorithms

- Borgs et. al. (2016) created algorithm to solve significant Pagerank problem, i.e. find all nodes with Pagerank value greater than Δ , in $O(\frac{n}{\Delta})$
- Bahmini, Chowdhury, and Goel (2010) created algorithm to estimate personalized pagerank and global pagerank in $O(\frac{n \ln(m)}{\epsilon^2})$.




Current: [2]

Paths: []




Current: $[2, 1]$

Paths: $[]$




Current: $[2, 1, 3]$

Paths: $[]$



Current: [2]
Paths: [[2, 1, 3]]




Current: [1]
Paths: [[2, 1, 3], [2, 4]]


$$\tilde{\pi}_v = \frac{X_v}{nR/\epsilon}$$

where X_v is the number of paths containing v .

Computational Results




(a) Simple Example




(b) Karate

Computational Results

$$y = Cx^{-\alpha}$$




(a) Simple Example




(b) Karate

Computational Results



(a) SRW: $\alpha = .54$, NBRW: $\alpha = .56$



(b) SRW: $\alpha = .49$, NBRW: $\alpha = .52$

Figure: Simple Example: $|C - C_{NB}| = .034$
Karate: $|C - C_{NB}| = .0052$

Thank you!