Caleb McWhorter — *Solutions* "I like long walks, especially when they **MATH 108** Fall 2021

are taken by people that annoy me." HW 7: Due 11/04

-Fred Allen

Problem 1. (10pt) Write down the tableau associated to the following linear programming problem:

$$\max z = 3x_1 + x_2$$

$$2x_1 + 3x_2 \le 6$$

$$4x_1 + 2x_2 \le 8$$

$$-3x_1 + 4x_2 \ge 2$$

$$x_1, x_2 \ge 0$$

Solution. Before creating the tableau, we want to be sure the linear programming problem is in standard form (or as close as possible). Because this is a maximization problem, we require all the inequalities to be '\le \cdot' [This is excluding the final inequality $x_1, x_2 \ge 0$, because we require the variables to be nonnegative.] So we multiply both sides of the third inequality by -1 to obtain $3x_1 - 4x_2 \le -2$. Then our system is

$$\max z = 3x_1 + x_2$$

$$2x_1 + 3x_2 \le 6$$

$$4x_1 + 2x_2 \le 8$$

$$3x_1 - 4x_2 \le -2$$

$$x_1, x_2 \ge 0$$

We introduce slack variables s_1, s_2, s_3 so that...

$$2x_1 + 3x_2 + s_1 = 6$$

 $4x_1 + 2x_2 + s_2 = 8$
 $3x_1 - 4x_2 + s_3 = -2$

Moving everything to the left side in $z = 3x_1 + x_2$, we have $z - 3x_1 - x_2 = 0$. Then the associated tableau is...

Problem 2. (10pt) Assume the following is a tableau associated to a standard maximization problem. Write down the function being maximization and the system of constraints.

Solution. It is clear that we had three inequalities. Therefore, we had three slack variables, s_1, s_2, s_3 . Because there are seven columns, one of which must correspond to the b's, there must be three variables, x_1, x_2, x_3 . Looking at the last row and assuming we are trying to maximize a variable z, we must have $z - 5x_1 - 4x_2 - 5x_3 = 0$. But then $z = 5x_1 + 4x_2 + 5x_3$. Therefore, the standard maximization problem was...

$$\max z = 5x_1 + 4x_2 + 5x_3$$

$$x_1 + 2x_2 + x_3 \le 100$$

$$2x_1 + 8x_2 + 2x_3 \le 150$$

$$x_1 + x_2 + x_3 \le 200$$

$$x_1, x_2, x_3 \ge 0$$

Problem 3. (10pt) Solve the following linear programming problem:

$$\max z = x_1 + 6x_2 + 3x_3$$
$$x_1 + x_2 + 2x_3 \le 4$$
$$x_1 + 2x_2 + x_3 \le 4$$
$$x_1, x_2, x_3 \ge 0$$

Solution. This linear programming problem is already in standard form for a maximization. We introduce slack variables s_1 and s_2 so that...

$$x_1 + x_2 + 2x_3 + s_1 = 4$$

 $x_1 + 2x_2 + x_3 + s_2 = 4$

Moving everything to the left side in $z = x_1 + 6x_2 + 3x_3$, we have $z - x_1 - 6x_2 - 3x_3 = 0$. Then the associated tableau is...

We find our first pivot position:

So x_x is the entering variable and s_2 is the exiting variable. We make this pivot entry 1 by $\frac{1}{2}R_2 \to R_2$.

Now we perform the first step of the simplex method: $-R_2 + R_1 \rightarrow R_1$ and $6R_2 + R_3 \rightarrow R_3$.

Because there are no negatives in the bottom row, the simplex method is complete. We see that the maximum value for z is 12 (the bottom-rightmost entry) and occurs at $(x_1, x_2, x_3, s_1, s_2) = (0, 2, 0, 2, 0)$, i.e. $x_1 = 0, x_2 = 2, x_3 = 0, s_1 = 2, s_2 = 0$.

Problem 4. (10pt) Find the dual problem to...

$$\min z = 5x_1 + 4x_2$$

$$x_1 + 7x_2 \ge 7$$

$$x_1 + 3x_2 \ge 9$$

$$x_1, x_2 \ge 0$$

Solution. First, observe that this linear programming problem is a minimization and is already in standard form. Therefore, the associated matrix is...

$$\begin{pmatrix} 1 & 7 & 7 \\ 1 & 3 & 9 \\ 5 & 4 & 0 \end{pmatrix}$$

Taking the transpose, we find...

$$\begin{pmatrix} 1 & 1 & 5 \\ 7 & 3 & 4 \\ 7 & 9 & 0 \end{pmatrix}$$

Therefore, the dual problem is...

$$\max w = 7y_1 + 9y_2$$
$$y_1 + y_2 \le 5$$
$$7y_1 + 3y_2 \le 4$$
$$y_1, y_2 \ge 0$$

Problem 5. (10pt) Solve the following linear programming problem:

$$\min z = 2x_1 + 3x_2$$

$$2x_1 + x_2 \ge 1$$

$$x_1 + 3x_2 \ge 1$$

$$x_1, x_2 \ge 0$$

Solution. This is linear programming problem is a minimization and it is already in standard form. Therefore, the associated matrix is...

$$\begin{pmatrix} 2 & 1 & 1 \\ 1 & 3 & 1 \\ 2 & 3 & 0 \end{pmatrix}$$

The transpose is then...

$$\begin{pmatrix} 2 & 1 & 2 \\ 1 & 3 & 3 \\ 1 & 1 & 0 \end{pmatrix}$$

Therefore, the associated maximization problem (the dual problem) is...

$$\max w = y_1 + y_2$$
$$2y_1 + y_2 \le 2$$
$$y_1 + 3y_2 \le 3$$
$$y_1, y_2 \ge 0$$

Because $w = y_1 + y_2$, we know that $w - y_1 - y_2 = 0$. We introduce slack variables s_1, s_2 so that...

$$2y_1 + y_2 + s_1 = 2$$
$$y_1 + 3y_2 + s_2 = 3$$

Therefore, the associated tableau is...

We find our first pivot position:

So y_1 is the entering variable and s_1 is the exiting variable. We make this pivot entry 1 by $\frac{1}{2}R_1 \to R_1$.

Now we perform the first step of the simplex method: $-R_1 + R_2 \rightarrow R_2$ and $R_1 + R_3 \rightarrow R_3$.

There are still negatives in the last row. So we proceed with the next step of the simplex method. We find our pivot position:

So y_2 is the entering variable and s_2 is the exiting variable. We make the pivot position 1 by $\frac{2}{5}R_2 \to R_2$.

Now we perform the next step of the simplex method: $-\frac{1}{2}R_2 + R_1 \to R_1$ and $\frac{1}{2}R_2 + R_3 \to R_3$:

Because there are no remaining negative entries in the bottom row, the simplex method is complete. We have maximum value $w=\frac{7}{5}=1.4$ (the bottom-rightmost entry) occurring at $(y_1,y_2,s_1,s_2)=(\frac{3}{5},\frac{4}{5},0,0)=(0.6,0.8,0,0)$, i.e. $y_1=\frac{3}{5}=0.6,y_2=\frac{4}{5}=0.8,s_1=0,s_2=0$.

But this maximum value is the minimum value for our original minimization problem (the dual problem). Therefore, the minimum value is $z=\frac{7}{5}=1.4$ occurring at $(x_1,x_2,s_1,s_2)=(\frac{2}{5},\frac{1}{5},0,0)=(0.4,0.2,0,0)$, i.e. $x_1=\frac{2}{5}=0.4, x_2=\frac{1}{5}=0.2, s_1=0, s_2=0$