"I can be just as non-competitive as anybody. Matter of fact, I'm the most non-competitive, so I win." —Peter Griffin, Family Guy

Problem 1. (10pt) Showing all the steps according to order of operations, compute the following:

(a)
$$10 + 10 - 16 \cdot 0 + 2 + 2$$

(b)
$$(-1)^3 - 1 + 4^2/2$$

(c)
$$15 - (6 - 10) + 3^2$$

(d)
$$\frac{-4 - (2 - 4)^2}{3^2 - 1}$$

Problem 2. (10pt) Define the following sets:

$$A = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$$

$$B = \{2, 4, 6, 8, 10\}$$

$$C = \{1, 3, 5, 7, 9\}$$

$$D = \{2, 3, 5, 7\}E$$

$$= \{2, 3, 4, 6, 8, 9\}$$

Consider all these sets as subsets of A. Compute the following:

- (a) B^c
- (b) $B \cup D$
- (c) $E \setminus D$
- (d) $C \cap E$
- (e) |A|

Problem 3. (10pt) Define the following sets:

A = All males over 40 years old.

B = All people that have acted in a movie.

 $C=\mathsf{All}\ \mathsf{US}\ \mathsf{Presidents},$ alive or dead.


D = All persons under 6 ft tall.

Consider all of these sets as subsets of the set of all people alive. Being sure to completely justify your response, answer the following:

- (a) Find an element of $A \cap B$.
- (b) Is Jeff Bezos $\in A \cup C$? Is Jeff Bezos $\in C \cup D$?
- (c) Is George Washington $\in C B$?
- (d) Is Danny Devito $\in D^C$?
- (e) Are sets *B* and *C* disjoint? [Hint: Consider US Presidents from the last 50 years.]

Problem 4. (10pt) Determine whether the following relations are functions, being sure to justify your answer. If the relation is a function, determine its domain, codomain, and range. [For this problem, in determining a functions domain, codomain, and range, you may invoke the use/description of a graph.]

(a)



(b)

x	g(x)		
1.0	1.0		
1.5	4.3		
3.0	-6.1		
4.4	2.2		
6.8	1.0		

(c)
$$h(x,y) = x + y^4$$
.

(d) j(x) =the multiple of two closest to x.

Problem 5. (10pt) Suppose that f(x, y) is the function given by the following table:

$x \setminus y$	1	2	3	4
1	-2	7	4	-4
2	0	3	-1	1
3	5	-6	7	6
4	1	0	4	0

Showing all your work, compute the following:

- (a) f(3,2)
- (b) $f(3-1,2^2)$
- (c) 5f(3,1) 8

(d)
$$\frac{4 - f(3^2 + (-2)^3, 1)}{2f(1, 3)}$$

Problem 6. (10pt) Let rdwn(x) denote the largest integer that is *less than* x.

- (a) Find rdwn(x) for x = 0.5, 2.2, 5.9, 6.0, -1.5, -4.9, -7.
- (b) Explain why rdwn(x) is a function.
- (c) Being as accurate as possible, sketch a graph of rdwn(x) on the plot below.

