

Check-In 08/21. (*True/False*) The integral $\int x \sqrt[3]{x-2} dx$ can be treated as a ‘shifting integral’ by using the u -substitution $u = x - 2$.

Solution. The statement is *true*. We ‘want’ to be able to distribute the x across the cube-root but we cannot—this is not a valid operation. However, if we make the u -substitution $u = x - 2$, then we will be able to distribute in a way that makes this integral ‘routine.’ So, let $u = x - 2$, then $du = dx$. Moreover, because $u = x - 2$, we know that $x = u + 2$. But then...

$$\int x \sqrt[3]{x-2} dx = \int (u+2) \sqrt[3]{u} du = \int (u^{4/3} + 2u^{1/3}) du = \frac{3}{7} u^{7/3} + \frac{3}{4} \cdot 2u^{4/3} + C = \frac{3}{7} (x-2)^{7/3} + \frac{3}{2} (x-2)^{4/3} + C$$

Note that a computer algebra system may write the answer (though you will *not* be expected to) like this:

$$\frac{3}{7} (x-2)^{7/3} + \frac{3}{2} (x-2)^{4/3} + C = (x-2)^{4/3} \left(\frac{3}{7} (x-2) + \frac{3}{2} \right) + C = (x-2)^{4/3} \left(\frac{3}{7} x + \frac{9}{14} \right) + C = \frac{3}{14} (x-2)^{4/3} (2x+3) + C$$

Check-In 08/26. (*True/False*) Using integration-by-parts to evaluate $\int x \tan^{-1}(x) dx$, one chooses $u = \tan^{-1} x$ and $dv = x$.

Solution. The statement is *true*. Using LIATE, the first term that appears is ‘T’ for inverse trig. Therefore, we choose $u = \tan^{-1} x$. But then $dv = x$. We then fill in our box:

$\tan^{-1} x$	$\frac{x^2}{2}$
$\frac{1}{1+x^2}$	x

Using the ‘rule of 7’, we have...

$$\int x \tan^{-1} x dx = \frac{1}{2} x^2 \tan^{-1} x - \frac{1}{2} \int \frac{x^2}{1+x^2} dx$$

We now need only evaluate the integral on the right. Dividing $1+x^2$ into x^2 , we have a remainder of -1 , i.e. $\frac{x^2}{1+x^2} = 1 + \frac{-1}{1+x^2}$. Therefore, we have...

$$\frac{1}{2} \int \frac{x^2}{1+x^2} dx = \frac{1}{2} \int \left(1 + \frac{-1}{1+x^2} \right) dx = \frac{1}{2} (x - \tan^{-1} x) + C$$

But then...

$$\begin{aligned}
 \int x \tan^{-1} x \, dx &= \frac{1}{2} x^2 \tan^{-1} x - \frac{1}{2} \int \frac{x^2}{1+x^2} \, dx \\
 &= \frac{1}{2} x^2 \tan^{-1} x - \frac{1}{2} (x - \tan^{-1} x) + C \\
 &= \frac{1}{2} x^2 \tan^{-1} x - \frac{1}{2} x + \frac{1}{2} \tan^{-1} x + C \\
 &= \frac{x^2 \tan^{-1} x - x + \tan^{-1} x}{2} + C \\
 &= \frac{(x^2 + 1) \tan^{-1} x - x}{2} + C
 \end{aligned}$$

Check-In 08/28. (*True/False*) The integral $\int e^x \sin(3x) \, dx$ can be treated as an integration-by-parts ‘looping’ integral.

Solution. The statement is *true*. Using integration-by-parts for $\int e^x \sin(3x) \, dx$ would result in an integral that would ‘loop’ back to itself. Generally, an integrand of the form exponential · (sin or cos) or trig · trig will have this property. Using traditional integration-by-parts, by LIATE, we choose $u = \sin(3x)$ and $dv = e^x$. Filling out our box, we have...

$\sin(3x)$	e^x
$3 \cos(3x)$	e^x

Using the ‘rule of seven’, we then have...

$$\int e^x \sin(3x) \, dx = e^x \sin(3x) - \int 3e^x \cos(3x) \, dx$$

To integrate $\int 3e^x \cos(3x) \, dx$, we again use integration-by-parts. Using LIATE, we choose $u =$

$3 \cos(3x)$ and $dv = e^x$. Filling out the box, we have...

$3 \cos(3x)$	e^x
$-9 \sin(3x)$	e^x

Using the 'rule of seven', we then have

$$\int 3e^x \cos(3x) dx = 3e^x \cos(3x) - \int -9e^x \sin(3x) dx = 3e^x \cos(3x) + 9 \int e^x \sin(3x) dx$$

But then we have...

$$\int e^x \sin(3x) dx = e^x \sin(3x) - \int 3e^x \cos(3x) dx = \int e^x \sin(3x) dx = e^x \sin(3x) - \left(3e^x \cos(3x) + 9 \int e^x \sin(3x) dx \right)$$

Therefore, we have...

$$\int e^x \sin(3x) dx = e^x \sin(3x) - \left(3e^x \cos(3x) + 9 \int e^x \sin(3x) dx \right)$$

$$\int e^x \sin(3x) dx = e^x \sin(3x) - 3e^x \cos(3x) - 9 \int e^x \sin(3x) dx$$

$$10 \int e^x \sin(3x) dx = e^x \sin(3x) - 3e^x \cos(3x)$$

$$\int e^x \sin(3x) dx = \frac{e^x \sin(3x) - 3e^x \cos(3x)}{10} + C$$

$$\int e^x \sin(3x) dx = \frac{e^x}{10} (\sin(3x) - 3 \cos(3x)) + C$$

Alternatively, we can use an alternation of the tabular method of integration-by-parts. We choose $u = \sin(3x)$ and $dv = e^x$. We then have...

u	dv
$\sin(3x)$	e^x
$3 \cos(3x)$	e^x
$-9 \sin(3x)$	e^x

Therefore, we have...

$$\int e^x \sin(3x) dx = e^x \sin(3x) - 3 \cos(3x)e^x - 9 \int e^x \sin(3x) dx$$

Solving for our integral, we have...

$$\int e^x \sin(3x) \, dx = e^x \sin(3x) - 3 \cos(3x)e^x - 9 \int e^x \sin(3x) \, dx$$

$$10 \int e^x \sin(3x) \, dx = e^x \sin(3x) - 3e^x \cos(3x)$$

$$\int e^x \sin(3x) \, dx = \frac{e^x \sin(3x) - 3e^x \cos(3x)}{10} + C$$

$$\int e^x \sin(3x) \, dx = \frac{e^x}{10} (\sin(3x) - 3 \cos(3x)) + C$$