

Determining the direction of causality in the face of measurement error

13 February 2016

Gibran Hemani* and George Davey Smith

MRC Integrative Epidemiology Unit (IEU) at the University of Bristol, School of Social and Community Medicine, Bristol, UK

* Correspondence to: g.hemani@bristol.ac.uk

Abstract

With our ability to characterize the human phenome ever improving it is becoming increasingly common to use statistical tests to infer the causal relationships between correlated variables. A simple method for inferring if an exposure is on the causal pathway to an outcome is through mediation analysis, where the effect of an instrument on the outcome is tested before and after adjusting for the exposure. We show that in the face of measurement error this can lead to erroneous results, and that increasing sample size, rather than reducing bias, can often have the alarming effect of increasing certainty in the wrong answer. We argue that because measurement error is ubiquitous in phenotypic data, that mediation-based causal inference should be treated with caution. Finally, we demonstrate that Mendelian randomization is a method for causal inference that is robust to measurement error.

Introduction

With the explosion in data that measures the human phenome, of great interest is understanding the causal nature of the emerging putative correlations between measures of interest. Mediation is a commonly used statistical method for making causal inference in observational data¹⁻⁴, and it exists in many forms, from simple regression based systems to structural equation modeling. The principle behind mediation analysis can be explained as follows. Supposing an exposure has a causal effect on an outcome, then an ‘instrument’ that causes the exposure (a single nucleotide polymorphism (SNP), for example) should also influence the outcome. Therefore the influence of the instrument on the outcome conditional on the exposure should be zero. This forms the basis of a number of methods, such as the causal inference test (CIT)¹, which have been employed by a number of recent publications make causal inferences, often in large scale ‘omics datasets⁵⁻⁷.

While mediation based approaches are simple and convenient to implement, it is important to note where they may lead to unreliable results. One such mechanism is measurement error⁸⁻¹¹. Measurement (or observational) error is the difference between a measured value of a quantity and its true value. Such variability can arise through a whole plethora of mechanisms, which are often unique to the study design and difficult to avoid^{12,13}. Array technology is now commonly used to obtain high throughput phenotyping at low cost, but they come with the problem of having imperfect sensitivity, so for example methylation levels as measured by the Illumina450k chip are prone to have some amount of noise around the true value^{14,15}. Sensitivity is also an issue, for example if the unit of measurement of biological interest is the methylation level in a T cell, then measurement error of this value can be introduced by using methylation levels from whole blood samples because the measured value will be an assay of many cell types¹⁶.

Measurement error can indeed arise in more low-tech data too, for example when measuring body mass index (BMI) one is typically interested in using this as a proxy for obesity, but it is clear that the correlation between BMI and obesity is not perfect¹⁷. A similar problem of biological misspecification is unavoidable in disease diagnosis. Measurement error can also be introduced after the data has been collected, for example the transformation of non-normal data for the purpose of statistical analysis will lead to a new variable that

will typically have both bias and imprecision compared to the original variable. The sources of measurement error are not limited to this list¹³, and its impact has been explored in the context of mediation analysis in the epidemiological literature extensively^{8,11}.

An alternative statistical method that gained prominence at least in part as a solution to the problem of measurement error¹⁸ is Mendelian randomisation (MR)^{19,20}. Here an instrumental variable (typically a SNP) that has a robust, causal association with the exposure is used to proxy the exposure in an association test with the outcome. If there is no measurement error in the SNP then the causal association between the exposure and the outcome can be estimated by scaling the association between the SNP and the outcome by the association between the SNP and the exposure. This approach also has the useful property that it guards against unmeasured confounders driving the association between exposure and outcome; and if the biological effect of the SNP is understood then it can also aid against reverse causality driving the association.

Here, however, arises a potential issue. Often the biological effect of a SNP is not known, and therefore it can be difficult to determine for which of the two variables in a putative exposure-outcome association is the SNP a valid instrument. By definition, we expect that if the association is causal then a SNP for the exposure will be associated with the outcome, so if the researcher erroneously uses the SNP as an instrument for the outcome then they are likely to see an apparently robust causal association of outcome on exposure. Such a situation can arise in many scenarios. For example genome wide association studies (GWASs) that identify genetic associations for complex traits are by design hypothesis free and agnostic of genomic function, and it often takes years of follow up studies to understand the biological nature of a putative GWAS hit²¹. Another situation is where the causal direction between 'omic measures need to be determined, for example if a DNA methylation probe is associated with expression of an adjacent gene, then is a cis-acting SNP an instrument for the DNA methylation level, or the gene expression level⁷?

Mediation based methods, for example CIT and other related approaches typically attempt to use the data to resolve this issue using the following logic: if the SNP-exposure association is stronger than the SNP-outcome association then the causal effect must be in the direction of exposure to outcome. The same logic can indeed be applied to MR, but often when applied the sensitivity of these approaches to measurement error is not considered.

When being objective one must assume that measurement error is ubiquitous, and any measured variable is only an imperfect proxy of the biological quantity that the researcher intended to obtain. Here we show using theory and simulations how measurement error can lead to unreliable causal inference in the mediation-based CIT method, and we propose an extension to MR that, although does not eliminate the problem of measurement error, provides a framework in which to ascertain the causal direction when the biology of the instrument is not fully understood.

Methods

Simulations

CIT direction

MR causal test

Results

The influence of measurement error on CIT

Here we consider a simple causal model where an exposure, x , has an effect on an outcome of interest, y , and that there is a known SNP, g , which has a direct effect on the exposure. Measurement error of an exposure can be modeled as some transformation of the true value that leads to the observed value, $x_O = f(x)$. For example, we can define $f(x) = \alpha_m + \beta_m x + \epsilon_m$, where α_m and β_m influence the bias in the measurement of x ,

and ϵ_m represents the imprecision in the measurement of x . Here the true value of the exposure is partially explained by the genetic instrument, g , such that

$$x = \alpha_g + \beta_g + \epsilon_g$$

where β_g is the effect of the SNP on the exposure, and ϵ_g is the residual value of x ; and the outcome is partially explained by the exposure

$$y = \alpha_x + \beta_x + \epsilon_x$$

where β_x is the true effect of the exposure on the outcome. In the causal inference test (CIT), an omnibus p-value is generated from four hypothesis tests: 1) g is associated with x ; 2) g is associated with y ; 3) g is associated with $x|y$; and 4) g is independent of $y|x$. The 4th condition is necessary for causal inference, and can be expressed as $cov(g, y - \hat{y}) = 0$, where $\hat{y} = \hat{\alpha}_x + \hat{\beta}_x x_O$. When measurement error is introduced we can show through simple algebra that

$$\begin{aligned} cov(g, y - \hat{y}) &= cov(g, y) - cov(g, \hat{y}) \\ &= \beta_g \beta_x var(g) - D \beta_g \beta_x var(g) \end{aligned}$$

where

$$D = \frac{\beta_m^2 var(x)}{\beta_m^2 var(x) + var(\epsilon_m)}$$

Thus an observational study will find $cov(g, y - \hat{y}) = 0$ when the true model is causal only when $D = 1$. Therefore, if there is any measurement error that incurs imprecision (i.e. $var(\epsilon_m) \neq 0$) then there will remain an association between g and $y|x$, which is in violation of the 4th condition of the CIT. Note that measurement bias alone is insufficient to lead to a violation of the test statistic assumptions.

An alternative method to infer causality that is in principle robust to measurement error in the exposure variable, is Mendelian randomization (MR). The origins and applications of MR have been explained in detail, but the principle in which this is achieved is straightforward. Whereas mediation makes causal inference by contrasting the association of the instrument on the outcome before and after adjusting for the exposure, MR uses the instrument as a proxy for the exposure that is assumed to be devoid of measurement error and independent of potential confounders. In this setting, the problem is reformulated: instead of the researcher being agnostic about whether the instrument, g , has a direct influence on x or y , the researcher must select an instrument which is known to have a direct effect on x . Thus if g serves as a proxy for x , one can reason that the influence of g on y is the proportion of the effect of x on y that is explained by the effect of g on x . Hence the effect estimate β_{MR} of x on y is estimated as

$$\beta_{MR} = \frac{\beta_y}{\beta_g} = \frac{cov(y, g)}{cov(x_O, g)} = \frac{cov(\beta_x x + \epsilon_x, g)}{cov(x + \epsilon_x, g)}$$

which reduces to

$$\lim_{n \rightarrow \infty} \hat{\beta}_{MR} = \frac{\beta_x cov(x + \epsilon_x)}{cov(x + \epsilon_x)} = \beta_x$$

thus we obtain an estimate of the effect of x_O on y that is unbiased by measurement error in x .

References

1. Millstein, J., Zhang, B., Zhu, J. & Schadt, E. E. Disentangling molecular relationships with a causal inference test. *BMC genetics* **10**, 23 (2009).
2. Schadt, E. E. *et al.* An integrative genomics approach to infer causal associations between gene expression and disease. *Nature Genetics* **37**, 710–717 (2005).
3. Gayan, J., Gonzalez-Perez, A. & Ruiz, A. ‘Does replication groups scoring reduce false positive rate in SNP interaction discovery? Response’. *BMC Genomics* **11**, 403 (2010).
4. Tavtigian, S. V. *et al.* Rare, Evolutionarily Unlikely Missense Substitutions in ATM Confer Increased Risk of Breast Cancer. *The American Journal of Human Genetics* **85**, 427–446 (2009).
5. Koestler, D. C. *et al.* Integrative genomic analysis identifies epigenetic marks that mediate genetic risk for epithelial ovarian cancer. *BMC medical genomics* **7**, 8 (2014).
6. Liu, Y. *et al.* Epigenome-wide association data implicate DNA methylation as an intermediary of genetic risk in rheumatoid arthritis. *Nature biotechnology* **31**, 142–7 (2013).
7. Waszak, S. M. *et al.* Variation and genetic control of chromatin architecture in humans. *Cell* **162**, 1039–1050 (2015).
8. Cessie, S. le, Debeij, J., Rosendaal, F. R., Cannegieter, S. C. & Vandenbroucke, J. P. Quantification of bias in direct effects estimates due to different types of measurement error in the mediator. *Epidemiology (Cambridge, Mass.)* **23**, 551–60 (2012).
9. Nagarajan, R. & Scutari, M. Impact of noise on molecular network inference. *PloS one* **8**, e80735 (2013).
10. Shpitser, I., VanderWeele, T. & Robins, J. On the validity of covariate adjustment for estimating causal effects. *Proceedings of the Twenty Sixth Conference on Uncertainty in Artificial Intelligence (UAI-10)* 527–536 (2010).
11. Blakely, T., McKenzie, S. & Carter, K. Misclassification of the mediator matters when estimating indirect effects. *Journal of epidemiology and community health* **67**, 458–66 (2013).
12. Houle, D., Pélabon, C., Wagner, G. & Hansen, T. Measurement and meaning in biology. *The Quarterly Review of Biology* **86**, 3–34 (2011).
13. Hernán, M. a & Cole, S. R. Invited Commentary: Causal diagrams and measurement bias. *American journal of epidemiology* **170**, 959–62; discussion 963–4 (2009).
14. Harper, K. N., Peters, B. a & Gamble, M. V. Batch effects and pathway analysis: two potential perils in cancer studies involving DNA methylation array analysis. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* **22**, 1052–60 (2013).
15. Chen, Y.-a. *et al.* Discovery of cross-reactive probes and polymorphic CpGs in the Illumina Infinium HumanMethylation450 microarray. *Epigenetics : official journal of the DNA Methylation Society* **8**, 203–9 (2013).
16. Houseman, E. A. *et al.* DNA methylation arrays as surrogate measures of cell mixture distribution. *BMC bioinformatics* **13**, 86 (2012).
17. Ahima, R. S. & Lazar, M. A. Physiology. The health risk of obesity—better metrics imperative. *Science (New York, N.Y.)* **341**, 856–8 (2013).
18. Ashenfelter, O. & Krueger, A. B. Estimates of the Economic Return to Schooling from a New Sample of Twins. *The American Economic Review* **84**, 1157–1173 (1994).
19. Davey Smith, G. & Ebrahim, S. Mendelian randomization: prospects, potentials, and limitations. *International journal of epidemiology* **33**, 30–42 (2004).

- 20.Davey Smith, G. & Hemani, G. Mendelian randomization: genetic anchors for causal inference in epidemiological studies. *Human molecular genetics* **23**, R89—R98 (2014).
- 21.Claussnitzer, M. *et al.* FTO Obesity Variant Circuitry and Adipocyte Browning in Humans. *The New England journal of medicine* **373**, 895–907 (2015).