Forecast-aware latent modeling for longitudinal prediction of pathological complete response from breast MRI

Author information to be completed

Affiliations and corresponding author to be completed

Abstract

Longitudinal breast MRI can reveal treatment response, but endpoint classifiers do not explicitly model the future trajectory implied by an observed history. We investigate forecast-aware latent modeling for pathological complete response (pCR) prediction using three-phase dynamic contrast-enhanced MRI, baseline clinical variables, and assigned-arm scenarios. An autonomous predictor estimates future latent states, and an attention readout combines the predicted terminal state, its change, and direct history features. The cohort comprises 976 patients and 3,641 examinations, with 801 development patients and a fixed internal holdout of 175 patients, including 56 positive outcomes. Three principal variants distinguish prediction without future-state supervision, auxiliary-supervised direct prediction, and forecast-attention readout. We evaluate training seeds 42, 44, 2026, and 3407, averaging five fold probabilities within each seed before computing metrics and then equally averaging seed-specific metrics. Forecast attention achieved mean T0/T1/T2 AUROCs of 0.7438/0.7448/0.7459 and Macro AUROC/AP/F1 of 0.7449/0.6175/0.5596, compared with 0.7269/0.5924/0.5484 for the control without future-state supervision. Its Macro AUROC sample seed standard deviation was 0.0040. A supplementary mean-pooling control had similar Macro AUROC (0.7436) and higher F1 (0.5690). The forecast-attention versus control AUROC difference had approximate paired-bootstrap $p = 0.00306$ and nine-comparison Holm $p = 0.02754$, but did not survive adjustment over all 90 tests ($p = 0.26622$). Compared with auxiliary-supervised direct readout, forecast attention had lower specificity, which survived all-90 adjustment, and worse probability-quality point estimates. These exploratory findings require independent validation because the seed set and reporting policies were selected retrospectively.

Keywords: Breast MRI, Pathological complete response, Longitudinal imaging, Latent dynamics, Treatment response prediction

1. Introduction

Neoadjuvant systemic therapy provides an opportunity to monitor breast cancer before surgery and to study how an individual tumor responds over time. Pathological complete response (pCR) is a clinically meaningful endpoint associated with long-term outcomes, although its relationship to prognosis varies across biological subtypes and it should not be equated with a universally valid trial-level surrogate (Cortazar et al., 2014). Predicting pCR from imaging could support response assessment and the design of future adaptive studies. Such a prediction remains a prognostic estimate under an observed treatment context; it does not establish that changing treatment would improve an individual patient's outcome.

Dynamic contrast-enhanced MRI captures three-dimensional anatomy and contrast uptake patterns that evolve during therapy. Prior work in the I-SPY setting has demonstrated associations between MRI-derived tumor measurements and pathological response or subsequent outcomes (Hylton et al., 2012, 2016). Public multicenter imaging resources also make it possible to examine learned representations across repeated visits (Li et al., 2022; Newitt et al., 2021). However, a longitudinal prediction system must operate at the time a decision is considered: a baseline prediction can use baseline information, whereas an interim prediction can use the examination history accumulated by that interim visit. Future scans may inform representation learning on development patients but cannot become inputs to predictions for holdout patients at an earlier landmark.

Most direct classifiers optimize the endpoint of interest from observed images and clinical variables. This is a reasonable objective, but it does not explicitly require the representation to capture how the tumor state may evolve. A different strategy is to predict future imaging information and use the prediction to inform the endpoint readout. Image synthesis provides one route to such a representation, but prediction in a compact latent space can focus the training signal on state evolution without requiring a pixel-space image generator at inference. This distinction is useful for response prediction: the relevant question is whether the learned forecast contributes information to classification, rather than whether a generated image is

visually convincing.

Joint-embedding predictive learning offers a general mechanism for predicting target representations computed by a slowly updated encoder (Assran et al., 2023). In a longitudinal medical setting, adapting that principle requires explicit handling of partial histories, missing future observations, treatment context, and the distinction between an auxiliary training target and a classifier input. A model may benefit from a future-state loss while still making a direct prediction from the observed history. Conversely, directly including forecast features in the classifier changes both its information pathway and its parameterization. These effects should be examined through matched controls rather than attributed to a single generic "world model" component.

We investigate forecast-aware latent modeling for longitudinal pCR prediction from breast MRI. The model encodes three selected DCE phases at each available visit, fuses observed MRI with baseline clinical information and an assigned treatment-arm scenario, and autonomously rolls the current latent state forward to a terminal stage. A spatial attention mechanism summarizes the predicted terminal state and the predicted change from the current state for pCR classification. We compare the complete model with two direct-readout controls: one without future-state supervision and one with the same auxiliary supervision as the complete model. A supplementary mean-pooling model further examines the attention component.

The study makes three concrete contributions. First, it formulates a common representation and masked future-state objective that can be applied to available T0–T2 histories while using observed later scans only as development-time targets. Second, it provides a staged comparison that distinguishes the effect of auxiliary prediction from the joint effect of forecast readout and attention. Third, it reports landmark-specific discrimination, class-specific errors, calibration, hypothetical decision curves, subgroup heterogeneity, archived forecasting diagnostics, and paired patient-level uncertainty on a fixed internal holdout. The manuscript evaluates each variant with training seeds 42, 44, 2026, and 3407. Five fold probabilities are averaged within each seed before computing metrics; the metric values are then equally averaged across seeds and accompanied by descriptive seed standard deviations. This seed set and aggregation policy were chosen retrospectively, and patient-level inference conditions on the selected fitted predictors. Seed-selection and upstream-independence limits remain explicit.

2. Related work

MRI biomarkers and learned pCR prediction.. MRI-based response assessment has a substantial clinical and quantitative foundation. Functional tumor volume and related measurements have been studied in longitudinal breast MRI cohorts, including I-SPY (Hylton et al., 2012, 2016). The ACRIN 6698 study additionally examined diffusion-derived measurements in relation to response (Partridge et al., 2018). These studies motivate imaging-based monitoring but do not establish that a learned model using different inputs or a different evaluation split has equivalent clinical validity. Our experiments use selected DCE phases and baseline clinical variables; diffusion imaging is not an input to the present model.

Learning-based approaches can combine complementary image sequences or repeated examinations. Synthesis-based representation learning has been explored for multi-sequence 3D/4D MRI (Han et al., 2024), and recent studies have examined spatiotemporal or early-treatment information for pCR prediction (Tang et al., 2025; Du et al., 2025). Such studies provide relevant methodological context, but differences in patient selection, imaging availability, endpoints, and holdout construction prevent direct numerical comparisons with our results. We therefore do not claim superiority to published systems from cross-paper metric comparisons. Reimplementation of suitable external baselines under the same patient split remains necessary.

Predictive latent representations.. Predicting a target embedding rather than reconstructing the input image is a central idea in joint-embedding predictive learning (Assran et al., 2023). We use an exponential-moving-average target encoder to define latent states and supervise autonomous future predictions. The present objective is a supervised pCR loss combined with a masked multi-step latent-state loss. It does not activate a separate single-step teacher-forcing JEPA loss, image reconstruction loss, or latent-variance regularizer. This objective-level distinction matters when interpreting the internal experiment name NoJEPA: that control removes future-state supervision, while retaining the common fusion architecture.

Attention and volumetric image representation.. Transformer attention provides a flexible way to combine image tokens and contextual information (Vaswani et al., 2017). Compact convolutional representations can reduce the cost of processing volumetric MRI, and vector-quantized generative models provide one source of pretrained convolutional encoders (Esser et al., 2021). Here the pretrained codec contributes a continuous convolutional encoder. The quantizer and image decoder are bypassed, and the main experiments do not synthesize MRI volumes. The final attention weights summarize latent tokens that have already undergone spatial mixing; they are therefore not validated lesion segmentations or anatomical explanations.

3. Method

3.1. Longitudinal prediction problem

We consider prediction of pathological complete response (pCR) from a patient's available magnetic resonance imaging (MRI) history during neoadjuvant treatment. Canonical examination stages are indexed by $t \in \{0, 1, 2, 3\}$. At a




Figure 1: Forecast-aware latent modeling and matched controls. (A) Real three-phase MRI crops from one development patient illustrate the same baseline-defined plane at T0, T1, and T2; the example and slice were selected without outcomes or pCR predictions. Only visits available at the prediction landmark are inputs. A shared continuous encoder and spatial Transformer produce visit tokens. (B) Clinical values, availability masks, and an assigned-arm scenario condition fusion and autonomous prediction to T3. The EMA encoder supplies the detached current state; available future MRI supplies training targets only. Dashed orange arrows denote training-target or loss paths. (C) Attention weights pool the predicted terminal state and its change; these features are combined with direct pCR-query features. (D) The primary variants separate removal of future-state supervision from the complete forecast readout. ForecastMean is a supplementary pooling control. One fold model is illustrated. Within each of seeds 42, 44, 2026, and 3407, evaluation averages five fold probabilities per patient before computing metrics; reported performance then averages the four seed metrics, without ensembling probabilities across seeds. MRI tiles are illustrative crops, not generated images, segmentation ground truth, or attention maps.

prediction landmark $k \in \{0, 1, 2\}$, the model receives only examinations acquired and available by that landmark. Let $X_t \in \mathbb{R}^{3\times 32\times 128\times 128}$ denote the processed examination at stage t, and let $o_t^{(k)}$ indicate whether this examination is available in the observed prefix. Missing examinations retain their canonical stage rather than shifting later examinations forward. The three channels contain the pre-contrast acquisition, the first post-contrast acquisition, and a metadata-selected late post-contrast acquisition, respectively. The late acquisition is not assumed to have the same acquisition index for all examinations.

The observed information is

$$\mathcal{H}_k = \{(X_t, t) : o_t^{(k)} = 1\} \cup \{c_k, m_k, a_k, \tau_k\}, \qquad (1)$$

where $c_k \in \mathbb{R}^4$ contains the available clinical variables, m_k their availability indicators, a_k the available assigned treatment-arm category, and τ_k its declared availability stage. The clinical variables are age at screening, hormone-receptor positivity, HER2 positivity, and a binary MammaPrint variable. Age is standardized using the current training partition; binary variables retain their original coding. Unavailable values are set to zero after standardization and accompanied by an explicit mask. The clinical token embeds the concatenated values and masks. Treatment conditioning uses 13 released arm categories and an additional unknown category. In this retrospective experiment, assigned arm and available baseline clinical variables are assumed to be known at T0. This is an assigned-arm scenario; the categorical input does not specify actual doses, treatment intervals, or treatment changes, and it does not support a causal interpretation.

The output is $p_k = \Pr(y = 1 \mid \mathcal{H}_k)$, where $y \in \{0, 1\}$ is the final pCR label. Future MRI is stored separately as training supervision and is excluded from the prediction input. T3 can provide a future representation target during training, but is never a prediction landmark in the reported T0–T2 protocol.

3.2. Continuous MRI encoding and multimodal fusion

Each observed examination is encoded independently by an online visit encoder E_θ. Its convolutional front end shares the same single-channel weights across the three acquisition phases. Two stride-two three-dimensional convolutional stages, residual blocks, and group normalization produce 64 continuous feature channels per phase at one-quarter input resolution along each spatial axis. The three phase-specific feature maps are concatenated, adaptively averaged to a $2 \times 4 \times 4$ grid, and projected to a latent width of $d = 768$. This gives $N = 32$ spatial tokens per examination. Learned spatial positional embeddings and an eight-layer spatial Transformer then produce the visit representation $Z_t = E_\theta(X_t) \in \mathbb{R}^{N \times d}$.

The convolutional front end is initialized from the same pre-existing VQ-GAN-style codec checkpoint for all compared models and is fine-tuned. Here, the model uses continuous features from the codec encoder before its compression projection. It does not use discrete codebook assignments, a quantization objective, a decoder, or adversarial learning during downstream pCR training. [TO COMPLETE: Document the original codec pretraining dataset, patient-level overlap with the present partitions, checkpoint provenance, and applicable redistribution terms.]

Stage embeddings distinguish examination times. Observed image tokens, one clinical token, and one treatment-arm token form the multimodal context. An eight-layer fusion Transformer additionally receives eight learned transition-query tokens for each of the three canonical transitions and four pCR-query tokens. Transition-query slots are retained at every landmark, independently of which future targets happen to be available. Their conditioning is derived entirely from the observed prefix. Context tokens interact bidirectionally; appended transition and pCR queries use a causal attention mask and can attend to the context and preceding query positions. Padding masks remove unobserved image tokens. This produces transition features $A_{j+1}^{(k)} \in \mathbb{R}^{8 \times d}$ for transitions $j \to j + 1$, and pCR-query features $Q_k \in \mathbb{R}^{4 \times d}$.

All spatial, fusion, and prediction Transformers use pre-normalization blocks, GELU feed-forward layers of width 3072, and dropout 0.3, following the attention-based architecture of Vaswani et al. (2017). Attention has 16 heads with 64 channels per head: its internal width is therefore 1024, followed by projection back to the 768-dimensional residual stream. The clinical and treatment tokens condition forecasting through the shared fusion network rather than through a separately supplied future treatment trajectory.

3.3. EMA target states and autonomous forecasting

A target visit encoder $E_{\bar{\theta}}$ is initialized as a copy of the online encoder and updated after each optimizer step by exponential moving average (EMA):

$$\bar{\theta} \leftarrow \beta\bar{\theta} + (1 - \beta)\theta, \qquad \beta = 0.999. \qquad (2)$$

The target encoder remains in evaluation mode and receives no gradients. It defines a common representation space for the observed initial state and available future targets,

$$S_t = \mathrm{stopgrad}(E_{\bar{\theta}}(X_t)) \in \mathbb{R}^{N \times d}. \qquad (3)$$

Encoding a future target in this branch supplies a loss target only; it does not make that examination available to the forecasting or classification input.

The predictor starts from the latest actually observed state in $\mathcal{H}_k$ and recursively advances to T3. For the reported landmarks, this initial state is S_k. At each future transition, it uses the corresponding transition features and its previously generated states. A history input additionally embeds the difference between a source state and its latest available predecessor, with a separate indicator for whether a predecessor exists. Observed history is used where available; after the landmark, the history is extended using model predictions. Thus, a missing future examination removes supervision at that time but does not interrupt the simulated transition sequence.

The prediction network is an eight-layer time-causal Transformer. Each transition block contains its eight transition tokens and 32 spatial source-state tokens, together with learned spatial and temporal embeddings. Tokens can interact within a block; attention between transition blocks is directed from earlier to later blocks. A learned

root token remains available when some blocks are masked. The network predicts a residual state increment,

$$\widehat{S}_{j+1} = \widetilde{S}_j + G_{\phi,j}\left(\widetilde{S}_{k:j}, A^{(k)}_{k+1:j+1}, D_{k:j}\right), \qquad j = k, \ldots, 2, \tag{4}$$

where $\widetilde{S}_k = S_k$, subsequent $\widetilde{S}_j$ are generated states, and $D_{k:j}$ denotes the available predecessor differences and validity indicators. The residual output projection is initialized to zero, so prediction initially preserves the source state. The transition features summarize the complete legal prefix even when only a subset of previous state differences is available to a particular prediction block.

Forecasts are generated autonomously during training and inference. A true future state never replaces a predicted predecessor. The classifier likewise uses this autonomous trajectory. The configuration includes no teacher-forced next-state loss in these experiments. EMA encoding of real future examinations is used exclusively to evaluate representation error against available targets.

3.4. Prediction-conditioned pCR readout

The full model, ForecastAttention, combines a direct summary of the observed prefix with summaries of the predicted terminal state and its change from the current state. First, the four pCR-query features are averaged to obtain $q_k = \frac{1}{4}\sum_{r=1}^{4} Q_{k,r}$. For spatial token n, the forecast change is $\Delta S_{k,n} = \widehat{S}_{3,n} - S_{k,n}$. A learned scalar scorer defines

$$\alpha_{k,n} = \frac{\exp\left(w^\top \mathrm{LN}(\widehat{S}_{3,n})\right)}{\sum_{u=1}^{N} \exp\left(w^\top \mathrm{LN}(\widehat{S}_{3,u})\right)}. \tag{5}$$

The same weights pool terminal state and change,

$$v_k = \sum_{n=1}^{N} \alpha_{k,n} \widehat{S}_{3,n}, \qquad d_k = \sum_{n=1}^{N} \alpha_{k,n} \Delta S_{k,n}. \tag{6}$$

The final prediction is

$$p_k = \sigma\left(b + u^\top \mathrm{LN}([q_k; v_k; d_k])\right). \tag{7}$$

The scorer is initialized with $w = 0$, corresponding to uniform pooling at initialization. Classification gradients propagate through the generated terminal state, predictor, and transition-conditioning pathway. The observed EMA reference state remains detached. The attention weights summarize transformed spatial tokens and should not be interpreted as validated anatomical localization or tumor segmentation.

For DirectReadout, pCR is instead predicted directly from q_k using a layer-normalized linear head. Its future-state predictor is trained as an auxiliary task, but generated states are not explicit inputs to the pCR head. The direct head has input width d, whereas the full readout has width $3d$. Their comparison therefore evaluates the complete forecast-conditioned readout, including its state and change inputs and classifier width. An additional mean-pooling variant, when reported, retains the forecast-conditioned readout and replaces $\alpha_{k,n}$ by $1/N$.

3.5. Training objective and matched ablations

Let $\mathcal{B}_y$ be the labeled patients in a training batch, and let $m^{\mathrm{future}}_{i,t}$ indicate an actually available future target for patient i after its sampled landmark. The pCR objective is unweighted binary cross-entropy,

$$\mathcal{L}_{\mathrm{pCR}} = -\frac{1}{|\mathcal{B}_y|} \sum_{i \in \mathcal{B}_y} \left[y_i \log p_i + (1 - y_i) \log(1 - p_i) \right]. \tag{8}$$

Define $M_i = \sum_t m^{\mathrm{future}}_{i,t}$ and $\mathcal{B}_+ = \{i : M_i > 0\}$. Autonomous representation supervision is patient-equal:

$$\mathcal{L}_{\mathrm{roll}} = \frac{1}{|\mathcal{B}_+|} \sum_{i \in \mathcal{B}_+} \frac{1}{M_i} \sum_t m^{\mathrm{future}}_{i,t} \frac{\|\widehat{S}_{i,t} - S_{i,t}\|_1}{Nd}. \tag{9}$$

If a batch contains no future targets, this term is zero. Each available future examination has equal weight within a patient; the loss uses no horizon discount, lesion weighting, or fabricated target for a missing visit. Patients without future MRI still contribute pCR supervision.

The full model and DirectReadout minimize

$$\mathcal{L} = \mathcal{L}_{\mathrm{pCR}} + 0.5\mathcal{L}_{\mathrm{roll}}. \tag{10}$$

NoJEPA uses the direct pCR head with $\mathcal{L} = \mathcal{L}_{\mathrm{pCR}}$. This historical control name denotes removal of future-state auxiliary supervision, rather than removal of every transition-related module: it retains the shared fusion architecture and transition-query slots. Consequently, DirectReadout versus NoJEPA isolates the added autonomous representation objective under the same direct readout, whereas ForecastAttention versus DirectReadout assesses the added forecast-conditioned classification pathway. The latter comparison cannot isolate spatial attention alone.

All three models omit teacher-forced JEPA, image reconstruction, flow-matching, and variance-regularization losses. In particular, the future objective reported here is autonomous state prediction, despite the JEPA-related implementation lineage. During training, patients are shuffled once per epoch and one actual available T0–T2 landmark is sampled uniformly for each patient. This balances patient contributions without treating a patient with more visits as several independently sampled patients. The same clinical endpoint supervises each sampled prefix; validation assesses all available eligible prefixes using patient-equal weighting.

4. Dataset and preprocessing

4.1. Cohort construction and clinical variables

We retrospectively analysed longitudinal breast dynamic contrast-enhanced MRI (DCE-MRI) from the public I-SPY2 and ACRIN-6698 imaging resources Li et al. (2022); Newitt et al. (2021). Source imaging identifiers were linked to research identifiers using the accompanying official identity mapping, and outcomes and baseline clinical variables

were obtained from the official I-SPY2 Imaging Cohort 1 clinical spreadsheet. The final cohort comprised 976 patients with 3,641 usable examinations. Native imaging identifiers belonged to the ISPY2 and ACRIN-6698 namespaces for 712 and 264 patients, respectively; these identifiers were mapped to the same clinical release and do not define independent development and external validation cohorts. The official collection pages identify the source imaging and clinical release as CC BY 4.0. [TO COMPLETE: Confirm terms for redistributing derived assets and document the applicable ethics approval or waiver and original consent arrangements.]

Eligibility required an actual baseline examination, an available binary pCR label, usable baseline registered image assets, and a nonempty baseline localisation produced by the existing MAMA-MIA segmentation workflow. Of 980 clinically matched source candidates, three lacked a baseline examination and one had an empty or unavailable predicted baseline localisation. Their exclusion yielded 976 patients and 3,642 candidate examinations. One post-treatment examination was subsequently marked unavailable because its saved within-examination phase transforms displaced the postcontrast images outside the prescribed baseline crop. The patient's earlier examinations were retained, leaving 3,641 usable examinations. Missing follow-up examinations did not cause patient exclusion. No additional admission thresholds based on normalised cross-correlation, field-of-view ratios, functional tumour volume, or the availability of an accepted adjacent longitudinal pair were applied.

The endpoint was the released binary pCR field: 313 patients were labelled positive and 663 negative. [TO COMPLETE: Verify and state the release-specific pathological definition of pCR, including breast and nodal criteria and the handling of in-situ disease; do not infer this definition from the binary column alone.] Baseline covariates were age at screening, hormone-receptor (HR) status, HER2 status, and the released binary MammaPrint indicator. Age was available for 973 patients; all three binary covariates were available for the entire cohort. The official assigned treatment arm, represented by 13 categories, was also supplied to the model. The current experiments use an *assigned-arm scenario*: the assigned arm and available baseline covariates are assumed known at baseline. The spreadsheet does not establish prospective allocation timestamps, doses, cycle-level exposure, or interval treatment schedules, and the model does not reconstruct these quantities.

Examinations were indexed by the canonical study stages T0–T3 rather than continuous elapsed time. Availability at these stages was 976, 935, 875, and 855 patients, respectively (Table 1); 803 patients had all four examinations and 861 had T0–T2. The nominal source protocol acquired T0 before randomization, T1 after three weeks of the first regimen (paclitaxel with or without an experimental agent), T2 between paclitaxel and anthracycline regimens, and T3 after four cycles of anthracycline treatment and before surgery (Li et al., 2022). Because T0 MRI preceded randomization, supplying the eventual assigned arm at T0 is a retrospective conditional scenario, rather than a prospectively available predictor before treatment allocation. The source protocol used 1.5 or 3.0 Tesla scanners with dedicated breast coils and required the same scanner configuration within a patient. These are nominal protocol specifications, not a patient-level audit of timing or scanner compliance. Twenty-three patients had T0 only and remained eligible for pCR supervision. Their unavailable future MRI targets were masked, and no synthetic follow-up images were substituted. Model inputs and evaluation landmarks were restricted to T0, T1, and T2; a real T3 examination, when available in the training partition, could provide a future-state target. [TO COMPLETE: Report the actual examination intervals, acquisition centres, scanner distribution, and sequence-parameter ranges for the final analysed cohort.]

4.2. Patient partitioning and evaluation scope

The cohort was divided into 801 development patients and a fixed internal holdout of 175 patients, including 56 pCR-positive patients. All holdout patients had actual T0, T1, and T2 examinations, allowing matched longitudinal evaluation; 163 also had T3. The holdout retained 139 patients with complete T0–T2 observations from the historical benchmark and added 36 eligible patients. Additional patients were selected using outcome quotas and a fixed baseline-clinical distribution balancing rule with split seed 2026, without using model predictions or performance. This fixed cohort-construction seed is distinct from the training-seed comparison: the split is unchanged for training seeds 42, 44, 2026, and 3407. Development folds were stratified by pCR and examination-availability pattern; rare joint strata were combined within the same pCR class. Each development patient appeared in validation exactly once across the five folds. Fold 1 contained 640 training and 161 validation patients; the remaining folds each contained 641 training and 160 validation patients. Training, validation, and holdout identities were disjoint within each fold, and the same holdout was used for every fold model at each training seed.

This partition is an internal downstream benchmark. In particular, 29 of the 36 added holdout patients had belonged to the development population of an earlier experiment, so earlier trained models cannot be treated as held out on the entire new set. The present downstream models were trained using the new fold definitions. Patient independence of the supplied codec and segmentation pretraining checkpoints relative to the final holdout has not yet been established; that provenance must be audited before stronger claims of end-to-end independence. The complete-examination holdout eligibility also restricts inference about patients with missing early follow-up examinations.

4.3. Registration, fixed localisation, and normalisation

Within each examination, postcontrast phases were registered to the precontrast reference. Follow-up examinations

Table 1: Characteristics of the analysed cohort. Development statistics count each of the 801 development patients once, rather than pooling overlapping training folds. Age is reported as mean ± sample standard deviation among patients with observed age; other entries are counts.

Characteristic	All	Development	Holdout
Patients	976	801	175
pCR positive	313	257	56
pCR negative	663	544	119
Age (years)	48.7 ± 10.5	48.7 ± 10.6	48.9 ± 10.2
Missing age	3	3	0
HR positive	533	434	99
HER2 positive	240	195	45
Available T0	976	801	175
Available T1	935	760	175
Available T2	875	700	175
Available T3	855	692	163

were aligned to the patient's T0 reference using the saved longitudinal transforms, retaining a rigid fallback when the final deformable candidate was not usable. Baseline localisation used the existing MAMA-MIA nnU-Net prediction on the native first-postcontrast examination Garrucho et al. (2025). The centre of the largest 26-connected predicted component defined a fixed baseline crop, which was reused at every actual follow-up. Follow-up segmentation and follow-up-driven recentering were not used.

Each examination was represented by three channels: the precontrast phase, the first postcontrast phase, and the late phase explicitly specified in the examination metadata. The late channel was not selected by assuming a fixed phase number or taking the last available phase. Images were resampled to $32 \times 128 \times 128$ voxels with spacing $2.0 \times 0.7032 \times 0.7032$ mm in the slice and two in-plane dimensions, approximately covering $64 \times 90 \times 90$ mm. Linear interpolation was used for MRI intensities and nearest-neighbour interpolation for masks and acquisition support, with zero padding outside the acquired field of view. The localisation mask determined the crop; it was not a contemporaneous ground-truth tumour segmentation or a future observation supplied to the classifier.

Native metadata, baseline mask headers, registered reference grids, and saved transforms were maintained as matched versions. Explicit array-index mappings, rather than placeholder NIfTI affines, defined sampling positions. The exported arrays were checked for finite values, consistent coordinate mapping, and numerical agreement after compressed storage and readback. These checks establish data and geometry consistency but do not establish anatomical registration or segmentation accuracy. The fixed crop partly truncated the largest predicted baseline component for 85 patients; these patients were retained and the limitation was recorded.

Raw cropped intensities were stored once and shared across folds. For each fold, a single intensity mean and population standard deviation were fitted to nonzero precontrast foreground voxels from every unique actual examination of that fold's training patients. The same scalar transform was applied once to the foreground of all three channels, stages, and partitions; background and padding remained zero. Validation and holdout images were excluded from fitting. Age normalisation was likewise fitted to observed training-patient values; HR, HER2, and MammaPrint retained their official binary coding, with missingness represented explicitly.

5. Experimental design

5.1. Development protocol and landmark evaluation

The 801 development patients were partitioned into five patient-level folds. Fold 1 used 640 training and 161 validation patients; folds 2–5 each used 641 training and 160 validation patients. The same 175-patient holdout was evaluated by every fold model and every variant. Within each seed and variant, the five test evaluations are repeated evaluations of overlapping trained models on the same patients, rather than five independent test cohorts. Repeating training seeds also reuses this holdout. Patient identities were disjoint between training, validation, and holdout within the current split.

At each training epoch, a patient contributed one randomly selected available T0, T1, or T2 prefix. Only MRI observations available at or before the selected landmark were classifier inputs. Later observed visits, including T3, could supply latent targets for the auxiliary loss on training patients. Missing targets were masked. The 23 patients with only T0 remained eligible for pCR training but supplied no future-state loss. At evaluation, all three landmarks were evaluated for the 175 holdout patients, each of whom had T0, T1, and T2. T3 was not an evaluation landmark and was not used as observed input for these predictions.

5.2. Model comparisons

The principal comparison comprises three variants trained with seeds 42, 44, 2026, and 3407 using the same five development folds. Within each variant and seed, the five fold-specific models form an equal-weight probability ensemble. Performance is then summarized across the

four seed-specific ensembles by an equal mean of their metric values. NoJEPA uses direct pCR readout and a zero future-state loss weight. DirectReadout uses the same direct readout with a future-state loss weight of 0.5. ForecastAttention retains that auxiliary weight and replaces the direct readout with a concatenation of direct pCR features, attention-pooled predicted terminal state, and attention-pooled predicted change. All variants retain the shared image encoder, clinical inputs, assigned-arm encoding, fusion architecture, and training schedule.

The DirectReadout–NoJEPA contrast most directly tests adding future-state supervision. The ForecastAttention–DirectReadout contrast jointly changes the use of forecast features, spatial pooling, and classifier input width. It cannot isolate attention alone. For that purpose, the supplementary ForecastMean comparison uses forecast readout with mean spatial pooling and otherwise matches the forecast-attention configuration. All manuscript-facing pCR results first average the five fold probabilities separately for each seed, patient, and landmark, then compute metrics within each seed, and finally average those metrics across the four seeds. Probabilities are not averaged across seeds. No VQ-latent reconstruction-head, change-weighted-loss, or state-conditioned-head variants are included.

5.3. Implementation and optimization

The network uses a model width of 768, eight layers in each of the spatial encoder, multimodal fusion module, and predictor, 16 attention heads with 64 dimensions per head, and feed-forward width 3,072. The 1,024-dimensional concatenated attention-head representation is projected back to the 768-dimensional residual stream. The spatial token grid is $2 \times 4 \times 4$. The configuration uses eight state queries, four pCR queries, dropout 0.3, and an EMA coefficient of 0.999. The pretrained convolutional encoder is fine-tuned jointly with the remaining network. Its training-patient provenance requires additional documentation, as discussed in Section 7.5.

We use AdamW (Loshchilov and Hutter, 2019) with an initial base learning rate of 10^{-4} for both the encoder and the main network, weight decay 0.05, batch size 16, no gradient accumulation, and gradient-norm clipping at 1.0. Training uses BF16 mixed precision. A ten-epoch warm-up increases the learning rate from 10^{-5} to 10^{-4}, followed by cosine decay to 10^{-6} at epoch 100. Training is capped at 200 epochs, with early stopping after 50 epochs without improvement. The best checkpoint is selected independently in each fold using validation patient-equal-prefix F1 at threshold 0.5. This is an F1 selection protocol, even though AUROC is a principal reporting measure. Deterministic execution was not enabled, so a fixed seed does not imply bitwise reproducibility across hardware or software changes.

5.4. Metrics, aggregation, and patient-level uncertainty

We report 14 complementary metrics: AUROC, non-interpolated average precision (AP), F1, accuracy, sensitivity, specificity, positive and negative predictive value (PPV/NPV), balanced accuracy, Matthews correlation coefficient (MCC), Youden index, negative log-likelihood (NLL), Brier score, and expected calibration error (ECE) with ten equal-width probability bins. Confusion-matrix counts are additionally retained. Thresholded primary metrics use 0.5. MCC and balanced accuracy help interpret performance under class imbalance (Chicco and Jurman, 2020); positive prevalence is $56/175 = 0.32$. No holdout threshold optimization or probability recalibration was performed.

Let $p_{isf\ell}$ be the probability from the model trained with seed $s \in \mathcal{S} = \{42, 44, 2026, 3407\}$ in fold f for patient i at landmark ℓ. Within each variant, the reporting estimand is

$$\begin{aligned} \bar{p}_{is\ell} &= \frac{1}{5}\sum_{f=1}^{5} p_{isf\ell}, \\ M_{s\ell} &= M\big(\{y_i, \bar{p}_{is\ell}\}_{i=1}^{175}\big), \\ M_{s,\text{macro}} &= \frac{1}{3}\sum_{\ell=0}^{2} M_{s\ell}, \\ \widehat{M}_{\ell} &= \frac{1}{4}\sum_{s\in\mathcal{S}} M_{s\ell}. \end{aligned} \tag{11}$$

Macro and seed averaging commute for these metric summaries; neither operation is a probability average across time or seeds. Thresholded metrics apply 0.5 separately to each seed's five-model ensemble probabilities. Main performance tables report the mean and sample standard deviation of the four seed-specific metrics, using the sample-variance denominator of three. Macro standard deviations are computed from the four seed-specific Macro values, rather than averaging landmark standard deviations. This spread describes the four selected training realizations. It is not a standard error, a patient-level interval, or evidence for the distribution of all possible training seeds. The estimand is the average performance of four five-model predictors; it is not the performance of a single twenty-model probability ensemble or a single model trained on the full development cohort. Individual-fold metrics remain audit-only.

For conditional patient-level uncertainty, we draw 2,000 bootstrap samples of the 175 patients with replacement. Each draw is shared across every seed, variant, and landmark. Within a draw, we recompute metrics separately from each seed-specific ensemble, average landmark metrics for Macro, and average the four seed metrics. The 2.5th and 97.5th percentiles give unadjusted intervals for this seed-mean metric or paired difference. Seeds are held fixed and are not resampled. The intervals therefore exclude retraining, retrospective seed-set selection, model selection, and aggregation-policy uncertainty. The five folds, four seeds, and three repeated landmarks do not increase the independent patient count beyond 175.

ROC and precision–recall curves are computed separately within each seed-specific probability ensemble, interpolated onto common false-positive-rate and recall grids, respectively, and then averaged across seeds for display. The numerical AUROC and non-interpolated AP are computed within seeds before averaging; their means need not equal the area under the displayed mean curve. Pointwise bands use the shared patient resamples and retain seed averaging. Reliability diagrams form five equal-count rank groups of 35 patients separately within each seed and average their group summaries across seeds. These are descriptive summaries of several predictors, distinct from the ten equal-width bins used to calculate each seed's ECE.

5.5. Exploratory paired significance checks

We use approximate, two-sided, null-centered paired patient-bootstrap tests for all 14 metrics, including AUROC (Davison and Hinkley, 1997). The tested quantity is the difference in the metric averaged across the four fixed seed-specific predictors. We generate $B = 99{,}999$ shared patient resamples with random-number seed 20261103. Every resample draws 175 patients with replacement and is shared across seeds, models, and landmarks. Metrics are recomputed within seeds and then equally averaged, preserving all pairing. No model is retrained, no seed is resampled, and no fold- or seed-level t test is used.

Let Δ_b^* denote the seed-mean metric difference in resample b and Δ_{obs} its observed value. We estimate the two-sided p value by

$$p = \frac{1 + \sum_{b=1}^{B} \mathbb{I}\big(|\Delta_b^* - \Delta_{\text{obs}}| \geq |\Delta_{\text{obs}}|\big)}{B + 1}. \tag{12}$$

Ties are included. The plus-one convention prevents Monte Carlo zero values, but does not make this bootstrap approximation finite-sample exact. Non-smooth statistics such as AP, F1, and binned ECE warrant particular caution.

The principal reporting family contains nine tests: the three pairwise comparisons among NoJEPA, DirectReadout, and ForecastAttention, each for Macro AUROC, AP, and F1. The secondary family contains 81 tests: 27 landmark-specific AUROC/AP/F1 comparisons among the principal models; 33 comparisons of the other 11 Macro metrics; 12 ForecastAttention–ForecastMean comparisons for AUROC/AP/F1 at T0, T1, T2, and Macro; and nine T2–T0 AUROC/AP/F1 comparisons within the three principal models. Holm adjustment (Holm, 1979) is applied separately within each family. The aggregate CSV additionally reports an all-90 Holm sensitivity analysis; separate-family adjustments do not control the familywise error rate over the union. We use $p_{\text{Holm}} < 0.05$, with $*$, $**$, and $***$ denoting adjusted values below 0.05, 0.01, and 0.001, respectively, and ns otherwise.

The 2,000-resample percentile intervals remain unadjusted and are not exact inversions of these tests. We compute p values from the bootstrap differences rather than deriving them from whether an interval includes zero. The seed set, aggregation policy, testing procedure, and reporting families were selected retrospectively after results were available. All inference is exploratory and conditions on these fitted predictors; it does not measure uncertainty from retraining or retrospective selection. OOF-threshold, subgroup, decision-curve, and training-probe summaries receive no additional significance tests.

As a numerical stability sensitivity check retained from the preceding three-seed analysis, we additionally repeat the ForecastAttention–DirectReadout Macro specificity test with 999,999 shared patient resamples and an independent random-number seed, 20261104. The same null-centered statistic is calculated directly from per-patient negative-class contributions. The other 89 test values are held fixed when recalculating the secondary-family and all-90 adjustments for this check. It does not replace the uniformly generated 99,999-resample results in the main tables or alter the reporting families.

5.6. Operating-point and hypothetical decision analyses

As a threshold sensitivity analysis, within each training seed, each development patient's out-of-fold (OOF) probability is supplied only by the fold model for which that patient was in validation. These patients also informed their fold-specific best-checkpoint selection, so the OOF scores are not a nested or independent calibration set. The disjoint validation predictions are concatenated separately for every seed at T0, T1, and T2, and Youden index is maximized on the resulting OOF scores, with the highest threshold resolving ties. The threshold is then applied to holdout ensemble probabilities. The OOF score for a development patient comes from one model, whereas the holdout score averages five models; their score distributions can differ. This analysis therefore examines threshold transfer and does not supply independent validation or calibration of the five-model ensemble. No holdout outcome selects a threshold. Bootstrap intervals hold the OOF-selected threshold fixed.

We additionally compute exploratory decision curves (Vickers and Elkin, 2006). At risk threshold q, the generic positive action is applied when predicted pCR probability is at least q. Its mathematical net benefit is

$$\text{NB}(q) = \frac{\text{TP}(q)}{n} - \frac{\text{FP}(q)}{n}\frac{q}{1-q}. \tag{13}$$

We calculate this separately from each seed's five-fold ensemble probabilities over thresholds 0.05–0.80, average the three landmark net benefits for Macro, and then average across the four seeds. Treat-all and treat-none are references. The calculation uses uncalibrated probabilities and does not validate a particular escalation, de-escalation, or other management action.

5.7. Exploratory subgroup and mechanism analyses

We examine released HR/HER2 combinations, age below or at least 50 years, native source namespace, and

saved predicted-component crop coverage. Age 50 is an exploratory reporting cutpoint. Subgroup AUROC/AP are suppressed when fewer than ten patients or fewer than five outcomes in either class are available. Subgroup metrics are computed within each seed's five-fold ensemble and then averaged across seeds. Paired within-subgroup patient resampling preserves seed, model, and landmark pairing; single-class draws are omitted for AUROC intervals and valid-draw counts retained. Intervals are unadjusted, and no subgroup-interaction hypothesis is tested. Source namespaces share the same clinical release and are not independent external cohorts.

Mechanism analysis is restricted to archived fixed-training-probe diagnostics collected every five epochs. These probes sample up to 32 training patients and are not full validation or holdout forecast evaluations. Unlike the missing-target-masked training objective, the archived forecast diagnostic includes only continuous observed visit chains: the real landmark, target, and every intermediate canonical visit must be available. A gap excludes that landmark–target pair, and eligible-patient counts therefore vary by horizon. We report a descriptive common epoch 20 and common-epoch trajectories. Forecast skill is $1 - \mathrm{L1}_{\mathrm{forecast}}/\mathrm{L1}_{\mathrm{copy}}$ within the same snapshot's EMA representation; the copy forecast holds the observed landmark state constant. This normalization avoids comparing raw latent errors between separately learned teacher coordinates. Predictor gradient norms from the pCR and weighted rollout objectives are descriptive checks of gradient routing, not independent statistical samples or clinical effects. These archived diagnostics are computed within each model's own EMA coordinates, averaged over the five folds within each seed, and then summarized across the four seeds. Only epochs with complete diagnostic coverage are used for common-epoch summaries. They do not ensemble latent states and should be distinguished from the probability-ensemble pCR evaluation.

5.8. Scope of the selected four-seed analysis

The choice of seeds 42, 44, 2026, and 3407 and within-seed probability ensembling was made retrospectively after holdout results from multiple seeds were available. Applying the same seed set, folds, and aggregation rule to every model preserves a matched architectural comparison, but does not remove selection bias. Four-seed standard deviations provide descriptive training variation for this selected subset; the patient-bootstrap intervals and tests condition on the subset. A confirmatory study should lock training seeds, aggregation, model selection, statistical comparisons, and operating-point policies before evaluating independent patients.

6. Results

6.1. Discrimination across four training seeds

All 60 principal-model runs and 20 supplementary mean-pooling runs completed training and formal evaluation. Eighty evaluation receipts and 240 registered report identities were verified. Every seed-specific predictor averaged five fold probabilities for each of the same 175 holdout patients, including 56 positive outcomes. We then calculated metrics within each seed and averaged the metrics across seeds 42, 44, 2026, and 3407. Table 2 reports means with sample seed standard deviations; conditional patient intervals and paired tests are reported separately.

ForecastAttention had mean T0/T1/T2 AUROCs of 0.7438/0.7448/0.7459 and AP values of 0.6188/0.6172/0.6164 (Fig. 2). Its Macro AUROC/AP/F1 was 0.7449/0.6175/0.5596, with seed standard deviations of 0.0040/0.0084/0.0142. Corresponding means were 0.7269/0.5924/0.5484 for NoJEPA and 0.7361/0.6027/0.5517 for DirectReadout. The full model had the highest mean AUROC and AP at every landmark among all four variants and the highest mean Macro F1 among the three principal variants. Its T2 F1 was 0.5621, compared with 0.5552 for DirectReadout and 0.5508 for NoJEPA. These are averages of metrics from four separately evaluated ensembles, not metrics of a combined predictor.

Under direct readout, adding future-state supervision increased mean Macro AUROC/AP/F1 by +0.0092/+0.0103/+0.0033. The complete forecast-attention readout gave further mean increases of +0.0087/+0.0147/+0.0080 relative to DirectReadout. These means conceal variation among the seeds. The direct-versus-NoJEPA Macro AUROC difference was positive at seeds 42, 44, and 3407 but negative at seed 2026 (0.7230 versus 0.7251). Likewise, ForecastAttention exceeded DirectReadout in Macro AUROC at seeds 44, 2026, and 3407 but was slightly lower at seed 42 (0.7419 versus 0.7428). At the newly included seed 3407, Macro AUROC was 0.7348/0.7386/0.7432 for NoJEPA/DirectReadout/ForecastAttention, respectively, but direct-readout F1 was slightly below NoJEPA (0.5654 versus 0.5661). The full-versus-NoJEPA Macro AUROC difference was positive at all four selected seeds. The supplementary seed tables retain these individual results; the subset does not establish uniform behavior across all possible seeds.

The supplementary ForecastMean had mean Macro AUROC/AP/F1 of 0.7436/0.6116/0.5690 (Table A.6). Attention had higher AUROC and AP point estimates, while mean pooling had the highest mean Macro F1 among all four variants. The attention-minus-mean differences were +0.0013 [−0.0085, 0.0114] for AUROC, +0.0059 [−0.0088, 0.0209] for AP, and −0.0093 [−0.0271, 0.0075] for F1. All three unadjusted patient intervals included zero. This comparison provides no uniform attention advantage.

Paired patient intervals qualify the descriptive rankings (Fig. 3). The mean Macro AUROC difference for ForecastAttention versus NoJEPA was +0.0179 [0.0071, 0.0297]. The corresponding DirectReadout–NoJEPA difference was +0.0092 [−0.0028, 0.0216], and the ForecastAttention–DirectReadout difference was +0.0087

Table 2: Performance on the fixed internal holdout (175 patients, 56 pCR positive). Entries are mean ± sample SD over seeds 42, 44, 2026, and 3407. Within each seed, five fold probabilities are averaged per patient before metrics; no probabilities are averaged across seeds. Macro averages T0–T2 metrics within each seed. F1 uses threshold 0.5. Paired Macro differences and adjusted tests are in Table 3.

Model	Landmark	AUROC	AP	F1
NoJEPA	T0	0.7282 ± 0.0054	0.5919 ± 0.0049	0.5434 ± 0.0183
NoJEPA	T1	0.7265 ± 0.0075	0.5927 ± 0.0063	0.5511 ± 0.0205
NoJEPA	T2	0.7261 ± 0.0062	0.5926 ± 0.0044	0.5508 ± 0.0105
NoJEPA	Macro	0.7269 ± 0.0062	0.5924 ± 0.0050	0.5484 ± 0.0160
DirectReadout	T0	0.7348 ± 0.0089	0.5974 ± 0.0064	0.5493 ± 0.0246
DirectReadout	T1	0.7375 ± 0.0108	0.6070 ± 0.0058	0.5505 ± 0.0303
DirectReadout	T2	0.7360 ± 0.0073	0.6038 ± 0.0083	0.5552 ± 0.0101
DirectReadout	Macro	0.7361 ± 0.0089	0.6027 ± 0.0044	0.5517 ± 0.0191
ForecastAttention	T0	0.7438 ± 0.0021	0.6188 ± 0.0078	0.5609 ± 0.0196
ForecastAttention	T1	0.7448 ± 0.0050	0.6172 ± 0.0083	0.5559 ± 0.0155
ForecastAttention	T2	0.7459 ± 0.0061	0.6164 ± 0.0105	0.5621 ± 0.0161
ForecastAttention	Macro	0.7449 ± 0.0040	0.6175 ± 0.0084	0.5596 ± 0.0142




Figure 2: Discrimination across seeds 42, 44, 2026, and 3407. (A–C) Mean ROC curves; (D–F) mean precision–recall curves. Curves are computed from five-fold ensemble probabilities separately within each seed, interpolated on a common grid, and equally averaged across seeds. Shading gives pointwise conditional 95% shared-patient-bootstrap intervals for the mean curve. Colored metric values are means of the directly computed seed-specific AUROC or non-interpolated AP; they need not equal the area under the displayed mean curve. The dotted PR reference is the 0.32 positive prevalence.

[−0.0055, 0.0230]. The full-versus-NoJEPA AP difference was +0.0250 [0.0065, 0.0458], whereas its F1 difference was +0.0112 [−0.0096, 0.0355]. All three full-versus-direct Macro intervals included zero. These intervals concern patient sampling conditional on the selected fitted predictors and are distinct from the descriptive seed standard deviations. The full-versus-NoJEPA Macro AUROC difference had an approximate paired-bootstrap raw $p = 0.00306$ and $p_{\text{Holm}} = 0.02754$ within the nine-comparison principal family. It was the only principal comparison below

the family-adjusted 0.05 criterion. Its adjusted value increased to 0.26622 when all 90 reported tests were considered together. The AP comparison had raw/within-family adjusted $p = 0.01254/0.10032$, and the full-versus-direct AUROC comparison had 0.21898/0.87592. Thus support for the complete-versus-control AUROC difference depends on the reporting family and remains exploratory. Nonsignificant comparisons do not establish equivalence.

6.2. Threshold behavior and class imbalance

At threshold 0.5, mean Macro sensitivity/NPV was 0.6667/0.8100 for ForecastAttention, compared with 0.6190/0.7987 for DirectReadout. These increases accompanied lower specificity (0.6639 versus 0.7073) and PPV (0.4850 versus 0.5010). The paired mean sensitivity difference was +0.0476 [0.0123, 0.0880], while the specificity difference was −0.0434 [−0.0643, −0.0241]. These unadjusted intervals describe an operating trade-off rather than a general classification improvement. The specificity decrease had raw $p = 0.00012$, secondary-family Holm $p = 0.00972$, and all-90 Holm $p = 0.01080$. It was the only secondary comparison meeting its family-adjusted 0.05 criterion and the only comparison meeting the all-90 criterion. The sensitivity increase did not pass its secondary-family criterion ($p_{\text{Holm}} = 0.99075$).

The additional 999,999-resample numerical stability check gave specificity $p = 0.000092$ (Monte Carlo standard error approximately 0.0000096). Holding the other 89 values fixed yielded secondary-family and all-90 adjusted values of 0.007452 and 0.008280, respectively, preserving the adverse-specificity conclusion. Main tables retain the uniformly generated 99,999-resample values.

The full model's seed-mean true-positive counts at T0/T1/T2 were 36.50/37.25/38.25, with false-positive counts 37.50/40.75/41.75. These are averages of integer counts from four correlated predictors, not fractional patients or a single combined confusion matrix. T1 added positive predictions while mean F1 decreased from 0.5609 at T0 to 0.5559; T2 F1 then increased to 0.5621. Later information can change error balance differently from ranking.

Among the principal variants, ForecastAttention had the highest mean Macro balanced accuracy (0.6653), while DirectReadout had slightly higher MCC (0.3127 versus 0.3122) and accuracy (0.6790 versus 0.6648). NoJEPA had balanced accuracy 0.6591, MCC 0.3033, and accuracy 0.6714. Every principal model's mean Macro accuracy was below the all-negative reference of 0.68 in this cohort. The supplementary ForecastMean reached balanced accuracy 0.6735, MCC 0.3268, and accuracy 0.6738. Accuracy, ranking, and class-specific errors therefore give different descriptive orderings.

OOF-selected Youden thresholds changed mean holdout Macro F1 from 0.5484 to 0.5515 for NoJEPA, 0.5517 to 0.5469 for DirectReadout, 0.5596 to 0.5599 for ForecastAttention, and 0.5690 to 0.5631 for ForecastMean (Table A.10). Thus the alternative threshold produced small increases for NoJEPA and ForecastAttention but decreases for the other variants. These results concern transfer of thresholds fitted on single-model OOF scores to within-seed five-model ensemble scores; they neither optimize a holdout threshold nor establish independent ensemble calibration.

6.3. Longitudinal information and probability quality

The paired seed-mean T2–T0 AUROC change was +0.0021 [−0.0030, 0.0079] for ForecastAttention, −0.0022 [−0.0070, 0.0021] for NoJEPA, and +0.0012 [−0.0034, 0.0058] for DirectReadout. The full model's corresponding AP change was −0.0025 [−0.0128, 0.0079] and F1 change +0.0011 [−0.0152, 0.0173]. All these longitudinal intervals included zero. The increasing mean AUROC for forecast attention across landmarks therefore remains a small descriptive trend, accompanied by slightly decreasing mean AP. Fig. A.8 shows outcome-conditioned distributions of changes in seed-specific ensemble probabilities.

Mean Macro NLL/Brier/ECE was 0.5900/0.2033/0.1469 for ForecastAttention, compared with 0.5817/0.2003/0.1288 for DirectReadout and 0.5898/0.2032/0.1299 for NoJEPA. Forecast attention had worse point estimates of all three probability-quality metrics than both direct-readout controls, although its NLL and Brier differences from NoJEPA were small. The better ranking means therefore did not translate into better probability-quality means. All 14 Macro metrics and their conditional patient intervals appear in Table A.9.

6.4. Hypothetical net benefit

Decision curves did not preserve the discrimination ranking across thresholds. At a hypothetical threshold of 0.30, mean Macro net benefit was 0.1163 for DirectReadout and 0.1031 for ForecastAttention. At 0.50, it was −0.0086 for NoJEPA, −0.0010 for DirectReadout, and −0.0152 for ForecastAttention. At 0.70 the ordering reversed, with +0.0002 for ForecastAttention and −0.0333 for DirectReadout. These values average seed-specific decision curves, use uncalibrated probabilities, and assume a generic positive action. They do not establish an acceptable clinical threshold or validate a management pathway.

6.5. Exploratory biological and acquisition-related subgroups

The four HR/HER2 groups had positive seed-mean Macro AUROC differences between ForecastAttention and NoJEPA (Fig. 5; Table B.15). The HR-positive/HER2-negative group had a difference of +0.0257 [−0.0010, 0.0556] among 68 patients, including 11 positive outcomes. Other differences were +0.0113 [−0.0457, 0.0730] in HR-positive/HER2-positive patients (31, 12 positive), +0.0425 [−0.0187, 0.1320] in HR-negative/HER2-positive patients (14, 7 positive), and +0.0219 [−0.0057,

Table 3: Exploratory principal Macro comparisons. All metrics use approximate, null-centered paired patient-bootstrap tests with 99,999 shared resamples. Within each draw, metrics are computed for each seed then equally averaged. All p values are two-sided. Holm_9 adjusts the nine entries here; Holm_{90} is the sensitivity analysis over all principal and secondary tests. Sig. refers to Holm_9. Direct and Attention abbreviate DirectReadout and ForecastAttention. $^{*}p_{\text{Holm}} < 0.05$, $^{**}p_{\text{Holm}} < 0.01$, $^{***}p_{\text{Holm}} < 0.001$; ns, not significant after the stated correction. Intervals are unadjusted 95% patient-bootstrap intervals (2,000 shared draws) and need not agree with adjusted test decisions. All analyses condition on the fixed, retrospectively selected seed42/44/2026/3407 ensembles; seeds are not resampled.

Comparison	Metric	Difference [95% CI]	p	$p_{\text{Holm},9}$	$p_{\text{Holm},90}$	Sig.
Direct – NoJEPA	AUROC	+0.0092 [-0.0028, +0.0216]	0.1273	0.7638	1.0000	ns
Direct – NoJEPA	AP	+0.0103 [-0.0046, +0.0255]	0.1721	0.8603	1.0000	ns
Direct – NoJEPA	F1	+0.0033 [-0.0135, +0.0184]	0.6713	1.0000	1.0000	ns
Attention – NoJEPA	AUROC	+0.0179 [+0.0071, +0.0297]	0.0031	0.0275	0.2662	*
Attention – NoJEPA	AP	+0.0250 [+0.0065, +0.0458]	0.0125	0.1003	1.0000	ns
Attention – NoJEPA	F1	+0.0112 [-0.0096, +0.0355]	0.3223	0.9668	1.0000	ns
Attention – Direct	AUROC	+0.0087 [-0.0055, +0.0230]	0.2190	0.8759	1.0000	ns
Attention – Direct	AP	+0.0147 [-0.0014, +0.0332]	0.0951	0.6654	1.0000	ns
Attention – Direct	F1	+0.0080 [-0.0165, +0.0373]	0.5598	1.0000	1.0000	ns




Figure 3: Paired seed-mean metric differences at T0, T1, T2, and Macro. Colors distinguish auxiliary supervision, additional forecast readout, and the complete-model contrast. Squares denote Macro estimates. Bars are unadjusted 95% patient-bootstrap intervals. Every draw is shared across seeds, models, and landmarks; metrics are computed within seed before averaging. Adjusted tests appear in Tables 3 and A.11. Folds, seeds, and repeated landmarks do not create independent test cohorts.

0.0516] in HR-negative/HER2-negative patients (62, 26 positive). All four unadjusted intervals included zero. These summaries do not establish an interaction or a confirmed subtype-specific benefit.

In the HR-positive/HER2-negative group, forecast attention's mean Macro sensitivity/specificity was 0.1591/0.9649, compared with 0.1364/0.9810 for NoJEPA. Thus low sensitivity at threshold 0.5 persisted despite the positive AUROC difference. DirectReadout had sensitivity/specificity 0.1212/0.9825 and a slightly higher mean AUROC than forecast attention (0.7125 versus 0.7087). The age groups also had positive full-versus-control differences, but their unadjusted exploratory intervals do not support an age-interaction conclusion. Source namespaces share one clinical release and do not constitute external validation. The partially clipped baseline-component group contained only 11 patients and one positive outcome; its AUROC and AP were suppressed under the minimum-outcome rule.

Table 4: Operating characteristics at threshold 0.5. Entries average the four seed-specific fivefold probability-ensemble metrics; Macro first averages T0–T2 within each seed. The same 175 patients appear in every evaluation.

Model	Landmark	Sens.	Spec.	PPV	NPV	BA	MCC
NoJEPA	T0	0.607	0.706	0.493	0.793	0.657	0.299
NoJEPA	T1	0.621	0.704	0.497	0.798	0.662	0.309
NoJEPA	T2	0.647	0.670	0.483	0.803	0.659	0.301
NoJEPA	Macro	0.625	0.693	0.491	0.798	0.659	0.303
DirectReadout	T0	0.598	0.729	0.511	0.795	0.664	0.317
DirectReadout	T1	0.607	0.721	0.507	0.797	0.664	0.316
DirectReadout	T2	0.652	0.672	0.485	0.804	0.662	0.306
DirectReadout	Macro	0.619	0.707	0.501	0.799	0.663	0.313
ForecastAttention	T0	0.652	0.685	0.493	0.808	0.668	0.318
ForecastAttention	T1	0.665	0.658	0.481	0.808	0.661	0.305
ForecastAttention	T2	0.683	0.649	0.481	0.815	0.666	0.313
ForecastAttention	Macro	0.667	0.664	0.485	0.810	0.665	0.312

Table 5: Probability quality: four-seed mean ± sample SD of Macro metrics. Each seed has its own fivefold probability ensemble. Lower is better; ECE uses ten equal-width bins.

Model	NLL	Brier	ECE
NoJEPA	0.5898 ± 0.0133	0.2032 ± 0.0054	0.1299 ± 0.0279
DirectReadout	0.5817 ± 0.0149	0.2003 ± 0.0068	0.1288 ± 0.0250
ForecastAttention	0.5900 ± 0.0194	0.2033 ± 0.0082	0.1469 ± 0.0342

6.6. Archived forecast and gradient diagnostics

Archived fixed-training-probe diagnostics describe individual models separately from the pCR holdout analysis (Fig. 6). At common epoch 20, mean copy-normalized forecast skills for DirectReadout were +0.0090, +0.0524, and +0.1590 at one-, two-, and three-step horizons. Corresponding values for ForecastAttention were −0.0509, −0.0516, and +0.0122 (Table B.16). Statistics were normalized within each snapshot's EMA coordinates, averaged across five folds within seed, and then averaged across the four seeds. Separate models' latent states were not ensembled. These diagnostics do not support attributing forecast attention's pCR ranking to uniformly more accurate latent forecasts.

The probe requires the real landmark, target, and every intermediate examination, whereas the training objective masks unavailable future targets. Eligible populations therefore vary by horizon; only 5–9 patients per probe were eligible at three steps. These measurements are not selected-checkpoint holdout forecast evaluations. The predictor's pCR-loss gradient norm was zero for DirectReadout and had mean 1.0123 for ForecastAttention; weighted rollout-gradient means were 0.1570 and 0.1662. These values support the expected gradient routing, but gradient magnitude is not a biological effect size. NoJEPA had no available trained-world diagnostic and was not assigned an artificial zero skill.

7. Discussion

7.1. Separating auxiliary supervision from forecast readout

Across the four selected seeds, auxiliary supervision increased the direct-readout mean Macro AUROC and AP, and the complete forecast-attention readout increased these means further. Forecast attention also had the highest mean Macro F1 among the principal variants. However, the auxiliary-supervision comparison changed sign in seed 2026, and the incremental full-versus-direct AUROC comparison changed sign in seed 42. These individual results remain visible alongside the summaries. A higher mean does not imply a favorable effect in every training realization.

The complete-versus-NoJEPA Macro AUROC comparison met the nine-test principal-family criterion ($p_{\text{Holm}} = 0.02754$), but not the all-90 sensitivity criterion (0.26622). The AP and F1 differences did not meet the principal-family criterion. This conditional evidence concerns the selected fitted models and reporting family; it does not remove retrospective selection bias or establish an independently confirmed architectural benefit. The supplementary mean-pooling control had slightly lower mean Macro AUROC and AP, but higher mean F1. Its paired AUROC difference from attention included zero. Attention is therefore not established as necessary for the observed ranking performance.

The DirectReadout–NoJEPA contrast most closely isolates future-state supervision. By comparison, the full-versus-direct contrast changes forecast input features, pooling, and classifier width together. Classifier-width-matched




Figure 4: Probability quality and hypothetical decision utility across four seeds. (A–C) Reliability summaries first form five equal-count probability-rank groups of 35 patients within each seed-specific five-fold ensemble and then average the group summaries across seeds. They differ from the ten equal-width bins used for ECE. (D–F) Net-benefit curves are computed separately within seeds and then averaged, with treat-all and treat-none references. Scores are not recalibrated. No clinical escalation or de-escalation strategy is validated.

direct readout, stopped-gradient forecast features, copy-state readout, and independent holdout forecast evaluation would help distinguish these contributions. The main models predict embeddings rather than generate MRI for classification. The internal name NoJEPA denotes removal of the future-state objective while retaining the shared fusion network and transition queries.

7.2. What additional visits change

The full model's mean AUROC increased modestly across T0–T2, but its paired T2–T0 interval included zero. Its mean AP decreased slightly, while F1 decreased at T1 before increasing at T2. Changes in true- and false-positive predictions explain why added observations, ranking, and a fixed-threshold decision rule need not improve together. Outcome-conditioned histograms summarize probability changes within each seed-specific ensemble; they do not demonstrate that every patient's prediction improves or that observed score changes are treatment effects.

T0 imaging preceded randomization, while the model is conditioned on the eventual assigned arm. This remains a retrospective conditional scenario. The arm encoding does not capture delivered dose, cycle timing, dose reductions, adherence, or treatment changes. Altering the arm token cannot identify an individual treatment effect without suitable causal assumptions and exposure data. Complete T0–T2 holdout eligibility also restricts the population to which the longitudinal comparison applies.

7.3. Classification, probability quality, and hypothetical utility

The additional metrics expose trade-offs behind the ranking means. Compared with auxiliary-supervised direct readout, forecast attention had higher mean sensitivity and NPV but lower specificity, PPV, and accuracy; mean MCC was almost equal. The specificity decrease passed both the 81-test secondary-family correction (0.00972) and the all-90 correction (0.01080), whereas the sensitivity increase did not pass its secondary-family criterion. Evidence for this adverse operating-point difference should accompany the favorable discrimination summary.

Forecast attention also had worse NLL, Brier score, and ECE point estimates than both direct-readout controls, although its NLL and Brier differences from NoJEPA were small. Its higher ranking means therefore do not imply better predicted probabilities. The OOF Youden




Figure 5: Exploratory subgroup and operating-point summaries, with metrics computed within seed and then equally averaged across seeds 42, 44, 2026, and 3407. (A) Macro AUROC differences between ForecastAttention and NoJEPA within HR/HER2 and age groups, with patient counts, positive outcomes, and unadjusted shared-patient intervals. No subgroup-interaction claim is made. (B) Mean specificity and sensitivity at threshold 0.5 across landmarks; colors identify models and markers identify landmarks.




Figure 6: Archived fixed-training-probe diagnostics across four seeds. (A) Copy-normalized forecast skill at common epoch 20. (B) Predictor gradient norms from pCR and weighted rollout losses. (C) Three-step skill across epochs with complete diagnostic coverage. Statistics are calculated within each snapshot's EMA coordinates, averaged across folds within seed, and summarized across seeds. Error bars in A and B and shading in C show the descriptive sample SD of the four seed-specific fold means. These are training diagnostics, distinct from pCR probability ensembling; real continuous visit chains are required, and three-step eligibility is only 5–9 patients per probe.

analysis slightly increased F1 for NoJEPA and forecast attention but decreased it for the other variants. Development OOF scores originate from one held-out fold model per patient, whereas test scores average five models within a seed. Transferring thresholds between these score distributions is a sensitivity analysis rather than independent ensemble calibration.

Net-benefit orderings changed with the hypothetical threshold: direct readout was higher at 0.30 and 0.50, while forecast attention was higher at 0.70. These are seed means under a generic positive-action utility, not evaluations of a defined care pathway. A clinically motivated threshold requires consequences for false-positive and false-negative actions to be specified and calibration to be fitted without holdout leakage.

7.4. Seed variation, subgroups, and forecasting diagnostics

The mean and sample standard deviation across four seeds describe four training realizations using identical development folds and holdout patients. They do not turn seeds into independent clinical replications. Patient-bootstrap intervals and tests answer a different question: how the mean metric of these fixed predictors varies under patient resampling. Because seeds are not resampled and models are not refitted, those intervals do not estimate uncertainty over the training algorithm or future seed choices.

Subgroup summaries are exploratory, use small denominators, and do not test subtype interactions. All four HR/HER2 full-versus-control Macro AUROC intervals included zero. The positive point estimates do not establish subtype-specific benefit; low sensitivity at 0.5 in the HR-positive/HER2-negative subgroup also persists. Source namespaces share one clinical release and do not provide external validation. Crop analysis is similarly constrained: the 11 patients with partial clipping contain only one positive outcome, preventing reliable discrimination assessment under the minimum-outcome rule. Reporting the denominator and suppressing AUROC/AP is preferable to interpreting an unstable estimate.

Archived fixed-training-probe diagnostics are separate from the pCR holdout analysis. Their copy-normalized skills and gradient norms are calculated in each snapshot's own EMA representation, averaged over folds within seed, and then summarized across seeds. They do not ensemble latent states or establish anatomical forecast fidelity. At epoch 20, auxiliary-supervised direct models had higher mean copy-normalized skill than forecast attention at all three horizons, despite the latter's higher mean pCR AUROC. This weakens an explanation based solely on more faithful forecasts. Gradient routing can support an implementation-level mechanism, but gradient magnitude is not a biological effect size. Independent forecast evaluation at the selected pCR checkpoints remains necessary.

7.5. Limitations and a confirmatory study

The seed set 42, 44, 2026, and 3407, within-seed probability ensembling, and reporting procedures were selected after holdout results were available. Patient-bootstrap inference conditions on these fitted ensembles and fixed OOF thresholds. It excludes retraining, seed-set selection, aggregation choice, model selection, and threshold-selection uncertainty. Four selected seeds do not establish broad training stability, and their sample standard deviations have limited precision. The reported mean describes four five-model predictors; it is not the performance of one twenty-model ensemble or a model trained on the full development cohort.

The 95% patient intervals are unadjusted. Holm corrections for nine principal and 81 secondary tests, together with the all-90 sensitivity analysis, do not undo retrospective choices. Null-centered bootstrap tests are approximate, with finite-sample limitations particularly for non-smooth statistics such as AP, F1, and binned ECE. The differing principal-family and all-90 conclusions should remain visible. Nonsignificance does not establish equivalence. A confirmatory study should lock seeds, aggregation, model and threshold selection, and statistical comparisons before evaluating independent patients.

The holdout is an internal downstream split. Twenty-nine patients had belonged to an earlier development split. Current supervised models were retrained using the new patient-disjoint folds and were not warm-started from those earlier pCR models, but codec and localization-model pretraining have not been fully audited for current-holdout overlap. This prevents a claim that every pipeline stage was evaluated on wholly unseen patients.

Fixed baseline localization and cropping may omit multifocal disease or changes outside the field of view. Geometry and array checks do not establish anatomical registration accuracy. Holdout patients have complete T0–T2 imaging, whereas development retains incomplete histories; missingness, acquisition variation, and subtype prevalence may limit transportability. Archived training probes do not replace best-checkpoint holdout forecast evaluation and have small long-horizon denominators.

The study remains an internal architectural comparison. Matched clinical-only, conventional imaging, and independent longitudinal baselines are still needed, along with external validation, anatomical quality review, and independently assessed calibration. Precise pCR pathology criteria, scan intervals, ethics determination, and upstream provenance also require final documentation before submission.

8. Conclusion

Across training seeds 42, 44, 2026, and 3407, forecast-attention readout had the highest mean AUROC and AP among all four variants and the highest mean Macro F1 among the principal variants. The supplementary mean-pooling control had similar mean Macro AUROC and higher F1. The complete-versus-NoJEPA Macro AUROC difference met the nine-test principal-family criterion but not the all-90 correction; lower specificity than auxiliary-supervised direct readout remained detectable under both secondary-family and all-90 correction. Probability-quality and hypothetical-utility summaries showed further trade-offs. These exploratory results motivate independent validation, complete upstream provenance, stronger matched baselines, and prospectively fixed training and statistical policies.

Data and code availability

Source imaging collections are available through The Cancer Imaging Archive, subject to their applicable access conditions and licenses. The present derived manifests,

processing lineage, and evaluation outputs are retained locally. [TO COMPLETE: Specify the public repository, versioned release, permitted derived-data artifacts, and access mechanism before submission; no public code-release claim is made in this draft.]

Ethics approval and consent

[TO COMPLETE: Provide the original studies' ethics and consent statements and the local institutional determination for this secondary analysis, including an approval or exemption identifier where applicable. Public availability alone is not an ethics approval.]

Funding

[TO COMPLETE: Provide grant numbers and the funders' roles, or confirm that no specific funding was received.]

Declaration of competing interest

[TO COMPLETE: Obtain and insert the actual declarations from every author.]

CRediT authorship contribution statement

[TO COMPLETE: Provide author names and agreed contributions; no authorship or contributions have been inferred.]

Declaration of generative AI and AI-assisted technologies in the manuscript preparation process

During preparation of this initial draft, OpenAI Codex was used to organize source evidence, generate reproducible aggregate tables and figures, and assist with English drafting. [TO COMPLETE: All authors must review and edit the manuscript, verify every statement and reference, and finalize the disclosure under the journal's current policy before submission.]

Appendix A. Supplementary pooling, metrics, and uncertainty

The supplementary ForecastMean uses seeds 42, 44, 2026, and 3407, the same five development folds, observed-history protocol, future-state objective, and forecast feature groups as ForecastAttention, replacing spatial attention with mean pooling. Metrics are computed from patient-wise five-fold probability means within each seed and then averaged across the four seeds. The per-seed results in Tables A.7 and A.8 preserve the individual training realizations underlying the reported mean and sample standard deviation.

Appendix A.1. Paired significance checks

Table 3 reports the nine principal Macro comparisons, with unadjusted 95% paired patient-bootstrap intervals, raw two-sided approximate bootstrap p values, within-family Holm adjustment, and all-90 Holm adjustment. The secondary family comprises 81 tests in Tables A.11–A.14. Every patient draw is shared across all four fixed seeds, models, and landmarks; metrics are computed within seeds before averaging. The independent patient count remains 175. A significance marker refers to the stated Holm-adjusted family; ns denotes $p_{\text{Holm}} \geq 0.05$. The existing 2,000-resample percentile intervals are unadjusted and are not exact inversions of the 99,999-resample null-centered tests. All testing and reporting choices remain retrospective and exploratory.

Appendix B. Supplementary subgroup and training-probe summaries

Appendix C. Reproducibility and reporting items to complete

The source package includes aggregate CSV files, source-report hashes, reproducible metric and figure scripts, the shared-patient-bootstrap procedure, and the official article class. The significance CSV includes raw p values, within-family Holm adjustment, and adjustment over all 90 tests as a sensitivity analysis. It contains no patient identifiers, patient-level prediction rows, original MRI volumes, or model checkpoints. The method figure includes only rendered, anonymous development-image slices. Every variant uses training seeds 42, 44, 2026, and 3407. Five fold probabilities are averaged per patient within each seed before computing pCR metrics, and seed-specific metric values are then equally averaged. Individual-fold metrics are retained for numerical audit; archived training diagnostics are summarized across folds within seed and then across seeds in their own latent coordinates.

[TO COMPLETE: Before submission, complete author and ethics statements; verify the precise pCR pathology definition; document codec and segmentation patient-level pretraining provenance; add matched external baselines and prospectively locked training and analysis policies; audit external-cohort independence; and confirm the journal's current submission and anonymization requirements.]

Table A.6: Supplementary spatial pooling comparison, mean ± sample SD over seeds 42,44,2026,3407. Within-seed fivefold probabilities are averaged before metrics, then seed metrics are averaged. ForecastMean and ForecastAttention share forecast readout and future-state supervision.

Model	Landmark	AUROC	AP	F1
ForecastMean	T0	0.7432 ± 0.0050	0.6086 ± 0.0077	0.5618 ± 0.0124
ForecastMean	T1	0.7446 ± 0.0056	0.6126 ± 0.0033	0.5752 ± 0.0147
ForecastMean	T2	0.7430 ± 0.0052	0.6136 ± 0.0069	0.5699 ± 0.0032
ForecastMean	Macro	0.7436 ± 0.0050	0.6116 ± 0.0050	0.5690 ± 0.0078
ForecastAttention	T0	0.7438 ± 0.0021	0.6188 ± 0.0078	0.5609 ± 0.0196
ForecastAttention	T1	0.7448 ± 0.0050	0.6172 ± 0.0083	0.5559 ± 0.0155
ForecastAttention	T2	0.7459 ± 0.0061	0.6164 ± 0.0105	0.5621 ± 0.0161
ForecastAttention	Macro	0.7449 ± 0.0040	0.6175 ± 0.0084	0.5596 ± 0.0142

Table A.7: Individual-seed results: patient-level fivefold probability ensembles, each evaluated on the same 175 patients (56 positive). Macro is the mean of three landmark metrics. These seed-specific estimates generate the main table mean and sample SD; they are not independent patient cohorts.

Model	Seed	Landmark	AUROC	AP	F1
NoJEPA	42	T0	0.7216	0.5945	0.5217
NoJEPA	42	T1	0.7176	0.5966	0.5299
NoJEPA	42	T2	0.7206	0.5957	0.5378
NoJEPA	42	Macro	0.7199	0.5956	0.5298
NoJEPA	44	T0	0.7275	0.5954	0.5354
NoJEPA	44	T1	0.7281	0.5981	0.5426
NoJEPA	44	T2	0.7281	0.5971	0.5469
NoJEPA	44	Macro	0.7279	0.5968	0.5416
NoJEPA	2026	T0	0.7291	0.5933	0.5538
NoJEPA	2026	T1	0.7246	0.5921	0.5538
NoJEPA	2026	T2	0.7216	0.5894	0.5606
NoJEPA	2026	Macro	0.7251	0.5916	0.5561
NoJEPA	3407	T0	0.7347	0.5846	0.5625
NoJEPA	3407	T1	0.7357	0.5842	0.5781
NoJEPA	3407	T2	0.7339	0.5882	0.5578
NoJEPA	3407	Macro	0.7348	0.5857	0.5661
DirectReadout	42	T0	0.7435	0.5881	0.5321
DirectReadout	42	T1	0.7461	0.6119	0.5487
DirectReadout	42	T2	0.7389	0.5914	0.5600
DirectReadout	42	Macro	0.7428	0.5971	0.5469
DirectReadout	44	T0	0.7372	0.6025	0.5669
DirectReadout	44	T1	0.7416	0.6115	0.5692
DirectReadout	44	T2	0.7410	0.6086	0.5672
DirectReadout	44	Macro	0.7399	0.6075	0.5678
DirectReadout	2026	T0	0.7224	0.5991	0.5246
DirectReadout	2026	T1	0.7216	0.5997	0.5085
DirectReadout	2026	T2	0.7251	0.6069	0.5469
DirectReadout	2026	Macro	0.7230	0.6019	0.5266
DirectReadout	3407	T0	0.7362	0.6000	0.5736
DirectReadout	3407	T1	0.7407	0.6049	0.5758
DirectReadout	3407	T2	0.7390	0.6084	0.5468
DirectReadout	3407	Macro	0.7386	0.6044	0.5654

Table A.8: Individual-seed results: patient-level fivefold probability ensembles, each evaluated on the same 175 patients (56 positive). Macro is the mean of three landmark metrics. These seed-specific estimates generate the main table mean and sample SD; they are not independent patient cohorts.

Model	Seed	Landmark	AUROC	AP	F1
ForecastAttention	42	T0	0.7408	0.6130	0.5366
ForecastAttention	42	T1	0.7423	0.6156	0.5410
ForecastAttention	42	T2	0.7426	0.6115	0.5410
ForecastAttention	42	Macro	0.7419	0.6134	0.5395
ForecastAttention	44	T0	0.7450	0.6256	0.5736
ForecastAttention	44	T1	0.7521	0.6278	0.5442
ForecastAttention	44	T2	0.7550	0.6315	0.5634
ForecastAttention	44	Macro	0.7507	0.6283	0.5604
ForecastAttention	2026	T0	0.7453	0.6111	0.5538
ForecastAttention	2026	T1	0.7437	0.6077	0.5669
ForecastAttention	2026	T2	0.7417	0.6079	0.5802
ForecastAttention	2026	Macro	0.7436	0.6089	0.5670
ForecastAttention	3407	T0	0.7441	0.6256	0.5797
ForecastAttention	3407	T1	0.7410	0.6175	0.5714
ForecastAttention	3407	T2	0.7444	0.6146	0.5638
ForecastAttention	3407	Macro	0.7432	0.6192	0.5716
ForecastMean	42	T0	0.7425	0.6188	0.5714
ForecastMean	42	T1	0.7407	0.6170	0.5891
ForecastMean	42	T2	0.7389	0.6168	0.5672
ForecastMean	42	Macro	0.7407	0.6175	0.5759
ForecastMean	44	T0	0.7431	0.6088	0.5625
ForecastMean	44	T1	0.7494	0.6121	0.5865
ForecastMean	44	T2	0.7471	0.6209	0.5736
ForecastMean	44	Macro	0.7465	0.6139	0.5742
ForecastMean	2026	T0	0.7497	0.6002	0.5692
ForecastMean	2026	T1	0.7495	0.6120	0.5600
ForecastMean	2026	T2	0.7477	0.6121	0.5714
ForecastMean	2026	Macro	0.7490	0.6081	0.5669
ForecastMean	3407	T0	0.7374	0.6066	0.5441
ForecastMean	3407	T1	0.7389	0.6091	0.5652
ForecastMean	3407	T2	0.7381	0.6049	0.5674
ForecastMean	3407	Macro	0.7381	0.6069	0.5589

Table A.9: All 14 macro metrics for the principal variants: point estimate [95% patient-bootstrap percentile interval]. Within each of seeds 42,44,2026,3407, five fold probabilities are averaged per patient, metrics are computed and averaged over T0–T2, then seed metrics are equally averaged. Threshold-dependent metrics use 0.5. Identical patient resamples preserve pairing across models and landmarks. Entries are marginal model intervals; paired differences are reported separately. Intervals condition on the fitted ensembles and exclude retraining and retrospective selection uncertainty. Downward arrows identify metrics for which lower is better.

Metric	NoJEPA	DirectReadout	ForecastAttention
AUROC	0.7269 [0.6425, 0.8014]	0.7361 [0.6523, 0.8087]	0.7449 [0.6618, 0.8155]
AP	0.5924 [0.4710, 0.7069]	0.6027 [0.4805, 0.7175]	0.6175 [0.4933, 0.7309]
F1	0.5484 [0.4443, 0.6406]	0.5517 [0.4496, 0.6440]	0.5596 [0.4576, 0.6468]
Accuracy	0.6714 [0.6052, 0.7338]	0.6790 [0.6148, 0.7405]	0.6648 [0.5967, 0.7267]
Sensitivity	0.6250 [0.5077, 0.7417]	0.6190 [0.5042, 0.7346]	0.6667 [0.5489, 0.7798]
Specificity	0.6933 [0.6126, 0.7689]	0.7073 [0.6318, 0.7800]	0.6639 [0.5804, 0.7431]
PPV	0.4911 [0.3786, 0.5994]	0.5010 [0.3879, 0.6118]	0.4850 [0.3791, 0.5890]
NPV	0.7980 [0.7179, 0.8718]	0.7987 [0.7188, 0.8716]	0.8100 [0.7303, 0.8821]
Balanced accuracy	0.6591 [0.5872, 0.7303]	0.6632 [0.5918, 0.7317]	0.6653 [0.5944, 0.7321]
MCC	0.3033 [0.1641, 0.4374]	0.3127 [0.1736, 0.4458]	0.3122 [0.1768, 0.4398]
Youden index	0.3183 [0.1744, 0.4607]	0.3263 [0.1835, 0.4635]	0.3305 [0.1887, 0.4642]
NLL ↓	0.5898 [0.5368, 0.6462]	0.5817 [0.5250, 0.6451]	0.5900 [0.5407, 0.6433]
Brier score ↓	0.2032 [0.1795, 0.2288]	0.2003 [0.1748, 0.2279]	0.2033 [0.1813, 0.2274]
ECE ↓	0.1299 [0.0984, 0.1959]	0.1288 [0.1017, 0.1924]	0.1469 [0.1094, 0.2101]

Table A.10: Threshold sensitivity analysis: four-seed mean Macro metrics. Each seed uses landmark-specific Youden thresholds from its disjoint validation-fold scores. OOF single-model scores differ from test fivefold ensemble scores; this is sensitivity analysis, not independent ensemble calibration. No test labels choose thresholds.

Model	Policy	F1	Sens.	Spec.	MCC
NoJEPA	0.5	0.548	0.625	0.693	0.303
NoJEPA	OOF validation Youden	0.551	0.638	0.683	0.305
DirectReadout	0.5	0.552	0.619	0.707	0.313
DirectReadout	OOF validation Youden	0.547	0.607	0.713	0.308
ForecastAttention	0.5	0.560	0.667	0.664	0.312
ForecastAttention	OOF validation Youden	0.560	0.662	0.670	0.314

Table A.11: Exploratory landmark comparisons. All metrics use approximate, null-centered paired patient-bootstrap tests (99,999 shared draws across seeds, models, and landmarks). All p values are two-sided. p_{Holm} adjusts jointly over the 81 secondary tests in Tables A.11–A.14. Direct and Attention abbreviate DirectReadout and ForecastAttention. $^{*}p_{\text{Holm}} < 0.05$, $^{**}p_{\text{Holm}} < 0.01$, $^{***}p_{\text{Holm}} < 0.001$; ns, not significant after the stated correction. Intervals are unadjusted 95% patient-bootstrap intervals (2,000 shared draws) and need not agree with adjusted test decisions. All analyses condition on the fixed, retrospectively selected seed42/44/2026/3407 ensembles; seeds are not resampled.

Comparison	Landmark	Metric	Difference [95% CI]	p	p_{Holm}	Sig.
Direct – NoJEPA	T0	AUROC	+0.0066 [-0.0052, +0.0186]	0.2672	1.0000	ns
Direct – NoJEPA	T0	AP	+0.0055 [-0.0109, +0.0217]	0.4901	1.0000	ns
Direct – NoJEPA	T0	F1	+0.0059 [-0.0197, +0.0288]	0.6153	1.0000	ns
Direct – NoJEPA	T1	AUROC	+0.0110 [-0.0016, +0.0234]	0.0883	1.0000	ns
Direct – NoJEPA	T1	AP	+0.0142 [-0.0035, +0.0325]	0.1187	1.0000	ns
Direct – NoJEPA	T1	F1	-0.0006 [-0.0169, +0.0145]	0.9405	1.0000	ns
Direct – NoJEPA	T2	AUROC	+0.0099 [-0.0022, +0.0233]	0.1151	1.0000	ns
Direct – NoJEPA	T2	AP	+0.0112 [-0.0023, +0.0277]	0.1443	1.0000	ns
Direct – NoJEPA	T2	F1	+0.0044 [-0.0171, +0.0259]	0.6796	1.0000	ns
Attention – NoJEPA	T0	AUROC	+0.0156 [+0.0039, +0.0282]	0.0159	1.0000	ns
Attention – NoJEPA	T0	AP	+0.0269 [+0.0072, +0.0486]	0.0123	0.9310	ns
Attention – NoJEPA	T0	F1	+0.0176 [-0.0091, +0.0479]	0.2247	1.0000	ns
Attention – NoJEPA	T1	AUROC	+0.0183 [+0.0074, +0.0298]	0.0024	0.1904	ns
Attention – NoJEPA	T1	AP	+0.0244 [+0.0047, +0.0459]	0.0223	1.0000	ns
Attention – NoJEPA	T1	F1	+0.0048 [-0.0167, +0.0298]	0.6818	1.0000	ns
Attention – NoJEPA	T2	AUROC	+0.0199 [+0.0087, +0.0323]	0.0021	0.1664	ns
Attention – NoJEPA	T2	AP	+0.0238 [+0.0040, +0.0456]	0.0240	1.0000	ns
Attention – NoJEPA	T2	F1	+0.0113 [-0.0098, +0.0341]	0.3156	1.0000	ns
Attention – Direct	T0	AUROC	+0.0090 [-0.0067, +0.0251]	0.2538	1.0000	ns
Attention – Direct	T0	AP	+0.0214 [+0.0008, +0.0420]	0.0369	1.0000	ns
Attention – Direct	T0	F1	+0.0116 [-0.0231, +0.0511]	0.5278	1.0000	ns
Attention – Direct	T1	AUROC	+0.0073 [-0.0062, +0.0211]	0.2851	1.0000	ns
Attention – Direct	T1	AP	+0.0102 [-0.0058, +0.0277]	0.2399	1.0000	ns
Attention – Direct	T1	F1	+0.0054 [-0.0216, +0.0371]	0.7102	1.0000	ns
Attention – Direct	T2	AUROC	+0.0099 [-0.0038, +0.0249]	0.1721	1.0000	ns
Attention – Direct	T2	AP	+0.0126 [-0.0066, +0.0337]	0.2178	1.0000	ns
Attention – Direct	T2	F1	+0.0069 [-0.0163, +0.0352]	0.5964	1.0000	ns

Table A.12: Exploratory Macro operating and probability-quality comparisons. Differences are first-named minus second-named model; negative Brier, NLL, or ECE differences favor the first model. All metrics use approximate, null-centered paired patient-bootstrap tests (99,999 shared draws across seeds, models, and landmarks). All p values are two-sided. p_{Holm} adjusts jointly over the 81 secondary tests in Tables A.11–A.14. Direct and Attention abbreviate DirectReadout and ForecastAttention. $^{*}p_{\text{Holm}} < 0.05$, $^{**}p_{\text{Holm}} < 0.01$, $^{***}p_{\text{Holm}} < 0.001$; ns, not significant after the stated correction. Intervals are unadjusted 95% patient-bootstrap intervals (2,000 shared draws) and need not agree with adjusted test decisions. All analyses condition on the fixed, retrospectively selected seed42/44/2026/3407 ensembles; seeds are not resampled.

Comparison	Metric	Difference [95% CI]	p	p_{Holm}	Sig.
Direct – NoJEPA	Accuracy	+0.0076 [-0.0048, +0.0205]	0.2365	1.0000	ns
Direct – NoJEPA	Sensitivity	-0.0060 [-0.0278, +0.0133]	0.5834	1.0000	ns
Direct – NoJEPA	Specificity	+0.0140 [-0.0007, +0.0304]	0.0765	1.0000	ns
Direct – NoJEPA	PPV	+0.0099 [-0.0058, +0.0259]	0.2026	1.0000	ns
Direct – NoJEPA	NPV	+0.0008 [-0.0093, +0.0095]	0.8704	1.0000	ns
Direct – NoJEPA	Balanced accuracy	+0.0040 [-0.0096, +0.0163]	0.5269	1.0000	ns
Direct – NoJEPA	MCC	+0.0094 [-0.0160, +0.0332]	0.4423	1.0000	ns
Direct – NoJEPA	Youden index	+0.0081 [-0.0192, +0.0326]	0.5269	1.0000	ns
Direct – NoJEPA	Brier	-0.0029 [-0.0071, +0.0014]	0.1814	1.0000	ns
Direct – NoJEPA	NLL	-0.0081 [-0.0182, +0.0028]	0.1311	1.0000	ns
Direct – NoJEPA	ECE	-0.0011 [-0.0149, +0.0206]	0.8963	1.0000	ns
Attention – NoJEPA	Accuracy	-0.0067 [-0.0257, +0.0105]	0.4824	1.0000	ns
Attention – NoJEPA	Sensitivity	+0.0417 [+0.0146, +0.0758]	0.0081	0.6318	ns
Attention – NoJEPA	Specificity	-0.0294 [-0.0516, -0.0096]	0.0085	0.6576	ns
Attention – NoJEPA	PPV	-0.0061 [-0.0279, +0.0139]	0.5651	1.0000	ns
Attention – NoJEPA	NPV	+0.0120 [-0.0018, +0.0287]	0.1130	1.0000	ns
Attention – NoJEPA	Balanced accuracy	+0.0061 [-0.0113, +0.0257]	0.5144	1.0000	ns
Attention – NoJEPA	MCC	+0.0089 [-0.0244, +0.0447]	0.6175	1.0000	ns
Attention – NoJEPA	Youden index	+0.0123 [-0.0227, +0.0513]	0.5144	1.0000	ns
Attention – NoJEPA	Brier	+0.0001 [-0.0045, +0.0044]	0.9765	1.0000	ns
Attention – NoJEPA	NLL	+0.0002 [-0.0097, +0.0096]	0.9688	1.0000	ns
Attention – NoJEPA	ECE	+0.0170 [-0.0041, +0.0245]	0.0520	1.0000	ns
Attention – Direct	Accuracy	-0.0143 [-0.0333, +0.0057]	0.1535	1.0000	ns
Attention – Direct	Sensitivity	+0.0476 [+0.0123, +0.0880]	0.0132	0.9908	ns
Attention – Direct	Specificity	-0.0434 [-0.0643, -0.0241]	0.0001	0.0097	**
Attention – Direct	PPV	-0.0160 [-0.0401, +0.0075]	0.1740	1.0000	ns
Attention – Direct	NPV	+0.0112 [-0.0055, +0.0303]	0.2187	1.0000	ns
Attention – Direct	Balanced accuracy	+0.0021 [-0.0184, +0.0247]	0.8475	1.0000	ns
Attention – Direct	MCC	-0.0004 [-0.0391, +0.0422]	0.9823	1.0000	ns
Attention – Direct	Youden index	+0.0042 [-0.0369, +0.0493]	0.8475	1.0000	ns
Attention – Direct	Brier	+0.0030 [-0.0033, +0.0086]	0.3119	1.0000	ns
Attention – Direct	NLL	+0.0083 [-0.0062, +0.0213]	0.2403	1.0000	ns
Attention – Direct	ECE	+0.0181 [-0.0152, +0.0301]	0.1537	1.0000	ns

Table A.13: Exploratory pooling comparisons: ForecastAttention minus ForecastMean. All metrics use approximate, null-centered paired patient-bootstrap tests (99,999 shared draws across seeds, models, and landmarks). All p values are two-sided. p_{Holm} adjusts jointly over the 81 secondary tests in Tables A.11–A.14. Direct and Attention abbreviate DirectReadout and ForecastAttention. $^{*}p_{\text{Holm}} < 0.05$, $^{**}p_{\text{Holm}} < 0.01$, $^{***}p_{\text{Holm}} < 0.001$; ns, not significant after the stated correction. Intervals are unadjusted 95% patient-bootstrap intervals (2,000 shared draws) and need not agree with adjusted test decisions. All analyses condition on the fixed, retrospectively selected seed42/44/2026/3407 ensembles; seeds are not resampled.

Landmark	Metric	Difference [95% CI]	p	p_{Holm}	Sig.
T0	AUROC	+0.0007 [-0.0102, +0.0115]	0.9013	1.0000	ns
T0	AP	+0.0102 [-0.0074, +0.0279]	0.2418	1.0000	ns
T0	F1	-0.0009 [-0.0244, +0.0218]	0.9385	1.0000	ns
T1	AUROC	+0.0002 [-0.0091, +0.0099]	0.9757	1.0000	ns
T1	AP	+0.0046 [-0.0104, +0.0205]	0.5589	1.0000	ns
T1	F1	-0.0193 [-0.0394, -0.0006]	0.0457	1.0000	ns
T2	AUROC	+0.0030 [-0.0074, +0.0139]	0.5712	1.0000	ns
T2	AP	+0.0027 [-0.0134, +0.0188]	0.7188	1.0000	ns
T2	F1	-0.0078 [-0.0262, +0.0121]	0.4088	1.0000	ns
Macro	AUROC	+0.0013 [-0.0085, +0.0114]	0.7957	1.0000	ns
Macro	AP	+0.0059 [-0.0088, +0.0209]	0.4206	1.0000	ns
Macro	F1	-0.0093 [-0.0271, +0.0075]	0.2802	1.0000	ns

Table A.14: Exploratory paired T2 minus T0 comparisons. Shared patient-bootstrap draws retain dependence between seeds and landmarks. All metrics use approximate, null-centered paired patient-bootstrap tests (99,999 shared draws across seeds, models, and landmarks). All p values are two-sided. p_{Holm} adjusts jointly over the 81 secondary tests in Tables A.11–A.14. Direct and Attention abbreviate DirectReadout and ForecastAttention. $^{*}p_{\text{Holm}} < 0.05$, $^{**}p_{\text{Holm}} < 0.01$, $^{***}p_{\text{Holm}} < 0.001$; ns, not significant after the stated correction. Intervals are unadjusted 95% patient-bootstrap intervals (2,000 shared draws) and need not agree with adjusted test decisions. All analyses condition on the fixed, retrospectively selected seed42/44/2026/3407 ensembles; seeds are not resampled.

Model	Metric	Difference [95% CI]	p	p_{Holm}	Sig.
NoJEPA	AUROC	-0.0022 [-0.0070, +0.0021]	0.3504	1.0000	ns
NoJEPA	AP	+0.0007 [-0.0079, +0.0088]	0.8711	1.0000	ns
NoJEPA	F1	+0.0074 [-0.0106, +0.0309]	0.4654	1.0000	ns
DirectReadout	AUROC	+0.0012 [-0.0034, +0.0058]	0.6145	1.0000	ns
DirectReadout	AP	+0.0064 [-0.0036, +0.0168]	0.2339	1.0000	ns
DirectReadout	F1	+0.0059 [-0.0211, +0.0409]	0.7005	1.0000	ns
ForecastAttention	AUROC	+0.0021 [-0.0030, +0.0079]	0.4508	1.0000	ns
ForecastAttention	AP	-0.0025 [-0.0128, +0.0079]	0.6360	1.0000	ns
ForecastAttention	F1	+0.0011 [-0.0152, +0.0173]	0.8942	1.0000	ns

Table B.15: Exploratory subgroup discrimination using seeds 42, 44, 2026, and 3407. Within each seed, five fold probabilities are averaged per patient before computing AUROC; Macro averages T0–T2 AUROCs. Model entries are means ± sample SD across the four seed metrics. Paired patient-bootstrap intervals for the mean ForecastAttention minus NoJEPA difference condition on all fitted models and keep the seed set fixed. Age 50 is an exploratory reporting cutpoint. Namespace groups are parts of one clinical release, not external cohorts.

Subgroup (n/pCR+)	NoJEPA	Full	Δ AUROC [95% interval]
HR+/HER2- (68/11)	0.683 ± 0.019	0.709 ± 0.015	+0.026 [-0.001, +0.056]
HR+/HER2+ (31/12)	0.635 ± 0.012	0.646 ± 0.026	+0.011 [-0.046, +0.073]
HR-/HER2+ (14/7)	0.828 ± 0.022	0.871 ± 0.029	+0.043 [-0.019, +0.132]
HR-/HER2- (62/26)	0.639 ± 0.005	0.661 ± 0.019	+0.022 [-0.006, +0.052]
Age <50 (90/29)	0.672 ± 0.012	0.688 ± 0.004	+0.016 [+0.000, +0.032]
Age ≥50 (85/27)	0.784 ± 0.006	0.803 ± 0.009	+0.019 [+0.002, +0.041]
ISPY2 namespace (125/39)	0.683 ± 0.008	0.701 ± 0.003	+0.018 [+0.003, +0.035]
ACRIN-6698 namespace (50/17)	0.844 ± 0.006	0.861 ± 0.011	+0.016 [+0.000, +0.034]




Figure A.7: Six complementary landmark summaries: AUROC, AP, F1, sensitivity, specificity, and Brier score. Metrics are computed separately from each seed's five-fold ensemble probabilities and then averaged across seeds 42, 44, 2026, and 3407. Error bars show descriptive sample seed standard deviations, while shading gives conditional 95% shared-patient-bootstrap intervals for the seed mean. The same 175 patients appear at every seed and landmark.

Table B.16: Archived epoch 20 fixed-training-probe diagnostics for seeds 42, 44, 2026, and 3407. Each entry is the mean ± sample SD of four seed-specific five-fold mean skills, distinct from probability ensembling for pCR. Skill is $1 - \mathrm{L1}_{forecast}/\mathrm{L1}_{copy}$ within each snapshot's EMA coordinates; positive is better than holding the landmark state constant. Forecast targets require a continuous actually observed visit chain. These are training-probe summaries, not best-checkpoint holdout forecast evaluations. NoJEPA has no trained-world diagnostic.

Model	Horizon 1 skill	Horizon 2 skill	Horizon 3 skill
NoJEPA	–	–	–
DirectReadout	+0.009 ± 0.018	+0.052 ± 0.015	+0.159 ± 0.073
ForecastAttention	-0.051 ± 0.020	-0.052 ± 0.036	+0.012 ± 0.018




Figure A.8: Outcome-conditioned changes in predicted pCR probability from T0 to T2. For each seed-specific five-fold ensemble, every patient contributes one probability change. Histograms summarize these within-seed changes and are then equally averaged across seeds; probabilities are not averaged across seeds before constructing a change distribution. Panels separate models and outcome classes. Repeated seed predictions are correlated observations of the same 175 patients and do not quadruple the sample size. Score changes do not establish individual treatment effects.

References

Assran, M., Duval, Q., Misra, I., Bojanowski, P., Vincent, P., Rabbat, M., LeCun, Y., Ballas, N., 2023. Self-Supervised Learning from Images with a Joint-Embedding Predictive Architecture, in: 2023 IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR), pp. 15619–15629. URL: https://doi.org/10.1109/cvpr52729.2023.01499, doi:10.1109/cvpr52729.2023.01499.

Chicco, D., Jurman, G., 2020. The advantages of the Matthews correlation coefficient (MCC) over F1 score and accuracy in binary classification evaluation. BMC Genomics 21, 6. doi:10.1186/s12864-019-6413-7.

Cortazar, P., Zhang, L., Untch, M., Mehta, K., Costantino, J.P., Wolmark, N., Bonnefoi, H., Cameron, D., Gianni, L., Valagussa, P., Swain, S.M., Prowell, T., Loibl, S., Wickerham, D.L., Bogaerts, J., Baselga, J., Perou, C., Blumenthal, G., Blohmer, J., Mamounas, E.P., Bergh, J., Semiglazov, V., Justice, R., Eidtmann, H., Paik, S., Piccart, M., Sridhara, R., Fasching, P.A., Slaets, L., Tang, S., Gerber, B., Geyer, C.E., Pazdur, R., Ditsch, N., Rastogi, P., Eiermann, W., von Minckwitz, G., 2014. Pathological complete response and long-term clinical benefit in breast cancer: the CTNeoBC pooled analysis. Lancet (London, England) 384, 164–172. URL: https://doi.org/10.1016/S0140-6736(13)62422-8, doi:10.1016/S0140-6736(13)62422-8.

Davison, A.C., Hinkley, D.V., 1997. Bootstrap Methods and their Application. Cambridge University Press. URL: https://doi.org/10.1017/CBO9780511802843, doi:10.1017/CBO9780511802843.

Du, S., Xie, W., Gao, S., Zhao, R., Wang, H., Tian, J., Liu, J., Liu, Z., Zhang, L., 2025. Early prediction of neoadjuvant therapy response in breast cancer using MRI-based neural networks: data from the ACRIN 6698 trial and a prospective Chinese cohort. Breast cancer research : BCR 27, 52. URL: https://doi.org/10.1186/s13058-025-02009-6, doi:10.1186/s13058-025-02009-6.

Esser, P., Rombach, R., Ommer, B., 2021. Taming Transformers for High-Resolution Image Synthesis, in: Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition, pp. 12873–12883. URL: https://openaccess.thecvf.com/content/CVPR2021/html/Esser_Taming_Transformers_for_High-Resolution_Image_Synthesis_CVPR_2021_paper.html.

Garrucho, L., Kushibar, K., Reidel, C.A., Joshi, S., Osuala, R., Tsirikoglou, A., Bobowicz, M., Del Riego, J., Catanese, A., Gwoździewicz, K., Cosaka, M.L., Abo-Elhoda, P.M., Tantawy, S.W., Sakrana, S.S., Shawky-Abdelfatah, N.O., Salem, A.M.A., Kozana, A., Divjak, E., Ivanac, G., Nikiforaki, K., Klontzas, M.E., García-Dosdá, R., Gulsun-Akpinar, M., Lafcı, O., Mann, R., Martín-Isla, C., Prior, F., Marias, K., Starmans, M.P.A., Strand, F., Díaz, O., Igual, L., Lekadir, K., 2025. A large-scale multicenter breast cancer DCE-MRI benchmark dataset with expert segmentations. Scientific data 12, 453. URL: https://doi.org/10.1038/s41597-025-04707-4, doi:10.1038/s41597-025-04707-4.

Han, L., Tan, T., Zhang, T., Huang, Y., Wang, X., Gao, Y., Teuwen, J., Mann, R., 2024. Synthesis-based imaging-differentiation representation learning for multi-sequence 3D/4D MRI. Medical image analysis 92, 103044. URL: https://doi.org/10.1016/j.media.2023.103044, doi:10.1016/j.media.2023.103044.

Holm, S., 1979. A simple sequentially rejective multiple test procedure. Scandinavian Journal of Statistics 6, 65–70. URL: https://www.jstor.org/stable/4615733.

Hylton, N.M., Blume, J.D., Bernreuter, W.K., Pisano, E.D., Rosen, M.A., Morris, E.A., Weatherall, P.T., Lehman, C.D., Newstead, G.M., Polin, S., Marques, H.S., Esserman, L.J., Schnall, M.D., 2012. Locally advanced breast cancer: MR imaging for prediction of response to neoadjuvant chemotherapy–results from ACRIN 6657/I-SPY TRIAL. Radiology 263, 663–672. URL: https://doi.org/10.1148/radiol.12110748, doi:10.1148/radiol.12110748.

Hylton, N.M., Gatsonis, C.A., Rosen, M.A., Lehman, C.D., Newitt, D.C., Partridge, S.C., Bernreuter, W.K., Pisano, E.D., Morris, E.A., Weatherall, P.T., Polin, S.M., Newstead, G.M., Marques, H.S., Esserman, L.J., Schnall, M.D., 2016. Neoadjuvant Chemotherapy for Breast Cancer: Functional Tumor Volume by MR Imaging Predicts Recurrence-free Survival-Results from the ACRIN 6657/CALGB 150007 I-SPY 1 TRIAL. Radiology 279, 44–55. URL: https://doi.org/10.1148/radiol.2015150013, doi:10.1148/radiol.2015150013.

Li, W., Newitt, D.C., Gibbs, J., Wilmes, L.J., Jones, E.F., Arasu, V.A., et al., 2022. I-SPY 2 Breast Dynamic Contrast Enhanced MRI Trial (ISPY2), Version 1. URL: https://doi.org/10.7937/TCIA.D8Z0-9T85, doi:10.7937/TCIA.D8Z0-9T85. data set.

Loshchilov, I., Hutter, F., 2019. Decoupled Weight Decay Regularization, in: International Conference on Learning Representations.

URL: https://arxiv.org/abs/1711.05101.

Newitt, D.C., Partridge, S.C., Zhang, Z., Gibbs, J., Chenevert, T., Rosen, M., et al., 2021. ACRIN 6698/I-SPY2 Breast DWI. URL: https://doi.org/10.7937/TCIA.KK02-6D95, doi:10.7937/TCIA.KK02-6D95. data set.

Partridge, S.C., Zhang, Z., Newitt, D.C., Gibbs, J.E., Chenevert, T.L., Rosen, M.A., Bolan, P.J., Marques, H.S., Romanoff, J., Cimino, L., Joe, B.N., Umphrey, H.R., Ojeda-Fournier, H., Dogan, B., Oh, K., Abe, H., Drukteinis, J.S., Esserman, L.J., Hylton, N.M., 2018. Diffusion-weighted MRI Findings Predict Pathologic Response in Neoadjuvant Treatment of Breast Cancer: The ACRIN 6698 Multicenter Trial. Radiology 289, 618–627. URL: https://doi.org/10.1148/radiol.2018180273, doi:10.1148/radiol.2018180273.

Tang, W., Jin, C., Kong, Q., Liu, C., Chen, S., Ding, S., Liu, B., Feng, Z., Li, Y., Dai, Y., Zhang, L., Chen, Y., Han, X., Liu, S., Chen, D., Weng, Z., Liu, W., Wei, X., Jiang, X., Zhou, Q., Mao, N., Guo, Y., 2025. Development and validation of an MRI spatiotemporal interaction model for early noninvasive prediction of neoadjuvant chemotherapy response in breast cancer: a multicentre study. EClinicalMedicine 85, 103298. URL: https://doi.org/10.1016/j.eclinm.2025.103298, doi:10.1016/j.eclinm.2025.103298.

Vaswani, A., Shazeer, N., Parmar, N., Uszkoreit, J., Jones, L., Gomez, A.N., Kaiser, L., Polosukhin, I., 2017. Attention Is All You Need, in: Advances in Neural Information Processing Systems. URL: https://arxiv.org/abs/1706.03762.

Vickers, A.J., Elkin, E.B., 2006. Decision Curve Analysis: A Novel Method for Evaluating Prediction Models. Medical Decision Making 26, 565–574. doi:10.1177/0272989X06295361.