
Trig Final (SLTN v656)

• You should have a calculator (like Desmos) and a unit-circle reference sheet.

Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The arc length is 13 meters. The radius is 3.6 meters. What is the angle measure in radians?




$$\theta = \frac{L}{r}$$
 $r = \frac{L}{\theta}$ $L = r\theta$

 $\theta = 3.611$ radians.


Question 2

Consider angles $\frac{-10\pi}{3}$ and $\frac{13\pi}{4}$. For each angle, use a spiral with an arrow head to **mark** the angle on a circle below in standard position. Then, find **exact** expressions for $\sin\left(\frac{-10\pi}{3}\right)$ and $\cos\left(\frac{13\pi}{4}\right)$ by using a unit circle (provided separately).



Find $sin(-10\pi/3)$

$$\sin(-10\pi/3) = \frac{\sqrt{3}}{2}$$




Find $cos(13\pi/4)$

$$\cos(13\pi/4) = \frac{-\sqrt{2}}{2}$$

Question 3

If $\tan(\theta) = \frac{-45}{28}$, and θ is in quadrant IV, determine an exact value for $\cos(\theta)$.

Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.




Solve the Pythagorean Equation

$$28^{2} + 45^{2} = C^{2}$$

$$C = \sqrt{28^{2} + 45^{2}}$$

$$C = 53$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant IV in a unit circle.



$$\cos(\theta) = \frac{28}{53}$$

Question 4

A mass-spring system oscillates vertically with an amplitude of 7.39 meters, a midline at y = -2.47 meters, and a frequency of 6.07 Hz. At t = 0, the mass is at the midline and moving down. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = -7.39\sin(2\pi6.07t) - 2.47$$

or

$$y = -7.39\sin(12.14\pi t) - 2.47$$

or

$$y = -7.39\sin(38.14t) - 2.47$$