Name: Date:

Trig Final (SLTN v653)

- You can use a calculator (like Desmos)
- You should have a unit-circle with special angles and coordinates marked.

Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The arc length is 10 meters. The radius is 3.2 meters. What is the angle measure in radians?



$$\theta = \frac{L}{r} \qquad r = \frac{L}{\theta} \qquad L = r\theta$$

$\theta = 3.125$ radians.

Question 2

Consider angles $\frac{11\pi}{3}$ and $\frac{-9\pi}{4}$. For each angle, use a spiral with an arrow head to **mark** the angle on a circle below in standard position. Then, find **exact** expressions for $\cos\left(\frac{11\pi}{3}\right)$ and $\sin\left(\frac{-9\pi}{4}\right)$ by using a unit circle (provided separately).





$\cos(11\pi/3) = \frac{1}{2}$

$\sin(-9\pi/4) = \frac{-\sqrt{2}}{2}$

Question 3

If $\tan(\theta) = \frac{15}{8}$, and θ is in quadrant III, determine an exact value for $\cos(\theta)$.

Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.



Solve the Pythagorean Equation

$$8^2 + 15^2 = C^2$$
$$C = \sqrt{8^2 + 15^2}$$
$$C = 17$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant III in a unit circle.



$$\cos(\theta) = \frac{-8}{17}$$

Question 4

A mass-spring system oscillates vertically with a frequency of 3.02 Hz, an amplitude of 6.57 meters, and a midline at $y = 8.28$ meters. At $t = 0$, the mass is at the midline and moving down. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = -6.57\sin(2\pi 3.02t) + 8.28$$

or

$$y = -6.57\sin(6.04\pi t) + 8.28$$

or

$$y = -6.57\sin(18.98t) + 8.28$$