
Trig Final (SLTN v669)

- You can use a calculator (like Desmos)
- You should have a unit-circle with special angles and coordinates marked.

Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The angle measure is 5 radians. The radius is 2.4 meters. How long is the arc in meters?




$$\theta = \frac{L}{r} \qquad r = \frac{L}{\theta} \qquad L = r\theta$$

L = 12 meters.


Question 2

Consider angles $\frac{-19\pi}{6}$ and $\frac{9\pi}{4}$. For each angle, use a spiral with an arrow head to **mark** the angle on a circle below in standard position. Then, find **exact** expressions for $\sin\left(\frac{-19\pi}{6}\right)$ and $\cos\left(\frac{9\pi}{4}\right)$ by using a unit circle (provided separately).



Find $sin(-19\pi/6)$

$$\sin(-19\pi/6) = \frac{1}{2}$$




Find $cos(9\pi/4)$

$$\cos(9\pi/4) = \frac{\sqrt{2}}{2}$$

Question 3

If $\sin(\theta) = \frac{21}{29}$, and θ is in quadrant II, determine an exact value for $\tan(\theta)$.

Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.




Solve the Pythagorean Equation

$$A^{2} + 21^{2} = 29^{2}$$

$$A = \sqrt{29^{2} - 21^{2}}$$

$$A = 20$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant II in a unit circle.



$$\tan(\theta) = \frac{\frac{21}{29}}{\frac{-20}{29}} = \frac{-21}{20}$$

Question 4

A mass-spring system oscillates vertically with a frequency of 3.97 Hz, an amplitude of 6.24 meters, and a midline at y = -2.69 meters. At t = 0, the mass is at the maximum height. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = 6.24\cos(2\pi 3.97t) - 2.69$$

or

$$y = 6.24\cos(7.94\pi t) - 2.69$$

or

$$y = 6.24\cos(24.94t) - 2.69$$