
Trig Final (SLTN v639)

- You can use a calculator (like Desmos)
- You should have a unit-circle with special angles and coordinates marked.

Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The arc length is 11 meters. The radius is 2.8 meters. What is the angle measure in radians?




$$\theta = \frac{L}{r} \qquad r = \frac{L}{\theta} \qquad L = r\theta$$

 $\theta = 3.929$ radians.


Question 2

Consider angles $\frac{7\pi}{3}$ and $\frac{-9\pi}{4}$. For each angle, use a spiral with an arrow head to **mark** the angle on a circle below in standard position. Then, find **exact** expressions for $\cos\left(\frac{7\pi}{3}\right)$ and $\sin\left(\frac{-9\pi}{4}\right)$ by using a unit circle (provided separately).



Find $cos(7\pi/3)$

$$\cos(7\pi/3) = \frac{1}{2}$$



Find $sin(-9\pi/4)$

$$\sin(-9\pi/4) = \frac{-\sqrt{2}}{2}$$

Question 3

If $tan(\theta) = \frac{63}{16}$, and θ is in quadrant III, determine an exact value for $cos(\theta)$.

Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.




Solve the Pythagorean Equation

$$16^{2} + 63^{2} = C^{2}$$

$$C = \sqrt{16^{2} + 63^{2}}$$

$$C = 65$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant III in a unit circle.



$$\cos(\theta) = \frac{-16}{65}$$

Question 4

A mass-spring system oscillates vertically with a midline at y = -3.52 meters, an amplitude of 4.73 meters, and a frequency of 7.62 Hz. At t = 0, the mass is at the midline and moving down. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = -4.73\sin(2\pi 7.62t) - 3.52$$

or

$$y = -4.73\sin(15.24\pi t) - 3.52$$

or

$$y = -4.73\sin(47.88t) - 3.52$$