

Name: _____


Date: _____

Trig Final (SLTN v664)

- You should have a calculator (like [Desmos](#)) and a [unit-circle](#) reference sheet.

Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The radius is 33 meters. The arc length is 97 meters. What is the angle measure in radians?




$$\theta = \frac{L}{r} \quad r = \frac{L}{\theta} \quad L = r\theta$$

$\theta = 2.939$ radians.


Question 2

Consider angles $-\frac{13\pi}{4}$ and $\frac{17\pi}{6}$. For each angle, use a spiral with an arrow head to **mark** the angle on a circle below in standard position. Then, find **exact** expressions for $\cos\left(-\frac{13\pi}{4}\right)$ and $\sin\left(\frac{17\pi}{6}\right)$ by using a unit circle (provided separately).



Find $\cos(-13\pi/4)$

$$\cos(-13\pi/4) = \frac{-\sqrt{2}}{2}$$




Find $\sin(17\pi/6)$

$$\sin(17\pi/6) = \frac{1}{2}$$

Question 3

If $\sin(\theta) = \frac{56}{65}$, and θ is in quadrant II, determine an exact value for $\tan(\theta)$.


Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.



Solve the Pythagorean Equation

$$\begin{aligned}A^2 + 56^2 &= 65^2 \\A &= \sqrt{65^2 - 56^2} \\A &= 33\end{aligned}$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant II in a unit circle.



$$\tan(\theta) = \frac{\frac{56}{65}}{\frac{-33}{65}} = \frac{-56}{33}$$

Question 4

A mass-spring system oscillates vertically with an amplitude of 7.03 meters, a midline at $y = -4.42$ meters, and a frequency of 8.7 Hz. At $t = 0$, the mass is at the midline and moving down. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = -7.03 \sin(2\pi 8.7t) - 4.42$$

or

$$y = -7.03 \sin(17.4\pi t) - 4.42$$

or

$$y = -7.03 \sin(54.66t) - 4.42$$