

Name: _____


Date: _____

Trig Final (Solution v4)

- You can use a calculator (like [Desmos](#))
- You should have a unit-circle with special angles and coordinates marked.

Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The radius is 3.8 meters. The angle measure is 4.9 radians. How long is the arc in meters?




$$\theta = \frac{L}{r} \quad r = \frac{L}{\theta} \quad L = r\theta$$

$L = 18.62$ meters.


Question 2

Consider angles $-\frac{15\pi}{4}$ and $\frac{10\pi}{3}$. For each angle, use a spiral with an arrow head to **mark** the angle on a circle below in standard position. Then, find **exact** expressions for $\sin\left(-\frac{15\pi}{4}\right)$ and $\cos\left(\frac{10\pi}{3}\right)$ by using a unit circle (provided separately).



Find $\sin(-15\pi/4)$

$$\sin(-15\pi/4) = \frac{\sqrt{2}}{2}$$



Find $\cos(10\pi/3)$

$$\cos(10\pi/3) = \frac{-1}{2}$$

Question 3

If $\cos(\theta) = \frac{-11}{61}$, and θ is in quadrant II, determine an exact value for $\tan(\theta)$.


Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.



Solve the Pythagorean Equation

$$\begin{aligned}11^2 + B^2 &= 61^2 \\ B &= \sqrt{61^2 - 11^2} \\ B &= 60\end{aligned}$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant II in a unit circle.



$$\tan(\theta) = \frac{\frac{60}{61}}{\frac{-11}{61}} = \frac{-60}{11}$$

Question 4

A mass-spring system oscillates vertically with an amplitude of 6.41 meters, a frequency of 2.51 Hz, and a midline at $y = -8.57$ meters. At $t = 0$, the mass is at the midline and moving down. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = -6.41 \sin(2\pi 2.51t) - 8.57$$

or

$$y = -6.41 \sin(5.02\pi t) - 8.57$$

or

$$y = -6.41 \sin(15.77t) - 8.57$$