Polynomial Operations PRACTICE (version 34)

1. Let polynomials p(x) and q(x) be defined below.

$$p(x) = -9x^5 + 8x^3 + 6x^2 + 7x + 4$$

$$q(x) = 3x^5 - 10x^4 + x^3 - 9x^2 - 8$$

Express the difference q(x) - p(x) in standard form.

2. Let polynomials a(x) and b(x) be defined below.

$$a(x) = 2x^2 - 8x - 4$$

$$b(x) = 5x + 6$$

Express the product $a(x) \cdot b(x)$ in standard form.

3. Express $(x+1)^4$ in standard (expanded) form.

Polynomial Operations PRACTICE (version 34)

4. Let polynomials f(x) and g(x) be defined below.

$$f(x) = 2x^3 + 16x^2 + 11x - 26$$

$$g(x) = x + 7$$

The quotient of $\frac{f(x)}{g(x)}$ can be expressed as a polynomial, h(x), and a remainder, R (a real number).

$$\frac{f(x)}{g(x)} = h(x) + \frac{R}{x+7}$$

By using synthetic division or long division, express h(x) in standard form, and find the remainder R.

5. Let polynomial f(x) still be defined as $f(x) = 2x^3 + 16x^2 + 11x - 26$. Evaluate f(-7).