
Trig Final (SLTN v675)

• You should have a calculator (like Desmos) and a unit-circle reference sheet.

Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The angle measure is 3.2 radians. The arc length is 91 meters. How long is the radius in meters?




$$\theta = \frac{L}{r}$$
 $r = \frac{L}{\theta}$ $L = r\theta$

r = 28.44 meters.


Question 2

Consider angles $\frac{-8\pi}{3}$ and $\frac{15\pi}{4}$. For each angle, use a spiral with an arrow head to \mathbf{mark} the angle on a circle below in standard position. Then, find \mathbf{exact} expressions for $\sin\left(\frac{-8\pi}{3}\right)$ and $\cos\left(\frac{15\pi}{4}\right)$ by using a unit circle (provided separately).



Find $sin(-8\pi/3)$

$$\sin(-8\pi/3) = \frac{-\sqrt{3}}{2}$$




Find $cos(15\pi/4)$

$$\cos(15\pi/4) = \frac{\sqrt{2}}{2}$$

Question 3

If $\cos(\theta) = \frac{65}{97}$, and θ is in quadrant IV, determine an exact value for $\tan(\theta)$.

Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.




Solve the Pythagorean Equation

$$65^{2} + B^{2} = 97^{2}$$

$$B = \sqrt{97^{2} - 65^{2}}$$

$$B = 72$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant IV in a unit circle.



$$\tan(\theta) = \frac{\frac{-72}{97}}{\frac{65}{97}} = \frac{-72}{65}$$

Question 4

A mass-spring system oscillates vertically with a midline at y = 5.36 meters, an amplitude of 4 meters, and a frequency of 7.16 Hz. At t = 0, the mass is at the minimum height. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = -4\cos(2\pi 7.16t) + 5.36$$

or

$$y = -4\cos(14.32\pi t) + 5.36$$

or

$$y = -4\cos(44.99t) + 5.36$$