
Trig Final (SLTN v649)

- You can use a calculator (like Desmos)
- You should have a unit-circle with special angles and coordinates marked.

Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The angle measure is 3.9 radians. The arc length is 110 meters. How long is the radius in meters?




$$\theta = rac{L}{r} \qquad r = rac{L}{ heta} \qquad L = r heta$$


r = 28.21 meters.

Question 2

Consider angles $\frac{13\pi}{6}$ and $\frac{-11\pi}{4}$. For each angle, use a spiral with an arrow head to **mark** the angle on a circle below in standard position. Then, find **exact** expressions for $\cos\left(\frac{13\pi}{6}\right)$ and $\sin\left(\frac{-11\pi}{4}\right)$ by using a unit circle (provided separately).



$$\cos(13\pi/6) = \frac{\sqrt{3}}{2}$$




Find $sin(-11\pi/4)$

$$\sin(-11\pi/4) = \frac{-\sqrt{2}}{2}$$

Question 3

If $tan(\theta) = \frac{15}{8}$, and θ is in quadrant III, determine an exact value for $sin(\theta)$.

Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.




Solve the Pythagorean Equation

$$8^{2} + 15^{2} = C^{2}$$

$$C = \sqrt{8^{2} + 15^{2}}$$

$$C = 17$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant III in a unit circle.



$$\sin(\theta) = \frac{-15}{17}$$

Question 4

A mass-spring system oscillates vertically with an amplitude of 7.52 meters, a frequency of 2.98 Hz, and a midline at y = 4.87 meters. At t = 0, the mass is at the midline and moving down. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = -7.52\sin(2\pi 2.98t) + 4.87$$

or

$$y = -7.52\sin(5.96\pi t) + 4.87$$

or

$$y = -7.52\sin(18.72t) + 4.87$$