
# Trig Final (SLTN v654)

- You can use a calculator (like Desmos)
- You should have a unit-circle with special angles and coordinates marked.

#### Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The arc length is 7.6 meters. The angle measure is 2.6 radians. How long is the radius in meters?




$$\theta = rac{L}{r} \qquad r = rac{L}{ heta} \qquad L = r heta$$


r = 2.923 meters.

## Question 2

Consider angles  $\frac{8\pi}{3}$  and  $\frac{-15\pi}{4}$ . For each angle, use a spiral with an arrow head to  $\mathbf{mark}$  the angle on a circle below in standard position. Then, find  $\mathbf{exact}$  expressions for  $\sin\left(\frac{8\pi}{3}\right)$  and  $\cos\left(\frac{-15\pi}{4}\right)$  by using a unit circle (provided separately).



$$\sin(8\pi/3) = \frac{\sqrt{3}}{2}$$



Find  $\cos(-15\pi/4)$ 

$$\cos(-15\pi/4) = \frac{\sqrt{2}}{2}$$

## Question 3

If  $\tan(\theta) = \frac{-24}{7}$ , and  $\theta$  is in quadrant IV, determine an exact value for  $\sin(\theta)$ .

Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.




Solve the Pythagorean Equation

$$7^{2} + 24^{2} = C^{2}$$

$$C = \sqrt{7^{2} + 24^{2}}$$

$$C = 25$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant IV in a unit circle.



$$\sin(\theta) = \frac{-24}{25}$$

#### Question 4

A mass-spring system oscillates vertically with a midline at y = 8.56 meters, an amplitude of 3.27 meters, and a frequency of 5.01 Hz. At t = 0, the mass is at the midline and moving up. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = 3.27\sin(2\pi 5.01t) + 8.56$$

or

$$y = 3.27\sin(10.02\pi t) + 8.56$$

or

$$y = 3.27\sin(31.48t) + 8.56$$