

Name: \_\_\_\_\_ Date: \_\_\_\_\_

## Polynomial Operations SOLUTIONS (version 24)

1. Let polynomials  $p(x)$  and  $q(x)$  be defined below.

$$p(x) = 8x^5 + 7x^4 + 2x^3 + 5x - 10$$

$$q(x) = -8x^5 - 7x^4 + 10x^3 - x^2 - 6$$

Express the difference  $p(x) - q(x)$  in standard form.

Get “unsimplified” forms. Then find  $p(x) - q(x)$  with addition/subtraction.

$$p(x) = (8)x^5 + (7)x^4 + (2)x^3 + (0)x^2 + (5)x^1 + (-10)x^0$$

$$q(x) = (-8)x^5 + (-7)x^4 + (10)x^3 + (-1)x^2 + (0)x^1 + (-6)x^0$$

$$p(x) - q(x) = (16)x^5 + (14)x^4 + (-8)x^3 + (1)x^2 + (5)x^1 + (-4)x^0$$

$$p(x) - q(x) = 16x^5 + 14x^4 - 8x^3 + x^2 + 5x - 4$$

2. Let polynomials  $a(x)$  and  $b(x)$  be defined below.

$$a(x) = -2x^2 + 5x - 4$$

$$b(x) = 6x + 5$$

Express the product  $a(x) \cdot b(x)$  in standard form.

You can use a table for multiplication.

*	$-2x^2$	$5x$	$-4$
$6x$	$-12x^3$	$30x^2$	$-24x$
$5$	$-10x^2$	$25x$	$-20$

$$a(x) \cdot b(x) = -12x^3 + 30x^2 - 10x^2 - 24x + 25x - 20$$

Combine like terms.

$$a(x) \cdot b(x) = -12x^3 + 20x^2 + x - 20$$

3. Express  $(x + 1)^6$  in standard (expanded) form.

Remember [the binomial theorem](#). It tells us to use Pascal’s triangle.

$$x^6 + 6x^5 + 15x^4 + 20x^3 + 15x^2 + 6x + 1$$

## Polynomial Operations SOLUTIONS (version 24)

4. Let polynomials  $f(x)$  and  $g(x)$  be defined below.

$$\begin{aligned}f(x) &= x^3 + 12x^2 + 28x + 11 \\g(x) &= x + 9\end{aligned}$$

The quotient of  $\frac{f(x)}{g(x)}$  can be expressed as a polynomial,  $h(x)$ , and a remainder,  $R$  (a real number).

$$\frac{f(x)}{g(x)} = h(x) + \frac{R}{x+9}$$

By using synthetic division or long division, express  $h(x)$  in standard form, and find the remainder  $R$ .

I prefer using synthetic division.

$$\begin{array}{r|rrrr}-9 & 1 & 12 & 28 & 11 \\ & & -9 & -27 & -9 \\ \hline & 1 & 3 & 1 & 2\end{array}$$

So,

$$\frac{f(x)}{g(x)} = x^2 + 3x + 1 + \frac{2}{x+9}$$

In other words,  $h(x) = x^2 + 3x + 1$  and the remainder is  $R = 2$ .

5. Let polynomial  $f(x)$  still be defined as  $f(x) = x^3 + 12x^2 + 28x + 11$ . Evaluate  $f(-9)$ .

You could do this the hard way.

$$\begin{aligned}f(-9) &= (1) \cdot (-9)^3 + (12) \cdot (-9)^2 + (28) \cdot (-9) + (11) \\ &= (1) \cdot (-729) + (12) \cdot (81) + (28) \cdot (-9) + (11) \\ &= (-729) + (972) + (-252) + (11) \\ &= 2\end{aligned}$$

Or, if you reference the [polynomial remainder theorem](#), you can state that you know  $f(-9)$  equals the remainder when  $f(x)$  is divided by  $x + 9$ . Thus,  $f(-9) = 2$ .