
Trig Final (SLTN v635)

- You can use a calculator (like Desmos)
- You should have a unit-circle with special angles and coordinates marked.

Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The angle measure is 3 radians. The radius is 40 meters. How long is the arc in meters?




$$\theta = \frac{L}{r} \qquad r = \frac{L}{\theta} \qquad L = r\theta$$

L = 120 meters.


Question 2

Consider angles $\frac{-15\pi}{4}$ and $\frac{7\pi}{3}$. For each angle, use a spiral with an arrow head to **mark** the angle on a circle below in standard position. Then, find **exact** expressions for $\sin\left(\frac{-15\pi}{4}\right)$ and $\cos\left(\frac{7\pi}{3}\right)$ by using a unit circle (provided separately).



Find $sin(-15\pi/4)$

$$\sin(-15\pi/4) = \frac{\sqrt{2}}{2}$$



Find $cos(7\pi/3)$

$$\cos(7\pi/3) = \frac{1}{2}$$

Question 3

If $\sin(\theta) = \frac{63}{65}$, and θ is in quadrant II, determine an exact value for $\cos(\theta)$.

Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.




Solve the Pythagorean Equation

$$A^{2} + 63^{2} = 65^{2}$$

$$A = \sqrt{65^{2} - 63^{2}}$$

$$A = 16$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant II in a unit circle.



$$\cos(\theta) = \frac{-16}{65}$$

Question 4

A mass-spring system oscillates vertically with an amplitude of 5.29 meters, a midline at y = 4.04 meters, and a frequency of 2.08 Hz. At t = 0, the mass is at the minimum height. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = -5.29\cos(2\pi 2.08t) + 4.04$$

or

$$y = -5.29\cos(4.16\pi t) + 4.04$$

or

$$y = -5.29\cos(13.07t) + 4.04$$