

Name: _____


Date: _____

Trig Final (SLTN v605)

- You should have a calculator (like [Desmos](#)) and a [unit-circle](#) reference sheet.

Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The arc length is 25 meters. The angle measure is 3.3 radians. How long is the radius in meters?




$$\theta = \frac{L}{r} \quad r = \frac{L}{\theta} \quad L = r\theta$$

$r = 7.576$ meters.


Question 2

Consider angles $\frac{15\pi}{4}$ and $-\frac{13\pi}{6}$. For each angle, use a spiral with an arrow head to **mark** the angle on a circle below in standard position. Then, find **exact** expressions for $\cos\left(\frac{15\pi}{4}\right)$ and $\sin\left(-\frac{13\pi}{6}\right)$ by using a unit circle (provided separately).



Find $\cos(15\pi/4)$

$$\cos(15\pi/4) = \frac{\sqrt{2}}{2}$$




Find $\sin(-13\pi/6)$

$$\sin(-13\pi/6) = -\frac{1}{2}$$

Question 3

If $\tan(\theta) = \frac{-12}{5}$, and θ is in quadrant II, determine an exact value for $\cos(\theta)$.

Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.




Solve the Pythagorean Equation

$$5^2 + 12^2 = C^2$$

$$C = \sqrt{5^2 + 12^2}$$

$$C = 13$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant II in a unit circle.



$$\cos(\theta) = \frac{-5}{13}$$

Question 4

A mass-spring system oscillates vertically with a midline at $y = -2.78$ meters, a frequency of 4.06 Hz, and an amplitude of 6.08 meters. At $t = 0$, the mass is at the midline and moving up. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = 6.08 \sin(2\pi 4.06t) - 2.78$$

or

$$y = 6.08 \sin(8.12\pi t) - 2.78$$

or

$$y = 6.08 \sin(25.51t) - 2.78$$