Polynomial Operations PRACTICE (version 22)

1. Let polynomials p(x) and q(x) be defined below.

$$p(x) = 8x^5 + x^4 + 3x^2 - 7x + 4$$

$$q(x) = x^5 - 2x^3 + 10x^2 + 9x - 5$$

Express the difference p(x) - q(x) in standard form.

2. Let polynomials a(x) and b(x) be defined below.

$$a(x) = -6x^2 + 3x - 9$$

$$b(x) = -5x + 4$$

Express the product $a(x) \cdot b(x)$ in standard form.

3. Express $(x+1)^6$ in standard (expanded) form.

Polynomial Operations PRACTICE (version 22)

4. Let polynomials f(x) and g(x) be defined below.

$$f(x) = -4x^3 + 27x^2 - 17x + 3$$

$$g(x) = x - 6$$

The quotient of $\frac{f(x)}{g(x)}$ can be expressed as a polynomial, h(x), and a remainder, R (a real number).

$$\frac{f(x)}{g(x)} = h(x) + \frac{R}{x - 6}$$

By using synthetic division or long division, express h(x) in standard form, and find the remainder R.

5. Let polynomial f(x) still be defined as $f(x) = -4x^3 + 27x^2 - 17x + 3$. Evaluate f(6).