
Trig Final (SLTN v622)

• You should have a calculator (like Desmos) and a unit-circle reference sheet.

Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The arc length is 21 meters. The angle measure is 3.8 radians. How long is the radius in meters?




$$\theta = \frac{L}{r}$$
 $r = \frac{L}{\theta}$ $L = r\theta$


r = 5.526 meters.

Question 2

Consider angles $\frac{-10\pi}{3}$ and $\frac{11\pi}{4}$. For each angle, use a spiral with an arrow head to **mark** the angle on a circle below in standard position. Then, find **exact** expressions for $\cos\left(\frac{-10\pi}{3}\right)$ and $\sin\left(\frac{11\pi}{4}\right)$ by using a unit circle (provided separately).



Find $cos(-10\pi/3)$



Find $sin(11\pi/4)$


$$\cos(-10\pi/3) = \frac{-1}{2}$$

$$\sin(11\pi/4) = \frac{\sqrt{2}}{2}$$

Question 3

If $\cos(\theta) = \frac{-65}{97}$, and θ is in quadrant III, determine an exact value for $\tan(\theta)$.

Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.




Solve the Pythagorean Equation

$$65^{2} + B^{2} = 97^{2}$$

$$B = \sqrt{97^{2} - 65^{2}}$$

$$B = 72$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant III in a unit circle.



$$\tan(\theta) = \frac{\frac{-72}{97}}{\frac{-65}{97}} = \frac{72}{65}$$

Question 4

A mass-spring system oscillates vertically with an amplitude of 5.83 meters, a frequency of 2.59 Hz, and a midline at y = 6.91 meters. At t = 0, the mass is at the midline and moving up. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = 5.83\sin(2\pi 2.59t) + 6.91$$

or

$$y = 5.83\sin(5.18\pi t) + 6.91$$

or

$$y = 5.83\sin(16.27t) + 6.91$$