Polynomial Factoring solution (version 631)

1. The quadratic formula says if $ax^2 + bx + c = 0$ then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$. Use the quadratic formula to solve the following equation.

$$x^2 + 8x + 34 = 0$$

Simplify your answer(s) as much as possible.

Solution

$$x = \frac{-(8) \pm \sqrt{(8)^2 - 4(1)(34)}}{2(1)}$$
$$x = \frac{-(8) \pm \sqrt{64 - 136}}{2(1)}$$

$$x = \frac{-8 \pm \sqrt{-72}}{2}$$

$$x = \frac{-8 \pm \sqrt{-36 \cdot 2}}{2}$$

$$x = \frac{-8 \pm 6\sqrt{2}\,i}{2}$$

$$x = -4 \pm 3\sqrt{2}\,i$$

Notice that i in NOT under the square-root radical symbol!!

2. Express the product of 7 + 8i and -5 - 3i in standard form (a + bi).

Solution

$$(7+8i) \cdot (-5-3i)$$

$$-35-21i-40i-24i^{2}$$

$$-35-21i-40i+24$$

$$-35+24-21i-40i$$

$$-11-61i$$

Polynomial Factoring solution (version 631)

3. Write function $f(x) = x^3 - x^2 - 36x + 36$ in factored form. I'll give you a hint: one factor is (x+6).

Solution

$$f(x) = (x+6)(x^2 - 7x + 6)$$

$$f(x) = (x+6)(x-6)(x-1)$$

4. Polynomial p is defined below in factored form.

$$p(x) = (x+7) \cdot (x+4)^2 \cdot (x-1) \cdot (x-6)^2$$

Sketch a graph of polynomial y = p(x).

