
Trig Final (SLTN v626)

• You should have a calculator (like Desmos) and a unit-circle reference sheet.

Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The angle measure is 4.9 radians. The radius is 2.5 meters. How long is the arc in meters?




$$\theta = \frac{L}{r}$$
 $r = \frac{L}{\theta}$ $L = r\theta$

L = 12.25 meters.


Question 2

Consider angles $\frac{-13\pi}{4}$ and $\frac{10\pi}{3}$. For each angle, use a spiral with an arrow head to **mark** the angle on a circle below in standard position. Then, find **exact** expressions for $\cos\left(\frac{-13\pi}{4}\right)$ and $\sin\left(\frac{10\pi}{3}\right)$ by using a unit circle (provided separately).



Find $cos(-13\pi/4)$

$$\cos(-13\pi/4) = \frac{-\sqrt{2}}{2}$$




Find $sin(10\pi/3)$

$$\sin(10\pi/3) = \frac{-\sqrt{3}}{2}$$

Question 3

If $\tan(\theta) = \frac{-72}{65}$, and θ is in quadrant II, determine an exact value for $\cos(\theta)$.

Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.




Solve the Pythagorean Equation

$$65^{2} + 72^{2} = C^{2}$$

$$C = \sqrt{65^{2} + 72^{2}}$$

$$C = 97$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant II in a unit circle.



$$\cos(\theta) = \frac{-65}{97}$$

Question 4

A mass-spring system oscillates vertically with a midline at y = -5.83 meters, a frequency of 2.73 Hz, and an amplitude of 4.46 meters. At t = 0, the mass is at the midline and moving down. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = -4.46\sin(2\pi 2.73t) - 5.83$$

or

$$y = -4.46\sin(5.46\pi t) - 5.83$$

or

$$y = -4.46\sin(17.15t) - 5.83$$