
Trig Final (SLTN v697)

• You should have a calculator (like Desmos) and a unit-circle reference sheet.

Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The angle measure is 5.3 radians. The arc length is 120 meters. How long is the radius in meters?




$$\theta = \frac{L}{r}$$
 $r = \frac{L}{\theta}$ $L = r\theta$

r = 22.64 meters.


Question 2

Consider angles $\frac{-13\pi}{6}$ and $\frac{13\pi}{4}$. For each angle, use a spiral with an arrow head to \mathbf{mark} the angle on a circle below in standard position. Then, find \mathbf{exact} expressions for $\sin\left(\frac{-13\pi}{6}\right)$ and $\cos\left(\frac{13\pi}{4}\right)$ by using a unit circle (provided separately).



Find $sin(-13\pi/6)$

$$\sin(-13\pi/6) = \frac{-1}{2}$$




Find $cos(13\pi/4)$

$$\cos(13\pi/4) = \frac{-\sqrt{2}}{2}$$

Question 3

If $\sin(\theta) = \frac{-72}{97}$, and θ is in quadrant III, determine an exact value for $\tan(\theta)$.

Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.




Solve the Pythagorean Equation

$$A^{2} + 72^{2} = 97^{2}$$

$$A = \sqrt{97^{2} - 72^{2}}$$

$$A = 65$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant III in a unit circle.



$$\tan(\theta) = \frac{\frac{-72}{97}}{\frac{-65}{97}} = \frac{72}{65}$$

Question 4

A mass-spring system oscillates vertically with a frequency of 7.51 Hz, an amplitude of 4.25 meters, and a midline at y = -2.09 meters. At t = 0, the mass is at the midline and moving up. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = 4.25\sin(2\pi 7.51t) - 2.09$$

or

$$y = 4.25\sin(15.02\pi t) - 2.09$$

or

$$y = 4.25\sin(47.19t) - 2.09$$