
Trig Final (SLTN v634)

- You can use a calculator (like Desmos)
- You should have a unit-circle with special angles and coordinates marked.

Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The arc length is 31 meters. The angle measure is 5.5 radians. How long is the radius in meters?




$$\theta = rac{L}{r} \qquad r = rac{L}{ heta} \qquad L = r heta$$

r = 5.636 meters.


Question 2

Consider angles $\frac{7\pi}{3}$ and $\frac{-9\pi}{4}$. For each angle, use a spiral with an arrow head to **mark** the angle on a circle below in standard position. Then, find **exact** expressions for $\cos\left(\frac{7\pi}{3}\right)$ and $\sin\left(\frac{-9\pi}{4}\right)$ by using a unit circle (provided separately).



Find $cos(7\pi/3)$

$$\cos(7\pi/3) = \frac{1}{2}$$



Find $sin(-9\pi/4)$

$$\sin(-9\pi/4) = \frac{-\sqrt{2}}{2}$$

Question 3

If $\tan(\theta) = \frac{-35}{12}$, and θ is in quadrant II, determine an exact value for $\sin(\theta)$.

Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.




Solve the Pythagorean Equation

$$12^{2} + 35^{2} = C^{2}$$

$$C = \sqrt{12^{2} + 35^{2}}$$

$$C = 37$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant II in a unit circle.



$$\sin(\theta) = \frac{35}{37}$$

Question 4

A mass-spring system oscillates vertically with a midline at y = -6.63 meters, a frequency of 2.13 Hz, and an amplitude of 4.85 meters. At t = 0, the mass is at the midline and moving down. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = -4.85\sin(2\pi 2.13t) - 6.63$$

or

$$y = -4.85\sin(4.26\pi t) - 6.63$$

or

$$y = -4.85\sin(13.38t) - 6.63$$