Polynomial Operations PRACTICE (version 29)

1. Let polynomials p(x) and q(x) be defined below.

$$p(x) = -10x^5 + 5x^3 + 3x^2 + 4x + 9$$

$$q(x) = -4x^5 - 6x^4 - 8x^3 + 2x^2 - 3$$

Express the sum of p(x) + q(x) in standard form.

2. Let polynomials a(x) and b(x) be defined below.

$$a(x) = 6x^2 - 5x - 4$$

$$b(x) = -2x - 8$$

Express the product $a(x) \cdot b(x)$ in standard form.

3. Express $(x+1)^4$ in standard (expanded) form.

Polynomial Operations PRACTICE (version 29)

4. Let polynomials f(x) and g(x) be defined below.

$$f(x) = -4x^3 + 25x^2 + 23x - 13$$

The quotient of $\frac{f(x)}{g(x)}$ can be expressed as a polynomial, h(x), and a remainder, R (a real number).

$$\frac{f(x)}{g(x)} = h(x) + \frac{R}{x - 7}$$

By using synthetic division or long division, express h(x) in standard form, and find the remainder R.

5. Let polynomial f(x) still be defined as $f(x) = -4x^3 + 25x^2 + 23x - 13$. Evaluate f(7).