

Name: _____


Date: _____

Trig Final (SLTN v604)

- You should have a calculator (like [Desmos](#)) and a [unit-circle](#) reference sheet.

Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The arc length is 2.6 meters. The radius is 2.3 meters. What is the angle measure in radians?




$$\theta = \frac{L}{r} \quad r = \frac{L}{\theta} \quad L = r\theta$$


$$\theta = 1.13 \text{ radians.}$$

Question 2

Consider angles $\frac{11\pi}{4}$ and $-\frac{17\pi}{6}$. For each angle, use a spiral with an arrow head to **mark** the angle on a circle below in standard position. Then, find **exact** expressions for $\cos\left(\frac{11\pi}{4}\right)$ and $\sin\left(-\frac{17\pi}{6}\right)$ by using a unit circle (provided separately).

Find $\cos(11\pi/4)$

$$\cos(11\pi/4) = \frac{-\sqrt{2}}{2}$$


Find $\sin(-17\pi/6)$

$$\sin(-17\pi/6) = \frac{-1}{2}$$

Question 3

If $\cos(\theta) = \frac{-16}{65}$, and θ is in quadrant II, determine an exact value for $\tan(\theta)$.


Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.



Solve the Pythagorean Equation

$$\begin{aligned}16^2 + B^2 &= 65^2 \\ B &= \sqrt{65^2 - 16^2} \\ B &= 63\end{aligned}$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant II in a unit circle.



$$\tan(\theta) = \frac{\frac{63}{65}}{\frac{-16}{65}} = \frac{-63}{16}$$

Question 4

A mass-spring system oscillates vertically with a frequency of 3.55 Hz, an amplitude of 5.22 meters, and a midline at $y = 7.58$ meters. At $t = 0$, the mass is at the minimum height. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = -5.22 \cos(2\pi 3.55t) + 7.58$$

or

$$y = -5.22 \cos(7.1\pi t) + 7.58$$

or

$$y = -5.22 \cos(22.31t) + 7.58$$