

Name: _____


Date: _____

Trig Final (SLTN v656)

- You should have a calculator (like [Desmos](#)) and a [unit-circle](#) reference sheet.

Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The arc length is 8.2 meters. The angle measure is 1.8 radians. How long is the radius in meters?




$$\theta = \frac{L}{r} \quad r = \frac{L}{\theta} \quad L = r\theta$$

$r = 4.556$ meters.


Question 2

Consider angles $\frac{11\pi}{4}$ and $-\frac{13\pi}{6}$. For each angle, use a spiral with an arrow head to **mark** the angle on a circle below in standard position. Then, find **exact** expressions for $\sin\left(\frac{11\pi}{4}\right)$ and $\cos\left(-\frac{13\pi}{6}\right)$ by using a unit circle (provided separately).



Find $\sin(11\pi/4)$

$$\sin(11\pi/4) = \frac{\sqrt{2}}{2}$$



Find $\cos(-13\pi/6)$

$$\cos(-13\pi/6) = \frac{\sqrt{3}}{2}$$

Question 3

If $\cos(\theta) = \frac{7}{25}$, and θ is in quadrant IV, determine an exact value for $\tan(\theta)$.

Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.




Solve the Pythagorean Equation

$$7^2 + B^2 = 25^2$$

$$B = \sqrt{25^2 - 7^2}$$

$$B = 24$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant IV in a unit circle.



$$\tan(\theta) = \frac{\frac{-24}{25}}{\frac{7}{25}} = \frac{-24}{7}$$

Question 4

A mass-spring system oscillates vertically with an amplitude of 2.26 meters, a midline at $y = -6.44$ meters, and a frequency of 8.57 Hz. At $t = 0$, the mass is at the midline and moving up. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = 2.26 \sin(2\pi 8.57t) - 6.44$$

or

$$y = 2.26 \sin(17.14\pi t) - 6.44$$

or

$$y = 2.26 \sin(53.85t) - 6.44$$