
Trig Final (SLTN v650)

- You can use a calculator (like Desmos)
- You should have a unit-circle with special angles and coordinates marked.

Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The arc length is 160 meters. The angle measure is 3 radians. How long is the radius in meters?




$$\theta = rac{L}{r} \qquad r = rac{L}{ heta} \qquad L = r heta$$

r = 53.33 meters.


Question 2

Consider angles $\frac{-19\pi}{6}$ and $\frac{13\pi}{4}$. For each angle, use a spiral with an arrow head to \mathbf{mark} the angle on a circle below in standard position. Then, find \mathbf{exact} expressions for $\sin\left(\frac{-19\pi}{6}\right)$ and $\cos\left(\frac{13\pi}{4}\right)$ by using a unit circle (provided separately).



Find
$$sin(-19\pi/6)$$

$$\sin(-19\pi/6) = \frac{1}{2}$$




Find $cos(13\pi/4)$

$$\cos(13\pi/4) = \frac{-\sqrt{2}}{2}$$

Question 3

If $\cos(\theta) = \frac{-65}{97}$, and θ is in quadrant II, determine an exact value for $\tan(\theta)$.

Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.




Solve the Pythagorean Equation

$$65^{2} + B^{2} = 97^{2}$$

$$B = \sqrt{97^{2} - 65^{2}}$$

$$B = 72$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant II in a unit circle.



$$\tan(\theta) = \frac{\frac{72}{97}}{\frac{-65}{97}} = \frac{-72}{65}$$

Question 4

A mass-spring system oscillates vertically with an amplitude of 8.68 meters, a midline at y = -2.3 meters, and a frequency of 3.63 Hz. At t = 0, the mass is at the maximum height. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = 8.68\cos(2\pi 3.63t) - 2.3$$

or

$$y = 8.68\cos(7.26\pi t) - 2.3$$

or

$$y = 8.68\cos(22.81t) - 2.3$$