
# Trig Final (Practice v1)

• You should have a calculator (like Desmos) and a unit-circle reference sheet.

#### Question 1

In the figure below, we see a circle and a central angle that subtends an arc. The angle measure is 2.5 radians. The arc length is 15 meters. How long is the radius in meters?




$$\theta = \frac{L}{r}$$
  $r = \frac{L}{\theta}$   $L = r\theta$ 


r = 6 meters.

## Question 2

Consider angles  $\frac{-19\pi}{6}$  and  $\frac{15\pi}{4}$ . For each angle, use a spiral with an arrow head to **mark** the angle on a circle below in standard position. Then, find **exact** expressions for  $\cos\left(\frac{-19\pi}{6}\right)$  and  $\sin\left(\frac{15\pi}{4}\right)$  by using a unit circle (provided separately).



Find  $cos(-19\pi/6)$ 



Find  $sin(15\pi/4)$ 


$$\cos(-19\pi/6) = \frac{-\sqrt{3}}{2}$$

$$\sin(15\pi/4) = \frac{-\sqrt{2}}{2}$$

## Question 3

If  $\tan(\theta) = \frac{-12}{5}$ , and  $\theta$  is in quadrant II, determine an exact value for  $\sin(\theta)$ .

Ignore any negatives and the quadrant, and draw a right triangle (based on SOHCAHTOA) in standard (quadrant I) orientation.




Solve the Pythagorean Equation

$$5^{2} + 12^{2} = C^{2}$$

$$C = \sqrt{5^{2} + 12^{2}}$$

$$C = 13$$

Rescale the triangle so the hypotenuse is 1. Reflect the triangle into Quadrant II in a unit circle.



$$\sin(\theta) = \frac{12}{13}$$

## Question 4

A mass-spring system oscillates vertically with an amplitude of 7.02 meters, a midline at y = -4.66 meters, and a frequency of 8.94 Hz. At t = 0, the mass is at the midline and moving down. Write an equation to model the height (y in meters) as a function of time (t in seconds).

Any of these equations would get full credit.

$$y = -7.02\sin(2\pi 8.94t) - 4.66$$

or

$$y = -7.02\sin(17.88\pi t) - 4.66$$

or

$$y = -7.02\sin(56.17t) - 4.66$$