



Towards a process-based understanding of rifted continental margins

Marta Pérez-Gussinyé, Jenny Collier, John Armitage, John Hopper, Zhen Sun, C. Ranero

► To cite this version:

Marta Pérez-Gussinyé, Jenny Collier, John Armitage, John Hopper, Zhen Sun, et al.. Towards a process-based understanding of rifted continental margins. *Nature Reviews Earth & Environment*, 2023, 4 (3), pp.166-184. 10.1038/s43017-022-00380-y . hal-04189136

HAL Id: hal-04189136

<https://ifp.hal.science/hal-04189136v1>

Submitted on 31 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Towards a process-based approach to the study of rifted continental margins

Marta Pérez-Gussinyé^{1†}, Jenny S. Collier^{2†}, John Armitage^{3†}, John R. Hopper^{4†}, Zhen Sun^{7†}, and C. R. Ranero^{6,7†}

(1) Marum, Bremen Universität, Bremen, Germany

(2) Department of Earth Science and Engineering, Imperial College London, London, UK

(3) Sciences de la Terre et Technologies de l'Environnement, IFP Energies Nouvelles, Rueil-Malmaison, France

(4) Geological Survey of Denmark and Greenland, Copenhagen, Denmark

(5) CAS Key laboratory of Ocean and Marginal Sea Geology, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou, 511458, China.

(6) Barcelona Center for Subsurface Imaging, ICM, CSIC, Barcelona, Spain.

(7) ICREA, Barcelona, Spain.

†e-mail: mpgussinye@marum.de, jenny.collier@imperial.ac.uk, john-joseph.armitage@ifpen.fr, zhensun@scsio.ac.cn, jrh@geus.dk, cranero@cmima.csic.es

Abstract |

Rifted margins form due to extension and break-up of the continental lithosphere. They host large natural resources and will likely play a key role for the future transition to a carbon-neutral economy. Their potential for becoming reservoirs for hydrothermally-induced mineral deposits, native hydrogen, and as sites for CO₂ storage and generation of geothermal energy, rests on the tectonic, sedimentary and hydrothermal processes that shape them. The interactions between these processes are complex and lead to a wide variety of margins, yet specific well studied margins have led to a tendency to classify them as two end-members, magma-poor or magma-rich, with a reductionist view of their evolution. In this review, we first summarize our current knowledge of the structural elements and key processes shaping margins and their variability. We show how a process-based understanding combined with high-quality observations, can be used to decipher margin evolution, with a focus on specific examples. Finally, we discuss how a better understanding of the feedbacks between deformation, sedimentation, magmatism and fluid-rock interactions, will be crucial to assess the potential of margins to support the new carbon-neutral economy, and suggest new pathways to integrate observations and modelling to de-risk exploration of these environments.

1. Introduction

Rifting of continents is a fundamental process of plate tectonics whereby the continental lithosphere extends and thins, leading to breakup [G] and the formation of new oceans. Rifted margins are found surrounding the once contiguous continents and contain some of the largest sediment accumulations worldwide (Figure 1). Rifting has several wide-ranging consequences. For example, the opening of new oceanic gateways can change ocean circulation, climate and the distribution of life^{1,2}. Breakup also generates CO₂ from decompression melting and thermogenic CH₄ during magma emplacement and cooling in sedimentary sections, which has consequences for understanding climate change during Earth history (e.g., Paleocene-Eocene thermal maximum^{3,4}). During the last stages of extension at some margins, seawater reaches and hydrates the mantle^{5,6}, generating H₂ and CH₄, which fuel chemosynthetic life in the deep seafloor, affecting the global carbon cycle⁷⁻⁹.

For the last 50 years rifted margins have been the target of hydrocarbon exploration, as 60% of world oil and gas reserves reside in these settings¹⁰. Today, as we shift towards a carbon neutral economy, the exploration of rifted margins will be driven by their potential for hosting hydrothermally-induced mineral deposits¹¹⁻¹³, native hydrogen reservoirs^{7,14}, CO₂ storage¹⁵, and as sites for generation of geothermal energy^{16,17}. These applications will require a significantly improved understanding of the tectonic, physio-chemical, hydrological and biological interactions occurring at margins. Bringing these processes, which occur at different temporal and spatial scales, into a common framework will be a key endeavour in future margin research.

The fundamental processes describing basin formation were developed in the 1970's and 80s¹⁸⁻²¹. Margins form due to extension of the continental lithosphere, which thins and subsides as the underlying asthenospheric mantle ascends, decompresses and melts. Extension culminates in the breakup of the former continent and the formation of new oceanic crust. Normal oceanic crust can be classed as 6-7 km thick, with an upper crust formed by extrusive basalts and dolerite dikes, overlying a gabbroic lower crust, and is known as Penrose-like oceanic crust^{22,23}. Traditionally, the volume of magmatism during rifting was taken as a key observable – with margins being classified as either magma-rich to magma-poor^{24,25}. This classification was made by analogy to that of oceanic spreading, where the competition between magmatic processes, which create new crust, and tectonic processes, which extend existing crust, shapes ridge architecture. The influence of tectonics and magmatism is itself controlled by spreading velocity, mantle temperature and composition. Continental margins, however, are further complicated by inheritance^{26,27}, so that the initial lithospheric thermal and compositional structure modulates rifting processes, resulting in a vast variety of margin architectures²⁸.

In this Review we show how a process-based understanding of rifting, based on observations and numerical modelling, illuminates the causes underlying the variety of extensional styles and magmatic budgets.

We first summarize our current knowledge of the structural elements (Section 2) and processes shaping margins (Section 3). Subsequently we demonstrate how a process-based understanding, combined with high-quality observations, can be used to decipher margin evolution across the global spectrum, with a focus on three examples (Section 4). Our intention is to avoid evolutionary templates for margin architecture, but instead give the tools to interpret their variability. Finally, we discuss how an understanding of the interactions between deformation, sedimentation, magmatism, and fluid rock-interactions will be crucial to determine the potential of margins for the new carbon-neutral economy (Section 5).

2. Anatomy of Rifted Margins

Rifted margins lie below sea-level such that drilling and seismic methods (wide-angle (WAS) [G] and multichannel (MCS) [G]) are key in determining their structure (Box 1). The discovery that continental breakup does not always immediately produce Penrose-like oceanic crust, as had been expected for extension under normal mantle temperature²⁴, resulted in new paradigms to understand extension and breakup processes. Exhumed serpentinised mantle with scarce magmatic products was first drilled at the continent-ocean transition (COT [G]) west of Iberia, WIM, which together with its conjugate [G], the Newfoundland margin, NF, became the archetypical magma-poor margins²⁹⁻³² (Figures 1, 2a, 3a).

Another major discovery was sequences of seaward dipping reflections (SDRs [G]) observed in MCS images of several margins^{33,34}. These were shown to be interbedded sequences of sediment and lava flows³⁵⁻³⁷. Margins with SDRs also have high p-wave seismic velocity lower crustal bodies, HVLC [G] (7.3-7.8 km/s) interpreted as magmatic intrusions³⁸, and typically significantly thicker than normal oceanic crust after breakup^{39,40}. These margins are collectively known as magma-rich (Figures 1, 2b, 3b) and are frequently associated with onshore large igneous provinces (LIPs [G]) that form by the intrusion and extrusion of large volumes of magmatism in very short geological periods, ≤ 2 Myr^{41,42} that are usually attributed to mantle plumes [G] (Figure 1a).

Magma-rich margins are easily identified as SDR sequences stand out in MCS images (Figure 2). Margins that do not exhibit SDRs in MCS images were often classified as magma-poor and interpreted to have exhumed mantle at their COT (e.g. the South China sea, the central South Atlantic margins, the East Indian margin⁴³⁻⁴⁶). This interpretation resulted in a binary classification of margins into either magma-poor or rich, which was strongly influenced by the study of the magma-poor margin ophiolites in the Alps^{47,48}. The identification of magma-poor margins from MCS data is however controversial as WAS is needed to clearly identify exhumed mantle at the COT (Box 1, Figure 3), and this ambiguity is becoming more recognised^{49,50}. Furthermore, several margins lacking SDRs appear to contain oceanic crust with magnetic anomaly lineations [G] abutting continental crust, without intervening exhumed mantle, such as the Woodlark Basin^{51,52}, north-eastern Brazilian margins⁵³⁻⁵⁵, Gulf of California⁵⁶, Ligurian⁵⁷, Red Sea⁵⁸, and the northeastern⁵⁹ and

northwestern South China Sea, SCS⁶⁰, where drilling confirmed an abrupt COT^{61,62}. These examples display a wide variety of tectonic architectures and there is not a common structural template that fits them all.

Below we summarize the current knowledge of structural elements across the spectrum of margins. We distinguish a proximal domain under the continental shelf and neighboring necking zone [G]; the distal domain, where the continental crust is less than 15 km thick, and the outer domain, where continental crust is ultra-thin and there is breakup (Figure 1b). Within any margin type, the available observations document variations in all aspects.

2.1 Magma-poor margins. Rifting at these margins is characterised by large brittle faults, practically absent syn-rift magmatism and crustal break-up followed by exhumation of the mantle at the COT²⁵. As there is much uncertainty in the basement composition of the COT, here we consider magma-poor margins only those whose COT has been characterized as exhumed mantle using integrated MCS and WAS data (see Box 1, Figure 1), while recognizing that some margins not classified here as such, may indeed be magma-poor.

From proximal to distal sectors. The continental platform is characterised by small offset faults dipping land- and oceanward and small degree of crustal thinning (> 25 km thick crust) distributed over a wide area. The necking zone, which records the thinning from the platform to the distal domain, is generally narrow (Figure 2a). In the distal domain, large fault bound rotated blocks are connected at depth to sub-horizontal reflectivity in the lower crust (Figure 2a, DR). As crust thins, rotated blocks become smaller and their bounding faults listric [G] (Figure 2a, L).

Towards the outer margin, faults merge at depth into a sub-horizontal reflection often interpreted as a detachment fault^{48,63,64} (Figure 2a, D). WAS data suggest that the detachment is directly underlain by serpentinitised mantle^{6,65} (Figure 3a), although a thin lower-crust layer underlying the detachment and overlying serpentinitised mantle might be possible⁶⁶. The lateral extent of these detachments varies from 30 -15 km (e.g., S and H reflectors, in the West Iberia, Armorican margins and Goban Spur^{48,64,67-69}). In some places, however, the continental crust thins to breakup without the involvement of such structures (Tagus Abyssal Plain in West Iberia^{70,71}).

Asymmetry. Magma-poor conjugate margins are often asymmetric (Figure 2a). This is the case of the Labrador – West Greenland margins⁷², and of the northern sector of the WIM-NF margins, where the WIM is wider and has more rotated blocks, as described above, and the NF displays fewer fault blocks, no obvious detachment, and abrupter crustal thinning^{73,74} (Figure 2a). To the south, central sectors of the WIM-NF margins (Iberia Abyssal plain and central Newfoundland^{75,76}) shows the opposite trend. However, modern coincident MCS and WAS data on conjugate margin pairs are rare, thus it is not clear whether all magma-poor margins are asymmetric. For example, the Porcupine failed rift has flanks that appear symmetric although one fault

vergence dominates at ultra-thin crust⁷⁷, indicating that asymmetry may emerge during terminal extension^{74,78,79}.

The Continent-Ocean Transition. The COT of magma-poor margins contains a series of margin-parallel ridges^{76,80}. Where drilled, in the WIM-NF, these ridges consist of serpentinised peridotite with a small proportion of magmatic products³² (Figure 2a). The P-wave velocity of exhumed mantle reduces with degree of serpentinisation, from ~ 8 to 4.5 km/s ⁸¹ (Figure 3a, Box 1) and is anisotropic⁸², complicating interpretation of WAS profiles. This seismic velocity range overlaps with that of different types of metamorphic, intrusive magmatic or oceanic rocks. However, a shallow steep velocity gradient and lack of abrupt crust-to-mantle velocity boundary (Moho reflection) characterise exhumed mantle domains⁸³ (Box 1).

The width of exhumed mantle and its seismic velocity structure is variable (Nova Scotia^{84,85}, Labrador sea^{72,86}, Orphan basin⁸⁷, Tyrrhenian^{88,89}, Deep Galicia margin⁶⁵, Tagus Abyssal Plain⁷¹, Flemish cap⁹⁰, Goban Spur⁹¹, Bay of Biscay⁶⁷), and may be asymmetric across conjugate margins (Figure 2a). For example, it is $\sim 160 \text{ km}$ wide in the central WIM⁸³, whereas in the central NF the COT is $\leq 30 \text{ km}$ wide⁷⁵, or may even be thin lower continental crust⁹².

The degree of magmatism at the COT may also be asymmetric (Figure 2a). Comparison of WAS data in the WIM and NF suggest the NF may be more magmatic^{75,92} (Figure 3a). Unfortunately, it is unclear whether some of the WAS velocity differences among the COTs reflect real variability in exhumation and serpentinisation processes, or are related to differences in velocity/depth models related to OBS spacing and modelling techniques (Box 1). Imaging small-scale structures, such as few-km-thick oceanic crust or discrete gabbroic bodies within serpentinised mantle, a narrower-than-typical OBS spacing of $10\text{--}15 \text{ km}$ is required and modern WAS data inversion techniques⁸⁵.

Oceanic Crust. A Penrose-like, $6\text{--}7 \text{ km}$ thick oceanic crust with a 2-gradient velocity structure and Moho reflection (Box 1) is not always observed oceanwards of exhumed mantle (Figure 2a), either because the seismic lines do not extend far enough, because of complexities in the 3D structure, or because the structure of oceanic crust in ultra-slow spreading crust differs from normal oceanic crust (Box 1). In West Iberia, WAS data show slightly thicker than normal oceanic crust abutting exhumed mantle at the location of the prominent J-magnetic-anomaly indicating an abrupt inception of seafloor spreading⁸³ (Figure 3a). However, oceanwards of this location, oceanic crust is thinner than normal ($\sim 4 \text{ km}$ thick^{71,83}). In the northern sector of the conjugate Newfoundland margin, WAS data indicate that continental crust transitions directly to anomalously thin, $1\text{--}3 \text{ km}$, igneous oceanic crust, with layer 3 absent in its thinnest portions, overlying serpentinised mantle⁷³ (Figure 2a); while in the central sector of the Newfoundland margin, thin oceanic crust may appear after $\leq 30 \text{ km}$ of mantle exhumation⁷⁵. These observations open different possible scenarios for the establishment of seafloor spreading (see Section 4).

2.2 Magma-rich margins.

These margins are characterized by voluminous magmatism during rifting. There is a complex interplay between tectonically induced crustal thinning and magmatism in the form of deep intrusions, diking, and extrusions⁹³⁻⁹⁵. In studies of magma-rich margins in the North Atlantic, distinct volcano-seismic facies and reflection characteristics were recognised that reflect different lava emplacement environments^{96,97} (Figure 4). Key to this interpretation is to recognise if extrusions erupted into subaerial, shallow water lacustrine/marine or deep marine environments. Seismic interpretation below SDRs is challenging because the high acoustic impedance contrast between sediment and basalt. Magmatism that re-thickens the thinned crust may also mask tectonically induced structures which complicates interpretation⁹⁸.

From Proximal to Distal sectors. The magmatic facies depend strongly on the inherited lithospheric structure²⁷. In the North Atlantic, pronounced crustal thinning beginning in the late Paleozoic and continuing throughout the Mesozoic occurred without significant magmatism, forming shallow marine basins. Initial volcanism erupted into shallow water resulting in hyaloclastite flows, tuff deposits, lava deltas, and landward flows that mark the proximal margin such as off Vøring and East Greenland^{96,99} (Figure 4). In contrast, the first volcanism in the South Atlantic appears to have erupted into thicker continental crust undergoing necking associated with early extension. This resulted in subaerial lava flows filling half grabens along landward dipping faults¹⁰⁰⁻¹⁰² (Figures 2b and 4). In the literature these are termed type I SDR to distinguish them from the main type II SDR sequence over the COT, which reach thickness of up to 15 km, significantly thicker than those imaged in the North Atlantic^{102,103}. The lower crust of the proximal margin and COT is typically marked by HVLC interpreted as evidence for intrusion of high MgO magmas at depth¹⁰⁴⁻¹⁰⁷. In the North Atlantic the presence of injected sills within the lower crust has been directly imaged within the HVLC⁹⁸. In the South Atlantic, the mid- and lower-crust shows high-amplitude discordant and anastomosing reflectivity, suggesting either ductile deformation during extension and/or injected sills^{108,109} (Figure 2b, DR).

The Continent-Ocean Transition. The COT is marked by the main SDR sequences. In the South Atlantic, the SDRs are spatially uniform and appear to form continuously (type II SDR, equivalent to 'inner SDR' on the Vøring margin¹⁰²; Figure 2b, 4). The reflectivity pattern of type II SDR closely resembles the Cenozoic lava sequences in eastern Iceland, which are built up into laterally extensive flow units with arcuate shapes and consistent dips towards the neo-volcanic zone^{34,110}. These lavas are believed to flow some kilometres away from the centres of eruption. Their seaward dip is attributed to the subsidence of the plate as it moves away from the axis^{111,112}. The reflectivity pattern below most type II SDRs exhibits much smaller amplitudes than below type I SDRs, consisting of a shallower seismically transparent package, underlain by discordant high amplitude reflections in the lower crust¹⁰⁰. This suggests that intrusion below the type II SDRs takes place in at most a very attenuated continental crust, so that most of the material is magmatic and exhibits less impedance contrast with the surrounding crust⁹⁸ (Figure 2b). On many margins, some extended ultra-thin continental crust cannot be excluded from the COT¹¹³. However, the pattern of reflectivity in combination with

the lack of faulted contacts between the type II SDRs and basement suggests a COT formed by subaerial oceanic spreading that produces thicker igneous crust than normal. This interpretation is supported by the recognition of linear magnetic anomalies very similar to submarine seafloor spreading anomalies¹¹⁴.

Oceanic crust. As the melt supply is reduced, volcanism evolves towards normal seafloor spreading. Along Greenland and Vøring, this reduction was gradual and resulted in progressive subsidence of the proto-spreading centre⁴⁰. Here a transition from subaerial to shallow marine volcanism is marked by a rough top basement reflection or an outer volcanic high interpreted as hyaloclastite flows^{96,115} (Figure 4). Eventually, the spreading system subsides to deeper water where the pressure prevents degassing and hyaloclastite formation (Figure 4). Large sheet flows form an outer set of SDR (Figure 3b). Well-developed linear magnetic anomalies can be interpreted in the crust that forms in these submarine environments. In contrast, in the South Atlantic, the peak magmatism coincided with the subaerial SDR stage, with the first submarine oceanic crust having a thickness of just 1 or 2 km more than normal¹¹⁶ (Figure 4).

Asymmetry. The distribution of magmatic products may vary both between the conjugate pairs and along strike of the province. For example, in the South Atlantic WAS data shows 4 times larger HVLC zones in the African compared to the South American side¹¹⁷. This conjugate asymmetry is also observed in the SDRs which are longer and flatter in the African side (Figure 2a). Overall, southwards of the Parana-Etendeka LIP the magma budget becomes smaller (and the distribution more asymmetric^{118,119}) whereas to the north there is little magmatism^{120,121}. Similarly, in the North Atlantic south of Iceland the volume of magmatic rocks is much larger along the Greenland than European sides (Figure 3b), whereas to the north of it is opposite¹²². All these variabilities point to the interaction of the plume material with pre-existing lithospheric structure during the generation of magma-rich margins¹²³⁻¹²⁵.

2.3 An example of an intermediate margin. Intermediate margins display no SDRs, indicative of excessive magmatism nor exhumed mantle at the COT, indicative of deficient magmatism. However, they may display geophysical evidence for robust magmatism as lower crustal magmatic bodies, as in the Angola margin¹²⁶, and detachment structures, as in the SCS^{127,128}, which resemble structures observed in both magma-rich and poor margins. Here we use the SCS as an example of, but not a template for, an intermediate margin. The South China Sea rifted a lithosphere formed by a Mesozoic subduction system. Bounded by the Yangjiang-Yitongansha Transfer Fault, the eastern margin broke up in the fore-arc area and the western margin broke up roughly along the volcanic arc^{129,130}. Therefore, the crustal structure, deformation style and the amount of the syn-rift magmatism varies greatly¹²⁸. The most distinguishable feature is that the eastern margin has up to 12 km thick HVLC, while the western margin usually bears sporadic thin layer (~2 km thick) of HVLC^{131,132}. Here, we focus on the north-eastern sub-basin where two IODP legs took place.

From Proximal to Distal sectors. Thinned continental crust in this segment extends asymmetrically across a large width, ~450 km in the north (Figure 2c). Thinning is progressive, but a clear necking zone is observed^{60,133}. The proximal margin extends over 200 km and has a laterally homogeneous crustal thickness of ~25-30 km. In this area, upper crust is extended by normal high-angle faults forming wide horsts and narrow and deep grabens (HG, Figure 2c). Seawards, the crust thins abruptly to ~7 km and subsequently the large-scale trend of crustal thinning is punctuated by local thickening below what are interpreted as large detachment faults^{127,134} (D, Figure 2c). In WAS data¹³⁵, this thickening is accommodated mainly by the addition to the lower crust of high-velocity layers, HVLC, with P-wave velocities of 7.0-7.3 km/s. Above the HVLC, high-amplitude discordant and anastomosing reflectivity in the lower crust is indicative of magmatic intrusion at higher crustal levels accompanied by intense ductile lower crustal flow^{127,134}. Unlike magma-rich margins, SDRs are not observed and instead sills are interpreted to occur at syn-rift sedimentary levels¹³⁶.

The Continent-Ocean Transition. The COT at intermediate margins is narrow, typically ≤ 30 km⁵⁹⁻⁶¹ (Figure 2c). In the SCS, increasing magmatic intrusions are associated with basement uplift and sills are emplaced in the overlying syn-rift sequences¹³⁴. Syn-intrusive forced folds in the sediment sequences suggest dike intrusion occurred in late stage rifting and became younger oceanward¹³⁶. Seawards, the HVLC diminishes in thickness and crustal breakup juxtaposes ultra-thin continental crust with normal Penrose-like oceanic crust¹³⁵.

Oceanic crust. The earliest mid-ocean ridge basalt that IODP drilled is from a volcanic edifice located in the outer margin, underlain by intra-crustal reflections which are interpreted as dykes cutting through ultra-extended crust¹³⁷. Seismic images from several segments support that magma instead of faults trigger the initial breakup in the SCS^{59,60}. At the IODP location, seismic images suggest that the COT may be composed of highly thinned continental crust overlain by erupted basalt and underplated by gabbroic lower crust that grade into and the early oceanic crust^{136,138}. The early oceanic crust is 5-6 km thick, and thickens slightly to 7-8 km oceanward over 100 kms^{139,140}.

3. Key parameters and processes

Rifted margins form by the interaction between tectonic, magmatic, hydrothermal, and surface processes modulated by the initial lithospheric structure and extension rates. These interactions are complex and lead to a variety of margin architectures and evolutions. In this section, lithospheric-scale numerical models are used to depict some of these interactions. These models summarize previously published results that show that the initial and evolving lithospheric strength determine the spatio-temporal distribution of deformation^{28,141}, fault geometry¹⁴², stratigraphy^{143,144} and magmatism¹⁴⁵⁻¹⁴⁷ (Figure 5, Suppl. Info and Movie S1-S4). Interactions not covered by these models are referenced from the published literature.

3.1 Lithospheric rheology. Observations of rifted margins such as the North Sea¹⁴⁸, Gulf of Corinth¹⁴⁹ and Suez¹⁵⁰ have shown that early extension is spatially distributed across numerous disconnected, short faults, which with time grow laterally and merge so that deformation becomes progressively localized in fewer, larger faults. The predominance in time and space of each of these phases, distributed vs. localized, determines the width and asymmetry of rifted margins, and is related to the importance of faulting vs. ductile thinning, which is a function of the evolving lithospheric rheology^{28,141,151}. In turn, this depends on lithospheric composition, geotherm, extension velocity and potential rift-plume interactions.

For strong lithospheres, initial deformation is coupled from the upper crust to the mantle, through narrow shear zones in the lower crust, leading to relatively rapid crustal thinning and mantle uplift (Suppl. Figure 2a, Movie S1). This makes the first phases of deformation already quite spatially localized and narrow^{28,141}, as in the Baikal and East African rifts, which develop close to strong cratons or ancient platforms^{28,152}.

When the lower crust is weak, initial deformation is characterized by viscous thinning of the lower crust over a broad area, which decouples deformation between upper crust and mantle and leads to comparatively slow crustal thinning and mantle uplift (Suppl. Figure 2b). This results in the formation of a wide system of horst and grabens, HG, underlain by little extended crust (Movie S3), as observed in the Basin and Range and the proximal margin of the South China sea, where the pre-rift lithosphere is thinner and weaker than in the previous examples^{28,127}.

As extension continues and the crust thins to $< \sim 15$ km conductive cooling becomes increasingly important^{153,154}. This cooling, which is more predominant for slower extension velocities, in addition to strain weakening [G], which reduces the strength of faults and shear zones where extension was previously localised, leads to progressive localisation of deformation, and its coupling from the upper crust to the mantle, through faults and narrow shear zones. For the same extension velocity, rifts that started in strong lithosphere will break-up quickly, forming two narrow symmetric conjugates^{28,141} (Figure 5a, Movie S1). Instead, those starting in weak lithospheres, will only localize after a prolonged phase of distributed extension, followed by rapid localization and break-up. Their architecture is dominated by horsts and grabens, HG, formed during the initial distributed phase (Figure 5c, Movie S3), resulting in wide, symmetric margins^{28,141}.

Symmetric margin development is disrupted when lateral migration of the deformation from one rift side towards the conjugate becomes dominant¹⁵⁵. This occurs when the lower crust becomes strong enough to couple deformation from upper crust to mantle, but is still ductile enough to prevent crustal breakup by faulting^{145,155,156} (Suppl. Figure 2c, Movie S2). In this case, conjugate margins evolve asymmetrically, via a system of oceanward younging, deep-penetrating fault/shear zones that work sequentially in time and nucleate in the hangingwall [G] of the previous deep penetrating structure. This process generates a wide margin where the sequential fault array [G], SF is active, and a narrow one which constitutes the hangingwall to the SF array⁷⁴

(Figure 5b, Movie S2). Lateral rift migration becomes more dominant with increasing extension velocity, as it allows focused localisation at high temperatures which keep the lower crust and mantle ductile, thus preventing break-up¹⁵⁵.

Conjugate margin asymmetry can also develop after a long phase of slow, distributed deformation, which results in cooling and strengthening of the extending areas and localisation of rifting into the side of the basin, where the crust is thicker and lithosphere weaker^{157,158} (Figure 5d, Movie S4).

3.2 Fault distribution and geometry. The evolution of fault systems and their relationship to ductile deformation is important for assessing the amount of extension accommodated by brittle vs. ductile processes, which impacts heat-flow. Fault system evolution strongly depends on lower crustal rheology^{143,159} (Movies S1-S4).

During early rifting in weak lower crust, deformation is accommodated by many, simultaneously active, small-offset faults, forming a system of horst and grabens, HG (Figure 5c-d, Movies S3-S4). With increasing extension, new generations of faults cut the previous ones, a process known as polyphase faulting, PF (Figure 5c and 5d), which may prevent fault recognition in MCS sections^{160,161}. For strong lower crust, early faults exhibit larger offsets and faulting quickly migrate basinward, where no previous large faults were active, and the PF is less intense (Movie S1, Figure 5a).

With increasing extension, deformation may migrate laterally, and fault geometry will depend on the relative strength of the lower crust. When it is relatively strong, the wide margin exhibits a series of oceanward rotated fault blocks, bounded by sequentially active faults, SF, that become progressively listric (SF, Figure 5b, Movie S2) and eventually merge at depth to form sub-horizontal structures resembling detachment faults^{74,142}. For relatively weaker crust, faults can attain longer offsets, as the flexural work needed to elastically accommodate fault offset is small, and acquire concave-downward geometries (CCD in Figure 5d, Movie S4), similar to those observed in oceanic core complexes¹⁶²⁻¹⁶⁵ and interpreted in the distal margins of the SCS (Figure 2c).

3.3 Sedimentation. In addition to rheology, numerical models have shown that syn-tectonic sedimentation influences deformation by increasing fault offset and thermally insulating the lithosphere, thus keeping it warmer for longer. The first effect is most important in the proximal margin sectors, while the second will decrease cooling in the distal areas where the crust is thin, increasing the ductility of the lower crust and mantle in heavily sedimented margins^{144,166-168}.

In terms of stratigraphic architecture, early models of extension assumed that faulting occurred synchronously with sediment deposition. Thus, syn-rift [G] sediments were classified as those exhibiting wedge-shape geometries against their bounding faults, whereas post-rift [G] sediments draped over previous topography, indicative for post-tectonic [G] deposition. Today, observational studies^{74,148,169,170} and numerical

models^{143,144} show that during rifting deformation migrates oceanward. This results in the syn-tectonic sediment [G] younging oceanward, and the upper syn-rift sedimentary section drapes over previous topography along the more proximal margin sections, i.e. has a post-tectonic geometry^{74,143,171} (Figure 5).

When the margin is wide, post-tectonic sequences can form large sag basins of syn-rift age, as observed in the central South Atlantic (e.g. Angola margin^{143,172}, Santos basin¹⁷³, Figure 5d). As a result of rift migration, unconformities [G] separating syn- and post-tectonic packages across a margin do not date continental breakup, as earlier assumed, but the age at which deformation terminated in a given area and migrated basinwards. Only unconformities across the outer margin should match breakup age¹⁴³. This is an important finding as such unconformities had been typically used to date breakup age.

3.4 Melting. During thinning of the continental crust, lava is extruded at magma-rich margins forming flows imaged as SDRs in MCS images⁹⁶. These extrusions are accompanied by intrusions of melt into the lower crust⁹⁸ (Figure 2b). Intrusions will change how extension is accommodated. For example, in Afar extension was initially accommodated by slip at the border faults, but presently extension is significantly accommodated by melt intrusion at the rift centre¹⁷⁴, which facilitates rifting¹⁷⁵. Repeated dike and sill intrusions that can generate the typical thickness of ~10 km of HVLC in periods of roughly 5 Myr, as observed magma-rich margins e.g. Hatton Bank^{27,98}, would increase crustal temperatures by around 800°C¹⁷⁶, potentially weakening the crust and promoting significant ductile deformation¹¹⁹.

On the other hand, not only will melting influence crustal strength, but the rheology of the crust will also impact melt generation during extension. Thus, unlike at mid-ocean ridges melting is not only dependent on extension velocity, mantle temperature, and composition, but also on lower crustal rheology^{145,177}. Spatially distributed extension tends to delay the onset of decompression melting as mantle uplift is slow, while localised extension tend to focus melting at the rift centre, thus resulting in a different spatial distribution of melt intrusion along the margin (Figure 5, Movies S1-S4).

In addition, the composition of lavas and the surface area of flows in Iceland suggests that magmatism is sensitive to surface processes, in this case the change in load due to deglaciation¹⁷⁸. Thus, there is a connectivity between surface processes and deep mantle processes and the response times are rapid¹⁷⁹. When the lower crust is weak, sediment loading can likewise impact the degree of decompression melting¹⁸⁰.

3.5 Hydrothermal circulation. The influence of hydrothermal circulation on the temperature field during extension, and on shaping fault geometry and margin architecture, is not yet understood. Models coupling deformation to hydrothermal circulation to our knowledge do not exist, mainly because of the complex non-linear problem of coupling both fluid flow for highly viscous fluids (water and brine) with the visco-elasto-plastic deformation of the crust and lithosphere. However, models have been used to look at the impact of hydrothermal circulation on crustal accretion at mid-ocean ridges¹⁸¹ and it has a profound effect on crustal temperatures and the depth of the melt lens. During margin formation hydrothermal circulation will likewise

favour mantle serpentinization as it will decrease the geothermal gradient, particularly at distal margins. In addition, understanding the evolution of hydrothermal system will be important in predicting the location of lithium rich brines and sediment-hosted metals¹¹.

4. Towards a process-based understanding of rifting

Having described how various geological attributes and processes influence margin evolution, we now focus on how they shape the structural elements observed across the wide spectrum of margins, which is crucial to assess their potential to host resources for the new green economy.

4.1 Rifting at magma-poor margins. Magma-poor margins form where extension velocities are on the ultra-slow range¹⁸² (< 20 mm/yr full spreading) or the mantle is originally cold¹⁸³ or depleted¹⁸⁴. These factors result in cold geothermal conditions which have several consequences.

Absence of magmatism and mode of extension. Under the above cited circumstances numerical models predict very little or no melt production and mantle exhumation at the COT¹⁸⁵. Initial rifting conditions suggest that extension in these margins probably started with a decoupled upper and lower crust^{74,186}. Due to the ultra-slow extension, conductive cooling became important with ongoing extension, resulting in an increased localisation and coupling of deformation throughout the crust⁵. This led to the formation of large faults, which continued as ductile shear zones at depth, and pulled the ductile lower crust towards their hangingwall as observed in WAS data⁷⁹. In numerical models⁶⁶ this pattern of extension leads to deformation fabrics that resemble the lower crustal reflectivity patterns, DR, often observed in magma-poor margins^{43,79} (Figure 2a) indicating the global predominance of coupling processes at these margins.

Faulting, serpentinisation and mantle exhumation. The history of faulting, serpentinisation and mantle exhumation influences the potential of magma-poor margins to host natural H₂ reservoirs, as H₂ is a by-product of serpentinisation. Early, simple 1D models of the rheological evolution of these margins predicted that cooling leads to embrittlement of the whole crust before detachments observed in the distal margin were formed⁵ (D in Figure 2a). This was thought to be a key stage during margin evolution, (also known as "coupling point"⁴³), as it allows faults to reach the mantle and bring enough water to serpentinise it⁵. Serpentinisation induced weakening would have facilitated the formation of low-angle detachments in the brittle field⁶³. However, new 2D dynamic simulation techniques allowing the reconstruction of seismic sections at fault-block scale (Figure 6), suggest that these detachments may form in the ductile field⁶⁶. During extension of the ultra-thin crust (< 10 km thick), the hangingwall of deep penetrating faults becomes hotter and ductile, so that the overlying faults propagate at depth into ductile shear zones active at low angles⁶⁶ as interpreted from 3D MCS data⁶³. Instead, the footwall [G] cools and becomes brittle allowing mantle serpentinisation in this area⁶⁶.

Crustal break-up and mantle exhumation. Final breakup occurs through either a pair of conjugate normal faults, forming two distal highs on each side of the new ocean floor, formed by either serpentinised

mantle, as drilled in the West Iberia margin²⁹ or continental crust, as interpreted in Newfoundland⁷³. Alternatively, interpretation from margin ophiolites in the Alps, suggests it occurs through concave-downward detachments exposing serpentinised mantle and generating and asymmetric COT¹⁶². Numerical models indicate that both situations are possible with normal faults forming in slightly colder conditions than concave-downward detachments¹⁶³ (Figure 5c,d, Movies S2, and S4). Mantle exhumation may occur through a series of faults that cut each other's footwall, known as flip-flop detachments, inferred to form the system of ridges observed in the COT of magma-poor margins¹⁸⁷.

Transition to seafloor spreading. The abrupt lateral change from exhumed mantle to oceanic crust west of Iberia indicates rapid initiation of seafloor spreading, perhaps supporting that the first oceanic crust was formed by mid-ocean ridge propagation⁷¹. In contrast a slow transition from exhumed mantle to thin oceanic crust as suggested for offshore central Newfoundland, might support the gradual establishment of a spreading centre by progressive thinning of the lithosphere^{73,75,92} and a gradual change from tectonic to magmatic processes¹⁸⁸. Such a progressive increase in magmatism would be promoted if rifting velocities increase during extension¹⁸⁹. In addition, the asymmetric distribution of exhumed mantle suggests that the location of deformation may have jumped, during the establishment of the steady-state oceanic spreading centre^{71,73}. The ridge propagation hypothesis would indicate that magma-poor margins form where the mantle is cold and only transitions to spreading upon ridge propagation and thus indicates strong lateral heterogeneity of the sub-lithospheric mantle. Instead, the progressive establishment of a mid-ocean ridge would suggest that mantle exposure is produced under ultra-slow velocities which increase with time as a result of extension dynamics¹⁸⁹.

4.2 Rifting at magma-rich margins. Excess magmatism at rifted margins is principally a function of the temperature of the mantle, but can be significantly modified by many secondary inherited attributes, such as lithosphere structure, asthenosphere composition, and processes such as small-scale convection¹⁹⁰.

Plume-lithosphere interaction. Observations show that magmatism is not distributed symmetrically around the LIP or assumed plume impact location, as classic models for magma-rich margins would suggest^{24,191}. These variabilities point to the importance of the interaction of the plume material with pre-existing lithospheric structure in the generation of magma-rich margins¹²²⁻¹²⁴. Since buoyant plume material will flow towards the thinnest lithosphere¹⁹², a key factor for the formation of magma-rich margins is the timing of the arrival of the thermal plume relative to the thinning of the continental lithosphere. In particular, if there is no prior extension, the mantle will likely cool significantly prior to eventual breakup leading to limited decompression melting of a depleted mantle as seen in the Seychelles-Laxmi Ridge margin²⁷. In contrast, significant pre-existing thin spots that drain plume material favours forming magma-rich margins^{123,124}.

Brittle vs. ductile deformation and emplacement of SDRs. How the continental crust extends during rifting at magma-rich margins is not clear, as seismic images are blurred beneath the thick SDR piles. In the

South Atlantic¹⁰², the first phases of extension appear to be accompanied by faulting that bounds the Type I SDRs (Figure 2b). Instead, in the North Atlantic⁹⁶, SDRs are all akin to Type II and appear to be deposited during increasing subsidence from the magma injection site, without the intervention of faulting (Figure 4). Structural issues such as these will become critical if these volcanic sequences are required for future carbon sequestration.

In the South Atlantic, close to the Parana-Etendeka flood basalts, the thickness of the SDR packages is largest, conjugate margins are symmetric and the crust appears to have undergone large ductile deformation^{102,119}, probably related to crustal heating resulting from magmatic intrusion. With distance from the onshore flood basalts, the amount of offshore magmatism decreases¹¹⁶ and the margins become asymmetric, suggesting relatively stronger (but still ductile) rheologies, as the smaller magma input provided less crustal heating.

Transition to seafloor spreading. In the South Atlantic the COT appears to be formed by sub-aerial spreading under large magmatic input, generating the long Type II SDRs sections. Seafloor spreading probably started when the plume material beneath the extending axis waned as the ocean basin opened and the plume had more space to laterally distribute. Decreasing magmatic input would generate increasing subsidence below a water depth where extrusives would not laterally flow and form SDRs but form pillow lavas. The difference in style of the SDRs formed in the South and North Atlantic show that elevation during SDR emplacement may vary significantly depending on the pre-rift conditions (Figure 4).

4.3 Rifting at intermediate margins. Intermediate margins are characterised by an abrupt transition from thinned continental to classic Penrose-like oceanic crust. These margins differ greatly in crustal architecture and we focus here only in one example, the South China Sea, where full extension velocity of the first oceanic crust is 50 mm/yr, much larger than at magma-poor margins, thus more magmatism is expected. Extension in the South China Sea probably started in a wide rift mode, promoted by a hydrated and weak lower crust and mantle resulting from its setting in a former volcanic arc^{59,129,130}. Wide rifting would have promoted the formation of horst and grabens over ~200 km in the proximal margin (Figure 2c and HG in Figure 5d) until localisation due to ongoing cooling took place in the distal margin. Here extension started to migrate towards the basin centre, through a series of faults accompanied by ductile lower crustal deformation, which, in conjunction with increasing heat released by the intruded magmatism may have promoted the formation of the interpreted core complexes under ductile conditions in this area (D in Figure 2c, CCD in Figure 5d). As the crust and lithosphere thinned, magma reached shallower levels and it intruded in the form of dykes in addition to underplated gabbro. With increasing extension, the continental crust is dissected by several faults and dykes giving way to the formation of classic Penrose-like oceanic crust.

5. Importance for the energy transition

Rifted margins are massive repositories of sedimentary, igneous and ultra-mafic rocks, are globally distributed and lie adjacent to large, coastal populations. Thus, they are likely to hold a pivotal role in the future transition to the new green economy. In the offshore environment, where there is the legacy of drilling by the hydrocarbon industry, there is a potential for a greater acceptance of the level of future exploitation that will be needed for the energy transition¹⁹³.

Whilst current Carbon Capture and Storage (CCS) projects have demonstrated the feasibility of storing CO₂ underground, to meet IPCC model targets injection operations will need to be increased by several orders of magnitude¹⁹⁴. This will require exploration for new reservoirs and careful consideration of long-term storage security, especially for saline aquifers where the natural rates of mineralisation are sluggish. In this respect, magma-rich margins may prove especially promising¹⁹⁵ as they contain abundant, reactive rocks such as basalts, where injected CO₂ rapidly mineralizes to form stable carbonates, as has been proven for the onshore CarbFix (Iceland) and Wallula (USA) projects¹⁹⁶.

Similarly, large amounts of H₂ are expected to be produced at the COTs of magma-poor margins due to hydration of the exposed mantle rocks. Molecular hydrogen may become an important new fuel for internal combustion engines. Promising reserves have been detected onshore where sections of these COTs are now exposed in ophiolites, such as in the Mauleon Basin in the Pyrenees¹⁴. This setting portrays key ingredients for long-term trapping of H₂, which are: a seal, e.g. impermeable clay or salt; a kitchen, e.g. iron-rich rocks; an hydraulic system that favours water-rock interaction; and reservoir temperatures of ~100-200° C, where H₂ is relatively inert¹⁴. In the offshore environment, present-day H₂ fluxes at the COT and whether it can be trapped and form extractable reservoirs is still unknown.

Mineral deposits in the sedimentary sections of rifted margins form by hydrothermal water-rock interaction. For example, base metals accumulate where there is a transition from thick to thin lithosphere¹¹. This suggests that margins that form adjacent to cratonic lithosphere might be future reservoirs of base metals. In rifts, such as the Rhine Graben, the startup Lithium de France will start geothermal exploration to extract both heat and Lithium from reservoirs¹⁹⁷.

In all these examples, new understanding is needed of the fault-block scale interactions between fluid flow at short time scales and the long-time scale lithosphere deformation, melt transport, and sedimentation that create the conditions favourable for storage and extraction. 3D seismic imaging, which proved so critical for hydrocarbon exploration, will need to be applied to new targets. Process-based models can help reduce risk through fitting numerical models of coupled lithosphere and basin formation to observations to map the most likely reservoir system at a large scale. In addition, the goal of understanding a given margin at the scales needed for quantitative resource prediction can be pursued by fusing simulations of rift dynamics with geological and geophysical data; commonly known as inversion and data assimilation (DA). A model with adequate dynamics could, for example, assimilate MCS images. A first step in this direction is the heuristic

nudging approach used by⁶⁶ (Figure 6), which sequentially assimilates interpretations of MCS data, and shows the potential of DA towards understanding tectonics at fault-block scale. In the future, advanced DA approaches should be developed for geodynamic modelling of rifted margins.

6. Summary and future Perspectives

Rifted margins form by the interaction between tectonic, sedimentary, magmatic, hydrothermal and surface processes modulated by the initial lithospheric structure and extension velocity. This leads to a variety of architectures where magma-rich and -poor margins are only the end-members of a as yet poorly-defined spectrum. In this review we documented that even within end-member types there is a rich variety in syn-rift magmatism, margin width, conjugate asymmetry and COT nature. Drilling at the South China sea has shown that conceptual templates of how margins evolve have skewed the interpretation of margins towards end-member models. Similarly, the contrasting patterns of magmatism at the North and South Atlantic magma-rich margins, shows how established conceptual models can be incomplete.

Margin studies require reference to all geophysical and geological data available, and an understanding of how they complement each other. Geophysical surveys, in particular WAS surveys, which are not typically carried out by industrial partners, need to be more common and, when existent, considered by the rest of the community. For this, it is important that the different lateral and depth resolutions of MCS and WAS data and the different information they convey is properly understood (Box 1). In addition, in sectors where the basement is laterally heterogenous, such as the COTs, instrument spacing needs to be sufficiently small, and modelling approaches exploit that resolution. One way to increase resolution is by using MCS data in between OBS instrument location, and jointly invert both MCS and WAS⁷¹ and/or undertake downward continuation of both MCS and OBS data¹⁹⁸, and when instrument spacing is sufficiently close use higher resolution full waveform inversion⁸⁵.

Geodynamic models have been pivotal in documenting how different processes may explain margin architecture. Initial lithospheric strength is key in determining rift width and asymmetry, and the spatio-temporal distribution of magmatism, faulting and sedimentation. In numerical models, however, the link between the original lithospheric structure and rift geometries is established in a general way. How the structure and age of the initial fold-belt lithosphere affects rift architecture has only recently begun to be investigated¹⁹⁹. 3D modelling approaches are necessary to understand how rift development is linked to a broader tectonic plate perspective²⁰⁰⁻²⁰².

Petroleum geoscience matured over many decades, meanwhile we are only at the beginning of exploration for hydrogen and an expanding need for base and rare-earth metals and CO₂ storage. The source rocks, fluid flow migration pathways, and reservoirs for these resources will require a renewed focus on the interactions between sediments, deformation, fluid flow and magmatism. The fusion of data and models that

couple sedimentation, deformation, fluid flow and magmatism at sufficiently small scales will be pivotal to make future accurate predictions for these new storage and energy needs.

Figure captions

Figure 1. Global distribution of magma-rich, intermediate or uncertain and magma-poor margins.

Oceanic crust age from²⁰³. Seaward dipping reflectors, SDRs, and onshore large-igneous provinces, with ages <200 Ma, along with the ages of the onset of the main large-igneous province phase are shown²⁰⁴. Magmatic margins are distinguished by their SDRs and comprise: NA: North Atlantic margins, associated to the North Atlantic Igneous province, NAIP; the Eastern North American margin and its conjugate in Africa, associated with the Central Atlantic Magmatic province, CAMP (with the African side classified as uncertain²⁰⁵); the South Atlantic magmatic margins, SA, associated with the Paraná-Etendeka; the India-Seychelles margins associated with the Deccan magmatic province, and the NW Australian margin. The Gulf of Mexico is magmatic in the East but not in the West^{206,207}. Magma poor margins exhibit exhumed mantle at their continent-ocean transition. Margins where exhumed mantle at the COT has been interpreted from wide-angle data are shown. These include the West Iberia, WIM, and Newfoundland, NF, the Bay of Biscay, BB, the Goban Spur, GB, the Porcupine Basin, PB, Rockall Trough, RT, Orphan Basin, OB, Labrador, L, the southern West Greenland, WG, and the Thyrreanean, THY. The South Australian and its Antarctic conjugate have been typically considered as magma-poor²⁰⁸⁻²¹⁰ but wide-angle data has not confirmed exhumed mantle yet, so a question mark is included. Also shown are ODP drillings mentioned in the text, as black circles. Red lines are wide-angle data shown in Figure 3.

Figure 2. Conceptual models for a) magma-poor, b) magma-rich and c) intermediate margins based on the West Iberia-Newfoundland margins, the South Atlantic magma-rich margins and the South China sea. As mentioned in the main text, within any of these margin types there is structural variability in terms of amount of width and amount of magmatism within the COT, and margin width, along strike as well as between conjugate margins. SDRs: seaward dipping reflectors, HVLC: high-velocity lower crustal body, DR: deep reflectivity in the lower crust, L: listric faults, HG: Horst and grabens, PR: peridotite ridge. Note that the horizontal scale differs in each panel.

Figure 3. Wide-angle p-wave seismic velocity models across two contrasting conjugate continental margins. a) the magma-poor Newfoundland-West Iberia margins^{83,92} and b) the magma-rich of SE Greenland-Hatton Bank in the North Atlantic^{113,115} (location in Figure 1). Models were generated by inverting wide-angle seismic and seismic reflection data, the latter were used to constrain the sediment and basement properties. Velocities at the continent-ocean transitions, COT, of magma-poor margins are interpreted as exhumed and serpentinised mantle, with various degrees of magmatic contributions. Typical, 2-layered magmatic oceanic

crust is found in the seaward end of West Iberia but not in the Newfoundland section. Magma-rich margins exhibit a high-velocity lower crustal body, HVLC, at their base. The thickness of the HVLC and its structural position is asymmetric in this example, indicating asymmetry in melting and extension. Towards the oceanic crust the HVLC thins. The distance over which this thinning occurs before typical oceanic crust is observed, can be quite variable on different margins. In coincident MCS data it is shown that the HVLC is overlain by seaward dipping reflections, SDRs. In the SE Greenland-Hatton Bank profiles, the oceanic crust inception is defined by the C24r magnetic anomaly¹¹⁵.

Figure 4. Formation of Seaward-Dipping Reflectors (SDRs) in the South and North Atlantic (adapted from^{96,102}. **a)** The geometries of the lava flows (colored blue, orange and green) depends on the tectonics (formation of large-offset faults or not), magma supply rate and interaction with standing water (lakes or sea-water – shown by the wavy line). The characteristic tilt results from either faulting or subsidence. In the North Atlantic, as the lava sequence erupted, the elevation fell from above to below sea-level. This caused two distinct lava flow accumulations separated by a hyaloclastic outer high (colored purple). In the South Atlantic, the whole sequence appears to have been erupted while above sea-level with two packages (fault bounded colored orange and non-fault bounded colored green) stacked on-top of each other. **b)** The difference in land elevation may be explained by differences in crustal thickness (dotted blue line) at the time of plume arrival, with the amount of pre-existing thinning larger in the North Atlantic¹⁷⁷. The melt production, shown by red line, experienced a peak at different times during rifting, such that in the North Atlantic, the oceanic crust is thicker than the SDRs, whereas in the South Atlantic it is reverse¹⁷⁷.

Figure 5. Effect of decreasing initial lithospheric strength in tectonic, sedimentary architecture and melting. Extension velocity is 10 mm/yr, i.e. within the ultra-slow domain for oceanic spreading, and the mantle temperature is 1300°C, thus models apply for magma-poor to intermediate margins (see Suppl. text for model setup). The lithospheric strength decreases from **a)** to **d)** (see Figure S1 and Table S1). Models 1 to 3, share the same crustal and lithospheric thickness (32 km and 120 km, respectively) and rheological properties for upper (wet quartzite), lower crust (wet anorthite) and mantle (dry and wet olivine in the lithospheric and asthenospheric mantle, respectively). The Moho temperature is 540°C in **a)**, 570°C in **b)** and 770°C in **c)**. Model shown in **d)** is as model shown in **c)** but has a thicker crust of 35 km. Red shade is plastic/brittle strain rate and blue is ductile strain rate. Grey shading shows accumulated plastic/brittle strain and represents areas where faulting occurred. Sediments are color coded by age since the start of rifting. Dark orange areas are a prediction of the thickness and location of underplated magmatic products. In the model the magma is only underplated, but in nature it will intrude the crust and extrude it as dykes. With decreasing initial rheological strength, margins become wider and the style of faulting, sediment architecture and the amount and distribution

of underplated/intruded magma changes. DH: distal highs, L: Listric faults, CCD: Concave-downward detachments.

Figure 6. Sequential nudging approach to simulate rift dynamics at fault-block scale (adapted from⁶⁶).

Comparison of **a)** two conjugate MCS lines along the West Iberia-Newfoundland margins, with, **b)** results of the modelling technique, Kinedyn^{66,211}. The interpretation of fault kinematics in the MCS line is incorporated during the dynamic model run, so that the model is nudged towards the MCS data observations. Modelling at fault-block scale yields the sediment and basement temperature field during the syn and post-rift margin evolution. In this case, the model was used to understand how brittle and ductile deformation evolve with time, and their influence on detachment, D, formation, and the spatial distribution of serpentinisation. In the future, such models can be used to predict the pattern of hydrothermal circulation at margins and their consequences for element exchange and deposition of minerals within the sedimentary sections. F1 to F7 are faults, B1 to B6 are fault-blocks. Red-blue-white stripes show the deformation within the crust, blue thick line is the Moho from the model, DR are deep reflections. Shown are also dredges and ODP wells. Age of sediment is shown in colors. Model is shown after 115 Myr of evolution.

Figure 7. Conceptual margin cross section showing potential for economic resources and processes

During rifting and perhaps thereafter higher than normal geothermal gradients will lead to hydrothermal circulation, releasing lithium rich brines and base metals in the sedimentary sections (A and B). In addition, sediments may host oil and gas reservoirs and may be used also as sites for CO₂ storage (C). Turbidites may pose a hazard for marine infrastructure and at the same time host unconventional oil and gas reservoirs (D). Finally, native H₂ produced during mantle hydration may be found in reservoirs which could be potentially used as energy sources (E). 1) Proximal margins and necking zone, 2) distal margin, 3) outer margin, 4) exhumed mantle, 5) oceanic crust. Note that magma-rich margins will look different and will host potential for CO₂ storage in their SDRs sequences.

Box 1 | Interpreting seismic data

Seismic data typically falls into two categories – “multi-channel” seismic (MCS) where both the seismic source and receiver are surface towed, and mainly records near-normal incidence reflected waves, and “wide-angle” seismic (WAS) where the source is surface towed and the receiver placed on the seabed, and records wide-angle refracted waves and reflected waves. WAS is currently the only technology to provide velocity-depth information across the full depth extent needed for continental margin studies.

In recent years MCS data imaging has been revolutionized by using longer hydrophone streamers and improved air gun technology, increasing our ability to suppress multiples and so image reflectivity at crustal

scale. However, MCS images are often shown in two-way-travel times which gives a highly distorted impression due to highly variable low-velocity water and sediment layers across a margin. Pre-stack depth migration (PSDM) was developed to address this issue, but it lacks sensitivity at depths larger than the streamer length²¹². This means that coincident PSDM and wide-angle sections, may exhibit up to some kms of difference in the depth of deep reflectors such as the Moho⁵⁵. In addition, wide-angle seismic models sample bulk properties, thus complex reflectivity seen on co-incident MCS lines which do not present a bulk property change, will not be observed in wide-angle data.

Although, absolute compressional wave velocities, V_p , of rocks, provided by wide-angle data do not intrinsically differentiate between rock type as their values overlap in certain velocity ranges, their V_p gradients allow to distinguish basement nature (Figure B1 and Figure B2). Over the years, mapping of these V_p gradients has continuously improved with increasing quality of field data and modelling technologies. Closely spaced OBS (<10 km) in conjunction with joint inversion of coincident MCS and wide-angle data provides to date, the most accurate depiction of velocity-depth structure and Moho depth⁷¹.

Continental domains are readily distinguished from oceanic domains based on velocity-depth structure (Figure B2). At magma-rich margins HVLC bodies are readily detected as they have absolute velocities higher than the surrounding lower continental rocks ($7.0 < V_p < 7.8$). Analysis of their velocities have been shown to be consistent with their volumes and is controlled by the depth and temperature of melting^{98,213,214}. Oceanic crust generally displays a two-layer structure, although at slow-spreading the contrast between the upper steep gradient layer and the lower gentle velocity gradient layer can be softened by the inclusion of a proportion of serpentinised mantle rocks^{215,216}. At ultra-slow oceanic spreading environments (< ~12 mm/yr spreading velocity) the oceanic crust may disappear altogether so that the mantle is exhumed on the seafloor and a Moho is missing^{217,218}. Care is therefore needed when interpreting lithology in such settings and across the COT of magma-poor margins.

Glossary

Breakup: The point (in time and space) where the thinning continents physically separate from each other.

COT: Continent-ocean transition. The area seawards of the thinned continental crust, which does not show seismic velocity-depth nor tectonic structures which are typical of thinned continental or oceanic crust.

Conjugate margins: Two sides of an ocean basin that before rifting were joined.

Footwall: The block that doesn't experience subsidence on one side of an active fault.

Hangingwall: The block that experiences subsidence on one side of an active fault.

HVLC: High velocity lower crustal bodies (V_p of 7.3-7.8 km/s) observed in wide-angle seismic data along some margins.

LIP: Large igneous provinces formed by the injection of large volumes of magmatism in very short geological periods, 1-2 Myr or less. Traditionally interpreted as formed by the impact of a new mantle plume.

Listric: A fault that has high-angle in its shallowest segments (~60-30°) and has much lower angle at depth.

MCS: Multichannel seismic data, also known as reflection seismics. Distance between the source and receiver is fixed. Provides high-resolution image but limited velocity information.

Magnetic anomaly lineations: Commonly interpreted to indicate seafloor spreading (oceanic crust). Formed when mafic magma cools below the Curie Point (580°C) and takes the polarity of the Earth's field at that time.

Necking zone: Where the crust thins from the continental platform (crustal thickness 28-25 km) towards the distal domain (crustal thickness of ~20-15 km or less). Necking zones can be spatially abrupt or occur over a large distance.

Post-rift: Period of time after breakup.

Post-tectonic sediment: sediment deposited after the activity of the underlying basement faults.

SDRs: Seaward-dipping reflectors. Formed of tilted, stacked, lava and sediment interbeds.

Sequential faulting: A system of faults that young oceanward and cut through the hangingwall of the previous ones.

Strain weakening: A reduction in the strength of the lithosphere due to mechanical damage. Strain weakening may result from the presence of fluids at faults, mineralization, reduction in grain size and the formation of crystallographic preferred orientations.

Syn-rift: Period of time before breakup.

Syn-tectonic sediment: sediment deposited during the activity of the underlying basement faults.

Unconformity: A boundary between sedimentary rocks caused by a period of erosion or a pause in sediment accumulation.

Underplating – igneous material added to the base of the crust.

WAS: Wide-angle seismic data, also known as refraction seismics. Distance between the source and receiver varies. Provides low-resolution image but important velocity information.

718 1 Dunkley Jones, T., Fauth, G. & LeVay, L. J. Expedition 388 scientific prospectus: Equatorial Atlantic
719 gateway. *Scientific Prospectuses International Ocean Discovery Program*,
720 doi:10.14379/iodp.sp.388.2019 (2019).

721 2 Planke, S., Berndt, C. & Alvarez Zarikian, C. A. Expedition 396 scientific prospectus: Mid-Norwegian
722 continental margin magmatism. *Scientific Prospectuses International Ocean Discovery Program*,
723 doi:10.14379/iodp.sp.396.2021 (2021).

724 3 Brune, S., Williams, S. E. & Müller, R. D. Potential links between continental rifting, CO₂ degassing
725 and climate change through time. *Nat Geosci* **10**, 941-946, doi:10.1038/s41561-017-0003-6 (2017).

726 4 Svensen, H. *et al.* Release of methane from a volcanic basin as a mechanism for initial Eocene global
727 warming. *Nature* **429**, 542-545, doi:10.1038/nature02566 (2004).

728 5 Pérez-Gussinyé, M. *et al.* Serpentinization and magmatism during extension at non-volcanic
729 margins: the effect of initial lithospheric structure. *Geological Society London, Special Publications*
730 **187**, doi:10.1144/gsl.Sp.2001.187.01.27 (2001).

731 6 Bayrakci, G. *et al.* Fault-controlled hydration of the upper mantle during continental rifting. *Nat*
732 *Geosci* **9**, 384-388, doi:10.1038/ngeo2671 (2016).

733 7 Albers, E., Bach, W., Pérez-Gussinyé, M., McCammon, C. & Frederichs, T. Serpentinization-driven H₂
734 production from continental break-up to mid-ocean ridge spreading: Unexpected high rates at the
735 West Iberia margin. *Front Earth Sc-Switz* **9**, doi:10.3389/feart.2021.673063 (2021).

736 8 Schwarzenbach, E. M., Früh-Green, G. L., Bernasconi, S. M., Alt, J. C. & Plas, A. Serpentinization and
737 carbon sequestration: A study of two ancient peridotite-hosted hydrothermal systems. *Chem Geol*
738 **351**, 115-133, doi:10.1016/j.chemgeo.2013.05.016 (2013).

739 9 Kelley, D. S. *et al.* A serpentinite-hosted ecosystem: The Lost City hydrothermal field. *Science* **307**,
740 1428-1434, doi:10.1126/science.1102556 (2005).

741 10 Levell, B., Argent, J., Dore, A. G. & Fraser, S. in *Petroleum Geology: From Mature Basins to New*
742 *Frontiers - Proceedings of the 7th Petroleum Geology Conference*. 823-830.

743 11 Hoggard, M. J. *et al.* Global distribution of sediment-hosted metals controlled by craton edge
744 stability. *Nat Geosci* **13**, doi:10.1038/s41561-020-0593-2 (2020).

745 12 Hitzman, M. W., Selley, D. & Bull, S. Formation of sedimentary rock-hosted stratiform copper
746 deposits through Earth history. *Economic Geology* **105**, 627-639, doi:10.2113/gsecongeo.105.3.627
747 (2010).

748 13 Wilkinson, J. J. in *Treatise on Geochemistry (Second Edition)* Vol. 13 (eds Heinrich D. Holland & Karl
749 K. Turekian) 219-249 (Elsevier, 2014).

750 14 Lefeuvre, N. *et al.* Native H₂ exploration in the western Pyrenean foothills. *Geochemistry,*
751 *Geophysics, Geosystems* **22**, doi:10.1029/2021GC009917 (2021).

752 15 Kelemen, P. B. & Matter, J. In situ carbonation of peridotite for CO₂ storage. *Proceedings of the*
753 *National Academy of Sciences* **105**, 17295-17300, doi:10.1073/pnas.0805794105 (2008).

754 16 Jolie, E. *et al.* Geological controls on geothermal resources for power generation. *Nature Reviews*
755 *Earth & Environment* **2**, 324-339, doi:10.1038/s43017-021-00154-y (2021).

756 17 Freymark, J. *et al.* The deep thermal field of the Upper Rhine Graben. *Tectonophysics* **694**, 114-129,
757 doi:10.1016/j.tecto.2016.11.013 (2017).

758 18 McKenzie, D. Some remarks on the development of sedimentary basins. *Earth and Planetary Science*
759 *Letters* **40**, 35-32 (1978).

760 19 Karner, G. D. & Watts, A. B. On isostasy at Atlantic-type continental margins. *J Geophys Res-Sol Ea*
761 **87**, 2923-2948, doi:10.1029/JB087iB04p02923 (1982).

762 20 Jarvis, G. T. & McKenzie, D. P. Sedimentary basin formation with finite extension rates. *Earth and*
763 *Planetary Science Letters* **48**, 42-52, doi:10.1016/0012-821X(80)90168-5 (1980).

764 21 Cochran, J. R. Effects of finite rifting times on the development of sedimentary basins. *Earth and*
765 *Planetary Science Letters* **66**, 289-302, doi:10.1016/0012-821X(83)90142-5 (1983).

766 22 Anonymous. Penrose field conference on ophiolites. *Geotimes* **17**, 24-25 (1972).

767 23 Christeson, G. L., Goff, J. A. & Reece, R. S. Synthesis of oceanic crustal structure from two-
768 dimensional seismic profiles. *Rev Geophys* **57**, 504-529, doi:10.1029/2019rg000641 (2019).

769 24 White, R. & Mckenzie, D. Magmatism at rift zones - the generation of volcanic continental margins
770 and flood basalts. *J Geophys Res-Solid* **94**, 7685-7729, doi:10.1029/JB094iB06p07685 (1989).

771 25 Sawyer, D. S., Coffin, M. F., Reston, T. J., Stock, J. M. & Hopper, J. R. COBBOOM: The continental
772 breakup and birth of oceans mission. *Scientific Drilling* **5**, 13-25, doi:10.2204/iodp.sd.5.02.2007
773 (2007).

774 26 Manatschal, G. *et al.* The role of inheritance in forming rifts and rifted margins and building
775 collisional orogens: a Biscay-Pyrenean perspective. *B Soc Geol Fr* **192**, doi:10.1051/bsgf/2021042
776 (2021).

777 27 Armitage, J. J., Collier, J. S. & Minshull, T. A. The importance of rift history for volcanic margin
778 formation. *Nature* **465**, 913-917, doi:10.1038/nature09063 (2010).

779 28 Buck, W. R. Modes of continental lithospheric extension. *J Geophys Res-Sol Ea* **96**, 20161-20178,
780 doi:10.1029/91jb01485 (1991).

781 29 Boillot, G. & Winterer, E. L. in *Proceedings of the Ocean Drilling Program, Scientific Results* Vol. 103
782 (eds G Boillot, E. L. Winterer, & A. W Meyer) 809–828 (Ocean Drilling Program, 1988).

783 30 Whitmarsh, R. B. & Sawyer, D. S. in *Proceedings of the Ocean Drilling Program, Scientific Results* Vol.
784 149 (eds R.B. Whitmarsh, D.S. Sawyer, A. Klaus, & D.G. Masson) 713–733 (Ocean Drilling Program,
785 1996).

786 31 Whitmarsh, R. B. & Wallace, P. J. in *Proceedings of the Ocean Drilling Program, Scientific Results* Vol.
787 173 (eds M.-O. Beslier, R.B. Whitmarsh, P.J. Wallace, & J. Girardeau) 1–36 (Ocean Drilling Program,
788 2001).

789 32 Tucholke, B. E. & Sibuet, J.-C. in *Proceedings of the Ocean Drilling Program, Scientific Results* Vol.
790 210 (eds B.E. Tucholke, J.-C. Sibuet, & A. Klaus) 1-56 (Ocean Drilling Program, 2007).

791 33 Hinz, K. A hypothesis on terrestrial catastrophes: wedges of very thick oceanward dipping layers
792 beneath passive continental margins—their origin and paleoenvironmental significance.
793 *Geologisches Jahrbuch, Reihe E, Geophysik* **22**, 3-28 (1981).

794 34 Mutter, J. C., Talwani, M. & Stoffa, P. L. Origin of seaward-dipping reflectors in oceanic-crust off the
795 Norwegian margin by subaerial sea-floor spreading. *Geology* **10**, 353-357, doi:10.1130/0091-
796 7613(1982)10<353:Oosrio>2.0.Co;2 (1982).

797 35 Roberts, D. G., Backman, J., Morton, A. C., Murray, J. W. & Keene, J. B. Evolution of volcanic rifted
798 margins - Synthesis of Leg-81 results on the west margin of Rockall Plateau. *Initial Rep Deep Sea* **81**,
799 883-911, doi:10.2973/dsdp.proc.81.139.1984 (1984).

800 36 Eldholm, O., Thiede, J. & Taylor, E. in *Proceedings of the Ocean Drilling Program, Scientific Results*
801 Vol. 104 *Proceedings of the Ocean Drilling Program* (eds O Eldholm *et al.*) 1033–1065 (1989).

802 37 Larsen, H. C. & Saunders, A. D. in *Proceedings of the Ocean Drilling Program, Scientific Results* Vol.
803 152 *Proceedings of the Ocean Drilling Program* (eds A.D. Saunders, H.C. Larsen, & S.W. Wise) 503–
804 533 (Ocean Drilling Program, 1998).

805 38 White, R. S. *et al.* Magmatism at rifted continental margins. *Nature* **330**, 439-444,
806 doi:10.1038/330439a0 (1987).

807 39 White, R. S., Mckenzie, D. & Onions, R. K. Oceanic crustal thickness from seismic measurements and
808 rare-Earth element inversions. *J Geophys Res-Sol Ea* **97**, 19683-19715, doi:10.1029/92jb01749
809 (1992).

810 40 Funck, T. *et al.* A review of the NE Atlantic conjugate margins based on seismic refraction data.
811 *Geological Society London, Special Publications* **447**, 171-205, doi:10.1144/sp447.9 (2017).

812 41 Coffin, M. F. & Eldholm, O. Large Igneous Provinces - Crustal structure, dimensions, and external
813 consequences. *Rev Geophys* **32**, 1-36, doi:10.1029/93rg02508 (1994).

- 42 Courtillot, V., Jaupart, C., Manighetti, I., Tapponnier, P. & Besse, J. On causal links between flood basalts and continental breakup. *Earth and Planetary Science Letters* **166**, 177-195, doi:10.1016/S0012-821x(98)00282-9 (1999).
- 43 Peron-Pinvidic, G., Manatschal, G. & Osmundsen, P. T. Structural comparison of archetypal Atlantic rifted margins: A review of observations and concepts. *Marine and Petroleum Geology* **43**, 21-47, doi:10.1016/j.marpetgeo.2013.02.002 (2013).
- 44 Franke, D. Rifting, lithosphere breakup and volcanism: Comparison of magma-poor and volcanic rifted margins. *Marine and Petroleum Geology* **43**, 63-87, doi:10.1016/j.marpetgeo.2012.11.003 (2013).
- 45 Hauptert, I., Manatschal, G., Decarlis, A. & Unternehr, P. Upper-plate magma-poor rifted margins: Stratigraphic architecture and structural evolution. *Marine and Petroleum Geology* **69**, 241-261, doi:10.1016/j.marpetgeo.2015.10.020 (2016).
- 46 Manatschal, G., Sutra, E. & Péron-Pinvidic, G. in *2nd Central & North Atlantic Conjugate Margins: Rediscovering the Atlantic, New insights, New Winds for an Old Sea*.
- 47 Froitzheim, N. & Manatschal, G. Kinematics of Jurassic rifting, mantle exhumation, and passive-margin formation in the Austroalpine and Penninic nappes (eastern Switzerland). *GSA Bulletin* **108**, 1120-1133, doi:10.1130/0016-7606(1996)108<1120:Kojrme>2.3.Co;2 (1996).
- 48 Whitmarsh, R. B., Manatschal, G. & Minshull, T. A. Evolution of magma-poor continental margins from rifting to seafloor spreading. *Nature* **413**, 150-154, doi:10.1038/35093085 (2001).
- 49 Harkin, C., Kusznir, N., Tugend, J., Manatschal, G. & McDermott, K. Evaluating magmatic additions at a magma-poor rifted margin: an East Indian case study. *Geophys J Int* **217**, 25-40, doi:10.1093/gji/ggz007 (2019).
- 50 Tugend, J. *et al.* Reappraisal of the magma-rich versus magma-poor rifted margin archetypes. *Geological Society London, Special Publications* **476**, doi:10.1144/sp476.9 (2020).
- 51 Taylor, B., Goodliffe, A., Martinez, F. & Hey, R. Continental rifting and initial sea-floor spreading in the Woodlark Basin. *Nature* **374**, 534-537, doi:10.1038/374534a0 (1995).
- 52 Taylor, B., Goodliffe, A. M. & Martinez, F. How continents break up: Insights from Papua New Guinea. *J Geophys Res-Sol Ea* **104**, 7497-7512, doi:10.1029/1998jb900115 (1999).
- 53 Loureiro, A. *et al.* Imaging exhumed lower continental crust in the distal Jequitinhonha basin, Brazil. *Journal of South American Earth Sciences* **84**, 351-372, doi:10.1016/j.jsames.2018.01.009 (2018).
- 54 Pinheiro, J. M. *et al.* Lithospheric structuration onshore-offshore of the Sergipe-Alagoas passive margin, NE Brazil, based on wide-angle seismic data. *Journal of South American Earth Sciences* **88**, 649-672, doi:10.1016/j.jsames.2018.09.015 (2018).
- 55 Evain, M. *et al.* Deep structure of the Santos Basin-Sao Paulo Plateau System, SE Brazil. *J Geophys Res-Sol Ea* **120**, 5401-5431, doi:10.1002/2014jb011561 (2015).
- 56 Lizarralde, D. *et al.* Variation in styles of rifting in the Gulf of California. *Nature* **448**, 466-469, doi:10.1038/nature06035 (2007).
- 57 Merino, I., Prada, M., Ranero, C. R., Sallarès, V. & Calahorrano, A. The structure of the continent-ocean transition in the Gulf of Lions from joint refraction and reflection travel-time tomography. *J Geophys Res-Sol Ea* **126**, doi:10.1029/2021JB021711 (2021).
- 58 Ligi, M. *et al.* Birth of an ocean in the Red Sea: Initial pangs. *Geochemistry, Geophysics, Geosystems* **13**, doi:10.1029/2012GC004155 (2012).
- 59 Cameselle, A. L., Ranero, C. R. & Barckhausen, U. Understanding the 3D formation of a wide rift: The central South China Sea rift system. *Tectonics* **39**, doi:10.1029/2019TC006040 (2020).
- 60 Cameselle, A. L., Ranero, C. R., Franke, D. & Barckhausen, U. The continent-ocean transition on the northwestern South China Sea. *Basin Res* **29**, 73-95, doi:10.1111/bre.12137 (2017).
- 61 Larsen, H. C. *et al.* Rapid transition from continental breakup to igneous oceanic crust in the South China Sea. *Nat Geosci* **11**, 782-789, doi:10.1038/s41561-018-0198-1 (2018).

862 62 Sun, Z. *et al.* South China Sea rifted margin. *Scientific Prospectuses International Ocean Discovery*
863 *Program* **367/368**, doi:10.14379/iodp.proc.367368.2018 (2018).

864 63 Lymer, G. *et al.* 3D development of detachment faulting during continental breakup. *Earth and*
865 *Planetary Science Letters* **515**, 90-99, doi:10.1016/j.epsl.2019.03.018 (2019).

866 64 Reston, T. J., Krawczyk, C. M. & Klaeschen, D. The S reflector west of Galicia (Spain): Evidence from
867 prestack depth migration for detachment faulting during continental breakup. *J Geophys Res-Sol Ea*
868 **101**, 8075-8091, doi:10.1029/95jb03466 (1996).

869 65 Davy, R. G. *et al.* Continental hyperextension, mantle exhumation, and thin oceanic crust at the
870 continent-ocean transition, West Iberia: New insights from wide-angle seismic. *J Geophys Res-Sol Ea*
871 **121**, 3177-3199, doi:10.1002/2016JB012825 (2016).

872 66 Liu, Z. *et al.* Lateral coexistence of ductile and brittle deformation shapes magma-poor distal
873 margins: An example from the West Iberia-Newfoundland margins. *Earth and Planetary Science*
874 *Letters* **578**, 117288, doi:10.1016/j.epsl.2021.117288 (2022).

875 67 Thinon, I. *et al.* Deep structure of the Armorican Basin (Bay of Biscay): a review of Norgasis seismic
876 reflection and refraction data. *J Geol Soc London* **160**, 99-116, doi:10.1144/0016-764901-103 (2003).

877 68 de Charpal, O., Guennoc, P., Montadert, L. & Roberts, D. G. Rifting, crustal attenuation and
878 subsidence in the Bay of Biscay. *Nature* **275**, 706-711, doi:10.1038/275706a0 (1978).

879 69 Krawczyk, C. M., Reston, T. J., Boillot, G. & Beslier, M.-O. in *Proceedings of the Ocean Drilling*
880 *Program, Scientific Results* Vol. 149 603-615 (1996).

881 70 Afilhado, A. *et al.* From unthinned continent to ocean: The deep structure of the West Iberia passive
882 continental margin at 38°N. *Tectonophysics* **458**, 9-50, doi:10.1016/j.tecto.2008.03.002 (2008).

883 71 Merino, I., Ranero, C. R., Prada, M., Sallarès, V. & Grevemeyer, I. The rift and continent-ocean
884 transition structure under the Tagus Abyssal Plain west of the Iberia. *J Geophys Res-Sol Ea* **126**,
885 doi:10.1029/2021JB022629 (2021).

886 72 Keen, C. E., Dickie, K. & Dafoe, L. T. Structural evolution of the rifted margin off northern Labrador:
887 The role of hyperextension and magmatism. *Tectonics* **37**, 1955-1972, doi:10.1029/2017TC004924
888 (2018).

889 73 Hopper, J. R. *et al.* Continental breakup and the onset of ultraslow seafloor spreading off Flemish
890 Cap on the Newfoundland rifted margin. *Geology* **32**, 93-96, doi:10.1130/G19694.1 (2004).

891 74 Ranero, C. R. & Perez-Gussinye, M. Sequential faulting explains the asymmetry and extension
892 discrepancy of conjugate margins. *Nature* **468**, 294-299, doi:10.1038/nature09520 (2010).

893 75 Van Avendonk, H. J. A. *et al.* Seismic velocity structure of the rifted margin of the eastern Grand
894 Banks of Newfoundland, Canada. *J Geophys Res-Sol Ea* **111** (2006).

895 76 Dean, S. M., Minshull, T. A., Whitmarsh, R. B. & Loudon, K. E. Deep structure of the ocean-continent
896 transition in the southern Iberia Abyssal Plain from seismic refraction profiles: The IAM-9 transect at
897 40 degrees 20 ' N. *J Geophys Res-Sol Ea* **105**, 5859-5885, doi:10.1029/1999jb900301 (2000).

898 77 Reston, T. J., Pennell, J., Stubenrauch, A., Walker, I. & Perez-Gussinye, M. Detachment faulting,
899 mantle serpentinization, and serpentinite-mud volcanism beneath the Porcupine Basin, southwest
900 of Ireland. *Geology* **29**, 587-590, doi:10.1130/0091-7613(2001)029<0587:Dfmsas>2.0.Co;2 (2001).

901 78 Reston, T. J. & Perez-Gussinye, M. Lithospheric extension from rifting to continental breakup at
902 magma-poor margins: rheology, serpentinisation and symmetry. *Int J Earth Sci* **96**, 1033-1046,
903 doi:10.1007/s00531-006-0161-z (2007).

904 79 Pérez-Gussinyé, M., Ranero, C. R., Reston, T. J. & Sawyer, D. Mechanisms of extension at
905 nonvolcanic margins: Evidence from the Galicia interior basin, west of Iberia. *J Geophys Res-Sol Ea*
906 **108**, doi:10.1029/2001jb000901 (2003).

907 80 Minshull, T. A., Dean, S. M. & Whitmarsh, R. B. The peridotite ridge province in the southern Iberia
908 Abyssal Plain: Seismic constraints revisited. *J Geophys Res-Sol Ea* **119**, 1580-1598,
909 doi:10.1002/2014jb011011 (2014).

910 81 Grevenmeyer, I., Ranero, C. R. & Ivandic, M. Structure of oceanic crust and serpentinization at
911 subduction trenches. *Geosphere* **14**, 395-418, doi:10.1130/Ges01537.1 (2018).

912 82 Watanabe, T., Kasami, H. & Ohshima, S. Compressional and shear wave velocities of serpentinized
913 peridotites up to 200 MPa. *Earth, Planets and Space* **59**, 233-244, doi:10.1186/BF03353100 (2007).

914 83 Grevenmeyer, I. *et al.* The continent-to-ocean transition in the Iberia Abyssal Plain. *Geology* **50**, 615-
915 619, doi:10.1130/g49753.1 (2022).

916 84 Lau, K. W. H., Nedimović, M. R. & Louden, K. E. Continent-ocean transition across the northeastern
917 Nova Scotian margin from a dense wide-angle seismic profile. *J Geophys Res-Sol Ea* **123**, 4331-4359,
918 doi:10.1029/2017JB015282 (2018).

919 85 Jian, H., Nedimović, M. R., Canales, J. P. & Lau, K. W. H. New insights into the rift to drift transition
920 across the northeastern Nova Scotian margin from wide-angle seismic waveform inversion and
921 reflection imaging. *J Geophys Res-Sol Ea* **126**, doi:10.1029/2021JB022201 (2021).

922 86 Chian, D. P., Louden, K. E. & Reid, I. Crustal structure of the Labrador Sea conjugate margin and
923 implications for the formation of nonvolcanic continental margins. *J Geophys Res-Sol Ea* **100**, 24239-
924 24253, doi:10.1029/95jb02162 (1995).

925 87 Welford, J. K., Dehler, S. A. & Funck, T. Crustal velocity structure across the Orphan Basin and
926 Orphan Knoll to the continent–ocean transition, offshore Newfoundland, Canada. *Geophys J Int* **221**,
927 37-59, doi:10.1093/gji/ggz575 (2019).

928 88 Prada, M. *et al.* Seismic structure of the Central Tyrrhenian basin: Geophysical constraints on the
929 nature of the main crustal domains. *J Geophys Res-Sol Ea* **119**, 52-70, doi:10.1002/2013JB010527
930 (2014).

931 89 Prada, M. *et al.* The complex 3-D transition from continental crust to backarc magmatism and
932 exhumed mantle in the Central Tyrrhenian basin. *Geophys J Int* **203**, 63-78, doi:10.1093/gji/ggv271
933 (2015).

934 90 Hopper, J. R. *et al.* A deep seismic investigation of the Flemish Cap margin: implications for the
935 origin of deep reflectivity and evidence for asymmetric break-up between Newfoundland and
936 Iberia. *Geophys J Int* **164**, 501-515, doi:10.1111/j.1365-246X.2006.02800.x (2006).

937 91 Bullock, A. D. & Minshull, T. A. From continental extension to seafloor spreading: crustal structure of
938 the Goban Spur rifted margin, southwest of the UK. *Geophys J Int* **163**, 527-546, doi:10.1111/j.1365-
939 246X.2005.02726.x (2005).

940 92 Eddy, D. R., Van Avendonk, H. J. A. & Shillington, D. J. Compressional and shear-wave velocity
941 structure of the continent-ocean transition zone at the eastern Grand Banks, Newfoundland.
942 *Geophys Res Lett* **40**, 3014-3020, doi:10.1002/grl.50511 (2013).

943 93 Rooney, T. O. The Cenozoic magmatism of East-Africa: Part I - Flood basalts and pulsed magmatism.
944 *Lithos* **286**, 264-301, doi:10.1016/j.lithos.2017.05.014 (2017).

945 94 Shillington, D. *et al.* Controls on rift faulting in the North Basin of the Malawi (Nyasa) Rift, East
946 Africa. *Tectonics* **39**, doi:10.1029/2019TC005633 (2020).

947 95 Scholz, C. A. *et al.* Intrarift fault fabric, segmentation, and basin evolution of the Lake Malawi
948 (Nyasa) Rift, East Africa. *Geosphere* **16**, 1293-1311, doi:10.1130/ges02228.1 (2020).

949 96 Planke, S., Symonds, P. A., Alvestad, E. & Skogseid, J. Seismic volcanostratigraphy of large-volume
950 basaltic extrusive complexes on rifted margins. *J Geophys Res-Sol Ea* **105**, 19335-19351,
951 doi:10.1029/1999jb900005 (2000).

952 97 Berndt, C., Planke, S., Alvestad, E., Tsikalas, F. & Rasmussen, T. Seismic volcanostratigraphy of the
953 Norwegian Margin: constraints on tectonomagmatic break-up processes. *J Geol Soc London* **158**,
954 413-426, doi:10.1144/jgs.158.3.413 (2001).

955 98 White, R. S. *et al.* Lower-crustal intrusion on the North Atlantic continental margin. *Nature* **452**, 460-
956 464, doi:10.1038/nature06687 (2008).

957 99 Peate, I. U., Larsen, M. & Leshner, C. L. The transition from sedimentation to flood volcanism in the
958 Kangerlussuaq Basin, East Greenland: basaltic pyroclastic volcanism during initial Palaeogene
959 continental break-up. *J Geol Soc London* **160**, 759 - 772 (2003).

960 100 Paton, D. A., Pindell, J., McDermott, K., Bellingham, P. & Horn, B. Evolution of seaward-dipping
961 reflectors at the onset of oceanic crust formation at volcanic passive margins: Insights from the
962 South Atlantic. *Geology* **45**, 439-442, doi:10.1130/G38706.1 (2017).

963 101 McDermott, C., Collier, J. S., Lonergan, L., Fruehn, J. & Bellingham, P. Seismic velocity structure of
964 seaward-dipping reflectors on the South American continental margin. *Earth and Planetary Science
965 Letters* **521**, 14-24, doi:10.1016/j.epsl.2019.05.049 (2019).

966 102 McDermott, C., Lonergan, L., Collier, J. S., McDermott, K. G. & Bellingham, P. Characterization of
967 seaward-dipping reflectors along the South American Atlantic margin and implications for
968 continental breakup. *Tectonics* **37**, 3303-3327, doi:10.1029/2017tc004923 (2018).

969 103 Stica, J. M., Zalan, P. V. & Ferrari, A. L. The evolution of rifting on the volcanic margin of the Pelotas
970 Basin and the contextualization of the Parana-Etendeka LIP in the separation of Gondwana in the
971 South Atlantic. *Marine and Petroleum Geology* **50**, 1-21, doi:10.1016/j.marpetgeo.2013.10.015
972 (2014).

973 104 Bauer, K. *et al.* Deep structure of the Namibia continental margin as derived from integrated
974 geophysical studies. *J Geophys Res-Sol Ea* **105**, 25829-25853, doi:10.1029/2000jb900227 (2000).

975 105 Holbrook, W. S. *et al.* Mantle thermal structure and active upwelling during continental breakup in
976 the North Atlantic. *Earth and Planetary Science Letters* **190**, 251-266, doi:10.1016/S0012-
977 821x(01)00392-2 (2001).

978 106 Breivik, A. J., Faleide, J. I., Mjelde, R. & Flueh, E. R. Magma productivity and early seafloor spreading
979 rate correlation on the northern Voring Margin, Norway - Constraints on mantle melting.
980 *Tectonophysics* **468**, 206-223, doi:10.1016/j.tecto.2008.09.020 (2009).

981 107 Voss, M., Schmidt-Aursch, M. C. & Jokat, W. Variations in magmatic processes along the East
982 Greenland volcanic margin. *Geophys J Int* **177**, 755-782, doi:10.1111/j.1365-246X.2009.04077.x
983 (2009).

984 108 Clerc, C., Jolivet, L. & Ringenbach, J. C. Ductile extensional shear zones in the lower crust of a
985 passive margin. *Earth and Planetary Science Letters* **431**, 1-7, doi:10.1016/j.epsl.2015.08.038 (2015).

986 109 Sapin, F., Ringenbach, J. C. & Clerc, C. Rifted margins classification and forcing parameters. *Scientific
987 Reports* **11**, doi:10.1038/s41598-021-87648-3 (2021).

988 110 Walker, G. P. L. Zeolite zones and dike distribution in relation to the structure of the basalts of
989 eastern Iceland. *The Journal of Geology* **68**, 515-528, doi:10.1086/626685 (1960).

990 111 Buck, W. R. The role of magmatic loads and rift jumps in generating seaward dipping reflectors on
991 volcanic rifted margins. *Earth and Planetary Science Letters* **466**, 62-69,
992 doi:10.1016/j.epsl.2017.02.041 (2017).

993 112 Morgan, R. L. & Watts, A. B. Seismic and gravity constraints on flexural models for the origin of
994 seaward dipping reflectors. *Geophys J Int* **214**, 2073-2083, doi:10.1093/gji/ggy243 (2018).

995 113 White, R. S. & Smith, L. K. Crustal structure of the Hatton and the conjugate east Greenland rifted
996 volcanic continental margins, NE Atlantic. *J Geophys Res-Sol Ea* **114**, doi:10.1029/2008jb005856
997 (2009).

998 114 Collier, J. S. *et al.* New constraints on the age and style of continental breakup in the South Atlantic
999 from magnetic anomaly data. *Earth and Planetary Science Letters* **477**, 27-40,
1000 doi:10.1016/j.epsl.2017.08.007 (2017).

1001 115 Hopper, J. R. *et al.* Structure of the SE Greenland margin from seismic reflection and refraction data:
1002 Implications for nascent spreading center subsidence and asymmetric crustal accretion during North
1003 Atlantic opening. *J Geophys Res-Sol Ea* **108**, doi:10.1029/2002jb001996 (2003).

1004 116 Taposeea, C. A., Armitage, J. J. & Collier, J. S. Asthenosphere and lithosphere structure controls on
1005 early onset oceanic crust production in the southern South Atlantic. *Tectonophysics* **716**, 4-20,
1006 doi:10.1016/j.tecto.2016.06.026 (2017).

1007 117 Becker, K. *et al.* Asymmetry of high-velocity lower crust on the South Atlantic rifted margins and
1008 implications for the interplay of magmatism and tectonics in continental breakup. *Solid Earth* **5**,
1009 1011-1026, doi:10.5194/se-5-1011-2014 (2014).

1010 118 Koopmann, H. *et al.* Segmentation and volcano-tectonic characteristics along the SW African
1011 continental margin, South Atlantic, as derived from multichannel seismic and potential field data.
1012 *Marine and Petroleum Geology* **50**, 22-39, doi:10.1016/j.marpetgeo.2013.10.016 (2014).

1013 119 Chauvet, F., Sapin, F., Geoffroy, L., Ringenbach, J. C. & Ferry, J. N. Conjugate volcanic passive
1014 margins in the austral segment of the South Atlantic - Architecture and development. *Earth-Sci Rev*
1015 **212**, doi:10.1016/j.earscirev.2020.103461 (2021).

1016 120 Fromm, T. *et al.* South Atlantic opening: A plume-induced breakup? *Geology* **43**, 931-934,
1017 doi:10.1130/g36936.1 (2015).

1018 121 Planert, L. *et al.* The wide-angle seismic image of a complex rifted margin, offshore North Namibia:
1019 Implications for the tectonics of continental breakup. *Tectonophysics* **716**, 130-148,
1020 doi:10.1016/j.tecto.2016.06.024 (2017).

1021 122 Á Horni, J. *et al.* Regional distribution of volcanism within the North Atlantic Igneous Province.
1022 *Geological Society London, Special Publications* **447**, 105-125, doi:10.1144/sp447.18 (2017).

1023 123 Steinberger, B., Bredow, E., Lebedev, S., Schaeffer, A. & Torsvik, T. H. Widespread volcanism in the
1024 Greenland-North Atlantic region explained by the Iceland plume. *Nat Geosci* **12**,
1025 doi:10.1038/s41561-018-0251-0 (2019).

1026 124 Morgan, J. P., Taramon, J. M., Araujo, M., Hasenclever, J. & Perez-Gussinye, M. Causes and
1027 consequences of asymmetric lateral plume flow during South Atlantic rifting. *Proceedings of the*
1028 *National Academy of Sciences of the United States of America* **117**, 27877-27883,
1029 doi:10.1073/pnas.2012246117 (2020).

1030 125 Nielsen, T. K., Larsen, H. C. & Hopper, J. R. Contrasting rifted margin styles south of Greenland:
1031 implications for mantle plume dynamics. *Earth and Planetary Science Letters* **200**, 271-286,
1032 doi:10.1016/S0012-821x(02)00616-7 (2002).

1033 126 Moulin, M. *et al.* Geological constraints on the evolution of the Angolan margin based on reflection
1034 and refraction seismic data (ZaiAngo project). *Geophys J Int* **162**, 793-810, doi:10.1111/j.1365-
1035 246X.2005.02668.x (2005).

1036 127 Deng, H. *et al.* South China Sea documents the transition from wide continental rift to continental
1037 break up. *Nature communications* **11**, 4583, doi:10.1038/s41467-020-18448-y (2020).

1038 128 Ding, W. *et al.* Lateral evolution of the rift-to-drift transition in the South China Sea: Evidence from
1039 multi-channel seismic data and IODP Expeditions 367&368 drilling results. *Earth and Planetary*
1040 *Science Letters* **531**, 115932, doi:10.1016/j.epsl.2019.115932 (2020).

1041 129 li, F. *et al.* Continental interior and edge breakup at convergent margins induced by subduction
1042 direction reversal: A numerical modeling study applied to the South China Sea margin. *Tectonics* **39**,
1043 doi:10.1029/2020TC006409 (2020).

1044 130 li, F., Sun, Z. & Yang, H. Possible spatial distribution of the Mesozoic volcanic arc in the present-day
1045 South China Sea continental margin and its tectonic implications. *J Geophys Res-Sol Ea*,
1046 doi:10.1029/2017JB014861 (2018).

1047 131 Sun, Z. *et al.* The rifting-breakup process of the passive continental margin and its relationship with
1048 magmatism: The attribution of the South China Sea. *Earth Science* **46**, 770-789,
1049 doi:10.3799/dqkx.2020.371 (2021).

1050 132 Wan, K., Xia, S., Cao, J., Sun, J. & Xu, H.-I. Deep seismic structure of the northeastern South China
1051 Sea: Origin of a high-velocity layer in the lower crust. *J Geophys Res-Sol Ea* **122**, 2831 - 2858 (2017).

1052 133 Yang, L. *et al.* The structure and evolution of deepwater basins in the distal margin of the northern
1053 South China Sea and their implications for the formation of the continental margin. *Marine and*
1054 *Petroleum Geology* **92**, 234-254, doi:10.1016/j.marpetgeo.2018.02.032 (2018).
1055 134 Zhang, C. *et al.* Syn-rift magmatic characteristics and evolution at a sediment-rich margin: Insights
1056 from high-resolution seismic data from the South China Sea. *Gondwana Research* **91**, 81-96,
1057 doi:10.1016/j.gr.2020.11.012 (2021).
1058 135 Pin, Y., Di, Z. & Zhaoshu, L. A crustal structure profile across the northern continental margin of the
1059 South China Sea. *Tectonophysics* **338**, 1-21, doi:10.1016/S0040-1951(01)00062-2 (2001).
1060 136 Sun, Z. *et al.* The role of magmatism in the thinning and breakup of the South China Sea continental
1061 margin: Special Topic: The South China Sea Ocean Drilling. *National Science Review* **6**, 871-876,
1062 doi:10.1093/nsr/nwz116 (2019).
1063 137 Wu, J., Liu, Z. & Yu, X. Plagioclase-regulated hydrothermal alteration of basaltic rocks with
1064 implications for the South China Sea rifting. *Chem Geol* **585**, 120569,
1065 doi:10.1016/j.chemgeo.2021.120569 (2021).
1066 138 Nirrengarten, M. *et al.* Extension modes and breakup processes of the southeast China-Northwest
1067 Palawan conjugate rifted margins. *Marine and Petroleum Geology* **113**, 104123,
1068 doi:10.1016/j.marpetgeo.2019.104123 (2020).
1069 139 Ding, W., Sun, Z., Dadd, K., Fang, Y. & Li, J. Structures within the oceanic crust of the central South
1070 China Sea basin and their implications for oceanic accretionary processes. *Earth and Planetary*
1071 *Science Letters* **488**, 115-125, doi:10.1016/j.epsl.2018.02.011 (2018).
1072 140 Yu, J. *et al.* Oceanic crustal structures and temporal variations of magmatic budget during seafloor
1073 spreading in the East Sub-basin of the South China Sea. *Mar Geol* **436**, 106475,
1074 doi:10.1016/j.margeo.2021.106475 (2021).
1075 141 Huismans, R. & Beaumont, C. Depth-dependent extension, two-stage breakup and cratonic
1076 underplating at rifted margins. *Nature* **473**, 74-78, doi:10.1038/nature09988 (2011).
1077 142 Buck, W. R. Flexural rotation of normal faults. *Tectonics* **7**, 959-973, doi:10.1029/TC007i005p00959
1078 (1988).
1079 143 Pérez-Gussinyé, M. *et al.* Lithospheric strength and rift migration controls on synrift stratigraphy
1080 and breakup unconformities at rifted margins: Examples from numerical models, the Atlantic and
1081 South China Sea margins. *Tectonics* **39**, doi:10.1029/2020TC006255 (2020).
1082 144 Theunissen, T. & Huismans, R. S. Long-term coupling and feedback between tectonics and surface
1083 processes during non-volcanic rifted margin formation. *J Geophys Res-Sol Ea* **124**, 12323-12347,
1084 doi:10.1029/2018JB017235 (2019).
1085 145 Ros, E. *et al.* Lower crustal strength controls on melting and serpentinization at magma-poor
1086 margins: Potential implications for the South Atlantic. *Geochemistry, Geophysics, Geosystems* **18**,
1087 4538-4557, doi:10.1002/2017GC007212 (2017).
1088 146 Armitage, J., Petersen, K. D. & Perez-Gussinye, M. The role of crustal strength in controlling
1089 magmatism and melt chemistry during rifting and break-up. *Geochemistry, Geophysics, Geosystems*
1090 **19**, doi:10.1002/2017gc007326 (2018).
1091 147 Lu, G. & Huismans, R. S. Melt volume at Atlantic volcanic rifted margins controlled by depth-
1092 dependent extension and mantle temperature. *Nature communications* **12**, 3894,
1093 doi:10.1038/s41467-021-23981-5 (2021).
1094 148 Cowie, P. A., Underhill, J. R., Behn, M. D., Lin, J. & Gill, C. E. Spatio-temporal evolution of strain
1095 accumulation derived from multi-scale observations of Late Jurassic rifting in the northern North
1096 Sea: A critical test of models for lithospheric extension. *Earth and Planetary Science Letters* **234**,
1097 401-419, doi:10.1016/j.epsl.2005.01.039 (2005).
1098 149 Mattei, M. *et al.* Tectonic evolution of fault-bounded continental blocks: Comparison of
1099 paleomagnetic and GPS data in the Corinth and Megara basins (Greece). *J Geophys Res-Sol Ea* **109**,
1100 doi:10.1029/2003JB002506 (2004).

1101 150 Gawthorpe, R. L. *et al.* Normal fault growth, displacement localisation and the evolution of normal
1102 fault populations: the Hammam Faraun fault block, Suez rift, Egypt. *Journal of Structural Geology*
1103 **25**, 883-895, doi:10.1016/S0191-8141(02)00088-3 (2003).
1104 151 Brune, S., Williams, S. E., Butterworth, N. P. & Muller, R. D. Abrupt plate accelerations shape rifted
1105 continental margins. *Nature* **536**, 201-204, doi:10.1038/nature18319 (2016).
1106 152 Petit, C. & Déverchère, J. Structure and evolution of the Baikal rift: A synthesis. *Geochemistry,*
1107 *Geophysics, Geosystems* **7**, doi:10.1029/2006GC001265 (2006).
1108 153 England, P. Constraints on extension of continental lithosphere. *J Geophys Res-Sol Ea* **88**, 1145-
1109 1152, doi:10.1029/JB088iB02p01145 (1983).
1110 154 Pérez-Gussinyé, M. & Reston, T. J. Rheological evolution during extension at nonvolcanic rifted
1111 margins: Onset of serpentinization and development of detachments leading to continental
1112 breakup. *J Geophys Res-Sol Ea* **106**, 3961-3975, doi:10.1029/2000jb900325 (2001).
1113 155 Brune, S., Heine, C., Perez-Gussinye, M. & Sobolev, S. V. Rift migration explains continental margin
1114 asymmetry and crustal hyper-extension. *Nature communications* **5**, 4014, doi:10.1038/ncomms5014
1115 (2014).
1116 156 Svartman Dias, A. E., Lavier, L. L. & Hayman, N. W. Conjugate rifted margins width and asymmetry:
1117 The interplay between lithospheric strength and thermomechanical processes. *J Geophys Res-Sol Ea*
1118 **120**, 8672-8700, doi:10.1002/2015JB012074 (2015).
1119 157 Tetreault, J. L. & Buiter, S. J. H. The influence of extension rate and crustal rheology on the evolution
1120 of passive margins from rifting to break-up. *Tectonophysics* **746**, 155-172,
1121 doi:10.1016/j.tecto.2017.08.029 (2018).
1122 158 Naliboff, J. & Buiter, S. J. H. Rift reactivation and migration during multiphase extension. *Earth and*
1123 *Planetary Science Letters* **421**, 58-67, doi:10.1016/j.epsl.2015.03.050 (2015).
1124 159 Neuharth, D. *et al.* Evolution of rift systems and their fault networks in response to surface
1125 processes. *Tectonics* **41**, doi:10.1029/2021TC007166 (2022).
1126 160 Reston, T. J. Polyphase faulting during the development of the west Galicia rifted margin. *Earth and*
1127 *Planetary Science Letters* **237**, 561-576, doi:10.1016/j.epsl.2005.06.019 (2005).
1128 161 Naliboff, J. B., Buiter, S. J. H., Peron-Pinvidic, G., Osmundsen, P. T. & Tetreault, J. Complex fault
1129 interaction controls continental rifting. *Nature communications* **8**, 1179, doi:10.1038/s41467-017-
1130 00904-x (2017).
1131 162 Lavier, L. L. & Manatschal, G. A mechanism to thin the continental lithosphere at magma-poor
1132 margins. *Nature* **440**, 324-328, doi:10.1038/nature04608 (2006).
1133 163 Lavier, L. L. & Buck, W. R. Half graben versus large-offset low-angle normal fault: Importance of
1134 keeping cool during normal faulting. *J Geophys Res-Sol Ea* **107**, doi:10.1029/2001JB000513 (2002).
1135 164 Rey, P. F., Teyssier, C. & Whitney, D. L. Extension rates, crustal melting, and core complex dynamics.
1136 *Geology* **37**, 391-394, doi:10.1130/g25460a.1 (2009).
1137 165 Brun, J. P., White, R. S., Hardman, R. F. P., Watts, A. B. & Whitmarsh, R. B. Narrow rifts versus wide
1138 rifts: inferences for the mechanics of rifting from laboratory experiments. *Philosophical Transactions*
1139 *of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences* **357**, 695-
1140 712, doi:10.1098/rsta.1999.0349 (1999).
1141 166 Buiter, S. J. H., Huismans, R. S. & Beaumont, C. Dissipation analysis as a guide to mode selection
1142 during crustal extension and implications for the styles of sedimentary basins. *J Geophys Res-Sol Ea*
1143 **113**, doi:10.1029/2007JB005272 (2008).
1144 167 Olive, J.-A., Behn, M. D. & Malatesta, L. C. Modes of extensional faulting controlled by surface
1145 processes. *Geophys Res Lett* **41**, 6725-6733, doi:10.1002/2014GL061507 (2014).
1146 168 Andres-Martinez, M., Perez-Gussinye, M., Armitage, J. & Morgan, J. P. Thermomechanical
1147 implications of sediment transport for the architecture and evolution of continental rifts and
1148 margins. *Tectonics* **38**, 641-665, doi:10.1029/2018tc005346 (2019).

1149 169 Peron-Pinvidic, G., Manatschal, G., Minshull, T. A. & Sawyer, D. S. Tectonosedimentary evolution of
1150 the deep Iberia-Newfoundland margins: Evidence for a complex breakup history. *Tectonics* **26**,
1151 doi:10.1029/2006tc001970 (2007).

1152 170 Masini, E., Manatschal, G. & Mohn, G. The Alpine Tethys rifted margins: Reconciling old and new
1153 ideas to understand the stratigraphic architecture of magma-poor rifted margins. *Sedimentology* **60**,
1154 174-196, doi:10.1111/sed.12017 (2013).

1155 171 Pérez-Gussinyé, M. *et al.* A tectonic model for hyperextension at magma-poor rifted margins: an
1156 example from the West Iberia–Newfoundland conjugate margins. *Geological Society London, Special*
1157 *Publications* **369**, doi:10.1144/sp369.19 (2013).

1158 172 Karner, G. D., Gambôa, L. A. P., Schreiber, B. C., Lugli, S. & Båbel, M. Timing and origin of the South
1159 Atlantic pre-salt sag basins and their capping evaporites. *Geological Society London, Special*
1160 *Publications* **285**, doi:10.1144/sp285.2 (2007).

1161 173 Neto Araujo, M., Pérez-Gussinyé, M. & Muldashev, I. Oceanward rift migration during formation of
1162 Santos–Benguela ultra-wide rifted margins. *Geological Society London, Special Publications* **524**,
1163 doi:10.1144/SP524-2021-123 (2022).

1164 174 Bastow, I. D. & Keir, D. The protracted development of the continent-ocean transition in Afar. *Nat*
1165 *Geosci* **4**, 248-250, doi:10.1038/Ngeo1095 (2011).

1166 175 Buck, W. R. The role of magma in the development of the Afro-Arabian Rift System. *Geological*
1167 *Society London, Special Publication* **259**, 43-54, doi:10.1144/gsl.sp.2006.259.01.05 (2006).

1168 176 Annen, C. & Sparks, R. S. J. Effects of repetitive emplacement of basaltic intrusions on thermal
1169 evolution and melt generation in the crust. *Earth and Planetary Science Letters* **203**, 937-955,
1170 doi:10.1016/S0012-821x(02)00929-9 (2002).

1171 177 Armitage, J. J. & Collier, J. S. The thermal structure of volcanic passive margins. *Petrol Geosci* **24**,
1172 393-401, doi:10.1144/petgeo2016-101 (2018).

1173 178 Jull, M. & McKenzie, D. The effect of deglaciation on mantle melting beneath Iceland. *J Geophys*
1174 *Res-Sol Ea* **101**, 21815-21828, doi:10.1029/96JB01308 (1996).

1175 179 Armitage, J. J., Ferguson, D. J., Petersen, K. D. & Creyts, T. T. The importance of Icelandic ice sheet
1176 growth and retreat on mantle CO₂ flux. *Geophys Res Lett* **46**, 6451-6458,
1177 doi:10.1029/2019GL081955 (2019).

1178 180 Sternai, P. Surface processes forcing on extensional rock melting. *Scientific Reports* **10**, 7711,
1179 doi:10.1038/s41598-020-63920-w (2020).

1180 181 Theissen-Krah, S., Iyer, K., Rüpke, L. H. & Morgan, J. P. Coupled mechanical and hydrothermal
1181 modeling of crustal accretion at intermediate to fast spreading ridges. *Earth and Planetary Science*
1182 *Letters* **311**, 275-286, doi:10.1016/j.epsl.2011.09.018 (2011).

1183 182 Sibuet, J. C., Srivastava, S. & Manatschal, G. Exhumed mantle-forming transitional crust in the
1184 Newfoundland-Iberia rift and associated magnetic anomalies. *J Geophys Res-Sol Ea* **112**,
1185 doi:10.1029/2005jb003856 (2007).

1186 183 Reston, T. J. & Morgan, J. P. Continental geotherm and the evolution of rifted margins. *Geology* **32**,
1187 133-136, doi:10.1130/G19999.1 (2004).

1188 184 Müntener, O. & Manatschal, G. High degrees of melt extraction recorded by spinel harzburgite of
1189 the Newfoundland margin: The role of inheritance and consequences for the evolution of the
1190 southern North Atlantic. *Earth and Planetary Science Letters* **252**, 437-452,
1191 doi:10.1016/j.epsl.2006.10.009 (2006).

1192 185 Pérez-Gussinyé, M., Morgan, J. P., Reston, T. J. & Ranero, C. R. The rift to drift transition at non-
1193 volcanic margins: Insights from numerical modelling. *Earth and Planetary Science Letters* **244**, 458-
1194 473, doi:10.1016/j.epsl.2006.01.059 (2006).

1195 186 Mohn, G. *et al.* Structural and stratigraphic evolution of the Iberia–Newfoundland hyper-extended
1196 rifted margin: a quantitative modelling approach. *Geological Society London, Special Publications*
1197 **413**, doi:10.1144/sp413.9 (2015).

1198 187 Reston, T. J. & McDermott, K. G. Successive detachment faults and mantle unroofing at magma-
1199 poor rifted margins. *Geology* **39**, 1071-1074, doi:10.1130/G32428.1 (2011).
1200 188 Gillard, M. *et al.* Birth of an oceanic spreading center at a magma-poor rift system. *Scientific Reports*
1201 **7**, 15072, doi:10.1038/s41598-017-15522-2 (2017).
1202 189 Brune, S., Heine, C., Clift, P. D. & Perez-Gussinye, M. Rifted margin architecture and crustal
1203 rheology: Reviewing Iberia-Newfoundland, Central South Atlantic, and South China Sea. *Marine and*
1204 *Petroleum Geology* **79**, 257-281, doi:10.1016/j.marpetgeo.2016.10.018 (2017).
1205 190 van Wijk, J. W., Huisman, R. S., ter Voorde, M. & Cloetingh, S. A. P. L. Melt generation at volcanic
1206 continental margins: no need for a mantle plume? *Geophys Res Lett* **28**, 3995-3998 (2001).
1207 191 Campbell, I. H. Testing the plume theory. *Chem Geol* **241**, 153-176,
1208 doi:10.1016/j.chemgeo.2007.01.024 (2007).
1209 192 Sleep, N. H. Lateral flow and ponding of starting plume material. *J Geophys Res-Sol Ea* **102**, 10001-
1210 10012, doi:10.1029/97jb00551 (1997).
1211 193 Ringrose, P. S. & Meckel, T. A. Maturing global CO₂ storage resources on offshore continental
1212 margins to achieve 2DS emissions reductions. *Scientific Reports* **9**, 17944, doi:10.1038/s41598-019-
1213 54363-z (2019).
1214 194 Krevor, S. Subsurface carbon dioxide and hydrogen storage in a sustainable energy future. *Nature*
1215 *Reviews Earth & Environment* (in press).
1216 195 Goldberg, D. S., Kent, D. V. & Olsen, P. E. Potential on-shore and off-shore reservoirs for CO₂
1217 sequestration in Central Atlantic magmatic province basalts. *Proceedings of the National Academy*
1218 *of Sciences* **107**, 1327-1332, doi:10.1073/pnas.0913721107 (2010).
1219 196 Snæbjörnsdóttir, S. Ó. *et al.* Carbon dioxide storage through mineral carbonation. *Nature Reviews*
1220 *Earth & Environment* **1**, 90-102, doi:10.1038/s43017-019-0011-8 (2020).
1221 197 <https://www.lithiumdefrance.earth/copie-de-les-sources>.
1222 198 Arnulf, A. F., Harding, A. J., Singh, S. C., Kent, G. M. & Crawford, W. Fine-scale velocity structure of
1223 upper oceanic crust from full waveform inversion of downward continued seismic reflection data at
1224 the Lucky Strike Volcano, Mid-Atlantic Ridge. *Geophys Res Lett* **39**, doi:10.1029/2012GL051064
1225 (2012).
1226 199 Peron-Pinvidic, G., Fourel, L. & Buiter, S. J. H. The influence of orogenic collision inheritance on
1227 rifted margin architecture: Insights from comparing numerical experiments to the Mid-Norwegian
1228 margin. *Tectonophysics* **828**, doi:10.1016/j.tecto.2022.229273 (2022).
1229 200 Neuharth, D., Brune, S., Glerum, A., Heine, C. & Welford, J. K. Formation of continental microplates
1230 through rift linkage: Numerical modeling and its application to the Flemish Cap and Sao Paulo
1231 Plateau. *Geochemistry, Geophysics, Geosystems* **22**, doi:10.1029/2020GC009615 (2021).
1232 201 Gouiza, M. & Naliboff, J. Rheological inheritance controls the formation of segmented rifted margins
1233 in cratonic lithosphere. *Nature communications* **12**, 4653, doi:10.1038/s41467-021-24945-5 (2021).
1234 202 Le Pourhiet, L. *et al.* Continental break-up of the South China Sea stalled by far-field compression.
1235 *Nat Geosci* **11**, 605-609, doi:10.1038/s41561-018-0178-5 (2018).
1236 203 Seton, M. *et al.* A global data set of present-day oceanic crustal age and seafloor spreading
1237 parameters. *Geochemistry, Geophysics, Geosystems* **21**, e2020GC009214,
1238 doi:10.1029/2020GC009214 (2020).
1239 204 Bryan, S. E. & Ferrari, L. Large igneous provinces and silicic large igneous provinces: Progress in our
1240 understanding over the last 25 years. *GSA Bulletin* **125**, 1053-1078, doi:10.1130/b30820.1 (2013).
1241 205 Biari, Y. *et al.* Structure and evolution of the Atlantic passive margins: A review of existing rifting
1242 models from wide-angle seismic data and kinematic reconstruction. *Marine and Petroleum Geology*
1243 **126**, 104898, doi:10.1016/j.marpetgeo.2021.104898 (2021).
1244 206 Eddy, D. R., Van Avendonk, H. J. A., Christeson, G. L. & Norton, I. O. Structure and origin of the rifted
1245 margin of the northern Gulf of Mexico. *Geosphere* **14**, 1804-1817, doi:10.1130/ges01662.1 (2018).

- 1246 207 Eddy, D. R. *et al.* Deep crustal structure of the northeastern Gulf of Mexico: Implications for rift
1247 evolution and seafloor spreading. *J Geophys Res-Sol Ea* **119**, 6802-6822, doi:10.1002/2014JB011311
1248 (2014).
- 1249 208 Direen, N. G., Stagg, H. M. J., Symonds, P. A. & Colwell, J. B. Dominant symmetry of a conjugate
1250 southern Australian and East Antarctic magma-poor rifted margin segment. *Geochemistry,*
1251 *Geophysics, Geosystems* **12**, doi:10.1029/2010GC003306 (2011).
- 1252 209 Ball, P., Eagles, G., Ebinger, C., McClay, K. & Totterdell, J. The spatial and temporal evolution of
1253 strain during the separation of Australia and Antarctica. *Geochemistry, Geophysics, Geosystems* **14**,
1254 2771-2799, doi:10.1002/ggge.20160 (2013).
- 1255 210 Gillard, M. *et al.* Tectonomagmatic evolution of the final stages of rifting along the deep conjugate
1256 Australian-Antarctic magma-poor rifted margins: Constraints from seismic observations. *Tectonics*
1257 **34**, 753-783, doi:10.1002/2015tc003850 (2015).
- 1258 211 Muldashev, I. A., Pérez-Gussinyé, M. & de Araújo, M. N. C. KineDyn: Thermomechanical forward
1259 method for validation of seismic interpretations and investigation of dynamics of rifts and rifted
1260 margins. *Phys Earth Planet In* **317**, 106748, doi:10.1016/j.pepi.2021.106748 (2021).
- 1261 212 Deregowski, S. M. Prestack depth migration by the 2-D boundary integral method. *4th Annual*
1262 *International Meeting, SEG, Expanded Abstracts*, 414–417 (1985).
- 1263 213 Kelemen, P. B. & Holbrook, W. S. Origin of thick, high-velocity igneous crust along the US east-coast
1264 margin. *J Geophys Res-Sol Ea* **100**, 10077-10094, doi:10.1029/95jb00924 (1995).
- 1265 214 Korenaga, J., Kelemen, P. B. & Holbrook, W. S. Methods for resolving the origin of large igneous
1266 provinces from crustal seismology. *J Geophys Res-Sol Ea* **107**, doi:10.1029/2001jb001030 (2002).
- 1267 215 Davy, R. G., Collier, J. S., Henstock, T. J. & Consortium, T. V. Wide-angle seismic imaging of two
1268 modes of crustal accretion in mature Atlantic Ocean crust. *J Geophys Res-Sol Ea* **125**,
1269 doi:10.1029/2019JB019100 (2020).
- 1270 216 Dunn, R. A., Arai, R., Eason, D. E., Canales, J. P. & Sohn, R. A. Three-dimensional seismic structure of
1271 the Mid-Atlantic Ridge: An investigation of tectonic, magmatic, and hydrothermal processes in the
1272 Rainbow Area. *J Geophys Res-Sol Ea* **122**, 9580-9602, doi:10.1002/2017JB015051 (2017).
- 1273 217 Corbalán, A. *et al.* Seismic velocity structure along and across the ultraslow-spreading Southwest
1274 Indian Ridge at 64° 30' E showcases flipping detachment faults. *J Geophys Res-Sol Ea* **126**,
1275 doi:10.1029/2021JB022177 (2021).
- 1276 218 Sauter, D. *et al.* Continuous exhumation of mantle-derived rocks at the Southwest Indian Ridge for
1277 11 million years. *Nat Geosci* **6**, 314-320, doi:10.1038/ngeo1771 (2013).

1278