

Ported alias Set-Pointed and Non-scalar Tutte Functions

Seth Chaiken

Computer Science Department

University at Albany, State Univ. of New York, USA

`sdc@cs.albany.edu`

`http://www.cs.albany.edu/~sdc/Matroids`

October 9, 2005

Traditionally a **Tutte Function** F

$$F : \begin{cases} \text{Matroids} \\ \text{(or Graphs)} \end{cases} \rightarrow \text{Commutative Ring } (R, +, \cdot)$$

$$F(\mathcal{N}) = g_e F(\mathcal{N}/e) + r_e F(\mathcal{N} \setminus e) \tag{A}$$

for all e not a loop or inthmus. g_e, r_e are parameters or 1.

$$F(\mathcal{N}_1 \oplus \mathcal{N}_2) = F(\mathcal{N}_1) \cdot F(\mathcal{N}_2) \tag{M}$$

We survey results where

- (1) $(R, +, \cdot)$ is replaced by discrete or other algebraic structures (Matroids!), or “Matroids” is replaced by matroid presentations;
- and** (2) given a distinguished subset P (**ports** or **set of points**), (A) is restricted to $e \notin P$.

1. $F : \text{Matroids} \rightarrow \text{Matroids}$.

2. $F : \text{Extensors} \rightarrow \text{Extensors}$.

The multiplication of exterior (aka Cayley, Grassmann) algebra is anticommutative: $\mathbf{N}_1 \mathbf{N}_2 = (\mathbf{N}_1 \wedge \mathbf{N}_2) = \mathbf{N}_2 \mathbf{N}_1 (-1)^{r(\mathbf{N}_1)r(\mathbf{N}_2)}$

3. $F : (\text{Oriented Matroids} \rightarrow \text{Commutative Ring where a substitution expresses (2) in the unimodular (regular) oriented case.}$

4. Algebraic expressions of (3) for P -unions and generalized parallel connections over P .

In all cases, “ports”, “set of points” P , for which deletion/contraction is forbidden, must satisfy $P \neq \emptyset$ for interesting results.

Result 1: (Construction of Reeski, Weinberg 70's; new observation here)

Given: Matroid $\mathcal{N}(P, E)$ has ground set $P \cup E$.

P_V and P_I are two disjoint copies of P :

$$P_V = \{p_V : p \in P\}; P_I = \{p_I : p \in P\}.$$

(The matroid!) $\mathcal{M}_E(\mathcal{N})(P_I \cup P_V) := (\mathcal{N}(P_I, E)) \cup \mathcal{N}^\perp(P_V, E)) / E$
satisfies

$$\mathcal{M}_E(\mathcal{N}) = \mathcal{M}_{E'}(\mathcal{N}/e) \cup_{\mathcal{B}} \mathcal{M}_{E'}(\mathcal{N} \setminus e) \text{ if } e \notin P \text{ and } E' = E \setminus e,$$

where $\cup_{\mathcal{B}}$ denotes union of matroid basis collections; and

$$\mathcal{M}_E(\mathcal{N}_1(E_1, P_1) \oplus \mathcal{N}_2(E_2, P_2)) = \mathcal{M}_{E_1}(\mathcal{N}_1(E_1, P_1)) \oplus \mathcal{M}_{E_2}(\mathcal{N}_2(E_2, P_2)).$$

where \oplus denotes matroid direct sum.

Proof: E is independent in $\mathcal{N}(P_I, E) \cup \mathcal{N}^\perp(P_V, E)$, so B is a basis of $\mathcal{M}_E(\mathcal{N})$ iff $A \subseteq E$, $B \cup A \cup (E \setminus A)$ is a basis of the union; thence either $e \in A$ or $e \in (E \setminus A)$.

Result 2. applies to a given **decomposable** $(\mathbf{N}(P, E) = \mathbf{v}_1 \mathbf{v}_2 \cdots \mathbf{v}_r$ of vectors) in the **exterior algebra** \mathcal{E} over vector space $k(P \cup E)$ with distinguished basis $(P \cup E)$.

We say an **extensor** is a decomposable element in \mathcal{E} .

Let $P \cup E$ label the columns of matrix N (full row rank r).

The (oriented) matroid of column dependencies (rank r) is determined by the *row space* of $N \leftrightarrow k$ -multiples of one extensor \mathbf{N} , such as $\mathbf{N} = \wedge(\text{row vectors of } N) \in \mathcal{E}$.

\mathcal{E} is quotient of the assoc. algebra generated by $k(P \cup E)$ modulo the ideal generated by \mathbf{v}^2 , $\mathbf{v} \in k(P \cup E)$. \mathcal{E} has dimension $2^{|P \cup E|} = 2^n$.

\mathcal{E} is graded: At rank r , \mathcal{E}_r has dim. $\binom{n}{r}$, $r = 0, 1, \dots, n$.

Natural coordinates of extensor $\mathbf{N}(P, E)$ are the $r \times r$ *determinants* in matrix N .

Exterior *sum* can be expressed by addition of the expansion coefficients under the basis of all 2^n subsets of $P \cup E$, (each subset with an element order fixed).

But, the sum of two extensors is not necessarily *decomposable*, not necessarily an *extensor*.

The exterior *product* of extensors for disjoint subspaces represents the **subspace join**.

When a ground set S is given (like $S = E \cup P$, distinguished basis for $k(E \cup P)$), **deletion** and **contraction** of $e \in S$, and **dualization**, known from multilinear algebra, represent the corresponding (oriented) matroid operations.

We define deletion/contraction so

$$\mathbf{N} = \mathbf{N} \setminus e + (\mathbf{N}/e)\mathbf{e}$$

Dualization: Copy the oriented matroid chirotope dualization formula (also known as Hodge star): Coefficients (determinants!) $\mathbf{N}^\perp[X] = \mathbf{N}[\overline{X}]\epsilon(\overline{X}, X)$.

Our Tutte-like function $\mathbf{M}_E(\mathbf{N}) : \text{Extensors} \rightarrow \text{Extensors}$.

Given N (matrix), construct N^\perp so their row spaces are orthogonal complements (N^\perp presents the (oriented) dual of the matroid from N).

Form the matrix: ($G = \text{diag}(g_e)$, $R = \text{diag}(r_e)$)

$$M = \left[\begin{array}{c|c|c} N(P) & 0 & N(E)G \\ \hline 0 & N^\perp(P) & N^\perp(E)R \end{array} \right]$$

with columns labelled by $P_I \cup P_V \cup E$.

Extensor \mathbf{M} over $k[g_e, r_e](P_V \cup P_I \cup E)$ is the product of M 's **row vectors**, and define $\mathbf{M}_E(\mathbf{N})$ by:

$$\mathbf{M} = \mathbf{M}_E(\mathbf{N})\mathbf{e}_1\mathbf{e}_2 \cdots \mathbf{e}_{|E|} + (\cdots)$$

Result 2: (2003)

$$\epsilon(P E) \; \mathbf{M}_E(\mathbf{N}(P, E)) =$$

$$\epsilon(P E') \; (g_e \mathbf{M}_{E'}(\mathbf{N}/e) + r_e \mathbf{M}_{E'}(\mathbf{N} \setminus e))$$

$$\epsilon(P_1 P_2 E) \; \mathbf{M}_E(\; \mathbf{N}_1(P_1, E_1) \; \mathbf{N}_2(P_2, E_2) \;) =$$

$$\epsilon(P_1 E_1) \epsilon(P_2 E_2) \; \mathbf{M}_{E_1}(\mathbf{N}_1(P_1, E_1)) \; \mathbf{M}_{E_2}(N_2(P_2, E_2))$$

Corollory:

$$\epsilon(P E) \mathbf{M}_E(\mathbf{N}) = \epsilon(P) \qquad \sum \qquad \mathbf{M}_{\emptyset}(\mathbf{N}/A|P) g_A r_A^{\rightarrow} . \quad (1)$$

$$A \subseteq E : \mathbf{r}_{\mathbf{N}} A = |A|,$$

$$\mathbf{r} \mathbf{N} - \mathbf{r}(\mathbf{N}/A|P) - \mathbf{r}_{\mathbf{N}} A = 0$$

Compare: (with $u = 0$ and $v = 0$)

$$R(\mathcal{N}(P, E)) = \sum_{A \subseteq E} [\mathcal{N}/A|P] \; g_A r_A^{\rightarrow} u^{\mathbf{N} - \mathbf{r}[\mathcal{N}/A|P] - \mathbf{r}_A} g_v^A |A| - \mathbf{r}_A .$$

Result 3: The ported, parametrized corank-nullity polynomials of oriented and non-oriented matroids

$$R_P(\mathcal{N}(P, E)) = \sum_{A \subseteq E} [\mathcal{N}/A|P] \, g_A r_A^{-u} r^{\mathcal{N}-r[\mathcal{N}/A|P]-r_A} g_{|A|-r_A}.$$

satisfy the ported Tutte equations.

(In the invariant case, this R is universal for Tutte P -invariants of oriented and non-oriented matroids)

$[\mathcal{N}/A|P]$ is a (commutative) *monomial* whose factors *are connected (oriented) matroids* over subsets of P .

(Las Vergnas “Big Tutte Polynomial” (’75, ’99), oriented/parametrized by sdc.)

R_P can distinguish some orientations of the same matroid when $|P| \geq 2$.

Result 4 pertains to three matroid combination operations:

Given $\mathcal{N}_1(P, E_1)$ and $\mathcal{N}_2(P, E_2)$ with only elements P in common.

(1.) Ported matroid union: $\mathcal{B}(\mathcal{N}_1 \cup \mathcal{N}_2) = \{B_1 \cup B_2 : B_i \in \mathcal{B}(\mathcal{N}_i)\}$

(2.) Duality conjugate \cup^* of \cup .

(3.) Given that P is a modular flat and a common submatroid in \mathcal{N}_1 and \mathcal{N}_2 , the generalized parallel connection.

When $|P| = 1$, both dual \cup and generalized parallel connection REDUCE to (one base point) parallel connection.

K =polynomial ring containing u and v . K_P =commutative K -module generated by monomials $[q_i]$ signifying matroids q_i over subsets of P , and $[\emptyset] = 1$.

Recall ported corank-nullity polynomial $R_P : \text{Matroids} \rightarrow K_P$.

Result 4: For each combination operation $(*_i, i = 1, 2, 3)$ there is a bilinear $M_i : K_P \times K_P \rightarrow K_P$ map and an $u^j v^k$ -valued function f_i such that

$$R_P(\mathcal{N}_1 *_i \mathcal{N}_2) = f_i(\mathcal{N}_1, \mathcal{N}_2, \mathcal{N}_1 *_i \mathcal{N}_2) M_i(R_P(\mathcal{N}_1), R_P(\mathcal{N}_2))$$

In other words, once $*_i$ is determined on pairs of matroids over subsets of P , the ported R_P for $\mathcal{N}_1 *_i \mathcal{N}_2$ can be calculated by formal multiplication

$$f_i(..) R_P(\mathcal{N}_1) *_i R_P(\mathcal{N}_2) = f_i(..) M_i(R_P(\mathcal{N}_1), R_P(\mathcal{N}_2))$$

and substitutions of $(1/f_i(q_a, q_b, q_a *_i q_b)) [q_a *_i q_b] \leftarrow [q_a] *_i [q_b]$.

Splitting Formulas generalize $F(\mathcal{N}_1 \oplus \mathcal{N}_2) = F(\mathcal{N}_1)F(\mathcal{N}_2)$ and solve this type of problem:

Given $\mathcal{N}_1(P \cup E_1)$, $\mathcal{N}_2(P \cup E_2)$, can we calculate the Tutte polynomial of $\mathcal{N}_1 *_i \mathcal{N}_2(P \cup E_1 \cup E_2)$ from Tutte polynomials of minors of \mathcal{N}_1 and \mathcal{N}_2 gotten by deletion/contraction of subsets of P ?

For one-point series and parallel connections, we can re-derive Brylawski's (1971) formulas by the above bilinear method (1989).

For generalized parallel connection, with P a modular flat and common restriction in both \mathcal{N}_1 and \mathcal{N}_2 , a splitting formula was obtained by Bonin and de Mier (2004).

Their formula, for the Tutte polynomial, is in terms of:

1. The lattice of flats $F \leq P$,
2. characteristic polys. of P/F , and
3. Tutte polynomials of \mathcal{N}_1/F and \mathcal{N}_2/F .

Our formulas are (1) for R_P (Big Tutte polynomial) and (2) are in terms of $R_P(\mathcal{N}_1) *_i R_P(\mathcal{N}_2)$. Our bilinear forms' coefficients and f in $R_P(\mathcal{N}_1 *_i \mathcal{N}_2) = f_i(\dots) R_P(\mathcal{N}_1) *_i R_P(\mathcal{N}_2)$:

$$1. * = \cup \quad (1989)$$

$$f(\mathcal{N}_1, \mathcal{N}_2, \mathcal{N}_1 \cup \mathcal{N}_2) = u^{\mathbf{r}_{\mathcal{N}_1 \cup \mathcal{N}_2} - \mathbf{r}_{\mathcal{N}_1} - \mathbf{r}_{\mathcal{N}_2}}.$$

$$2. * = \cup^* \quad (1989)$$

$$f(\mathcal{N}_1, \mathcal{N}_2, \mathcal{N}_1 \cup^* \mathcal{N}_2) = v^{\mathbf{r}_{\mathcal{N}_1} + \mathbf{r}_{\mathcal{N}_2} - \mathbf{r}_{\mathcal{N}_1 \cup^* \mathcal{N}_2}}.$$

3. Generalized Parallel P -Connection $*$ (1991) Since P is a modular flat, quotients $Q_i = \mathcal{N}_i / A_i | P$ correspond to flats in P , and $[Q_i] * [Q_j] = (1/f)[Q']$ where Q' is the quotient corresponding to the join of those flats in P .

$$f(\mathcal{N}_1, \mathcal{N}_2, \mathcal{N}_1 * \mathcal{N}_2) = v^{\mathbf{r}_{\mathcal{N}_1} + \mathbf{r}_{\mathcal{N}_2} - \mathbf{r}_{\mathcal{N}_1 * \mathcal{N}_2}}.$$

Brylawski's splitting formulas for $|P| = 1$ (series/union and parallel/co-union conn.) can be derived from our $R_P(\mathcal{N}_1) * R_P(\mathcal{N}_2)$ formula.

Can Bonin and de Meir's generalized parallel connection splitting formula for all $|P|$ also be derived that way?

Are there other expressions for the splitting function?