

Supplement For  
Beauty of  
**VEDIC SPEED**  
**MATHEMATICS**

**(Journey from Limited Intelligence to Human Bio-Calculator)**

**Version – 4 (27-03-2020)**

Paperback, eBook & **Video Course** on Vedic Speed Mathematics  
[www.Speed16.com/books/vm](http://www.Speed16.com/books/vm)


**Location of this file:** [www.Speed16.com/files/vms1.pdf](http://www.Speed16.com/files/vms1.pdf)

**Note:** We keep updating this file (adding new examples, new graphical representations etc.). Keep visiting for updates.

# MULTIPLICATION

## Multiplication using Base Method

**Case 1: When both numbers (multiplicand and multiplier) are less than the working base:**




Ex.1: 7×8	Ex.2: 6×7	Ex.3: 94×96	Ex.4: 90×89	Ex.5: 997×993	Ex.6: 950×930
B:10	B:10	B:100	B:100	B:1000	B:1000
 7   -3 8   -2 ----- 5   6	6   -4 7   -3 ----- 3   12 3+1   2 4   2	94   -6 96   -4 ----- 90   24	90   -10 89   -11 ----- 79   110 79+1   10 80   10	997   -3 993   -7 ----- 990   021	950   -50 930   -70 ----- 880   3500 880+3   500 883   500
56	42	9024	8010	990021	883500

## Multiplication using Criss Cross Method

**Case 1: Two Digit Numbers (2D×2D and 2D×1D) {D: Digit}**






### Graphical Representation:

Answer consists of three parts.

First Part:	Second Part:	Third Part:
 a   b c   d	 a   b c   d	 a   b c   d
(a×c)	(a×d + b×c)	(b×d)





## Case 2: Three Digit Numbers (3D×3D; 3D×2D and 3D×1D)




Answer consists of Five Parts.

First Part:	Second Part:	Third Part:	Fourth Part:	Fifth Part:
 a   b   c d   e   f	 a   b   c d   e   f	 a   b   c d   e   f	 a   b   c d   e   f	 a   b   c d   e   f
$(a \times d)$	$(a \times e + b \times d)$	$(a \times f + b \times e + c \times d)$	$(b \times f + c \times e)$	$(c \times f)$

## CASE 3: (4×3; 4×3; 4×2; 4×1)




Answer consists of seven parts.




First Part:	Second Part:	Third Part:	Fourth Part:
 a   b   c   d e   f   g   h	 a   b   c   d e   f   g   h	 a   b   c   d e   f   g   h	 a   b   c   d e   f   g   h
$(a \times e)$	$(a \times f) + (b \times e)$	$(a \times g) + (b \times f) + (c \times e)$	$(a \times h) + (b \times g) + (c \times f) + (d \times e)$




Fifth Part:	Sixth Part:	Seventh Part:
 a   b   c   d e   f   g   h	 a   b   c   d e   f   g   h	 a   b   c   d e   f   g   h
$(b \times h) + (c \times g) + (d \times f)$	$(c \times h) + (d \times g)$	$(d \times h)$

#### CASE 4: (5×5; 5×4; 5×3; 5×2; 5×1)

Answer consists of nine parts.

First Part:	Second Part:	Third Part:
		
(a×f)	(a×g)+(b×f)	(a×h)+(b×g)+(c×f)

Fourth Part:	Fifth Part:	Sixth Part:
		
(a×i)+(b×h)+(c×g)+(d×f)	(a×j)+(b×i)+(c×h)+(d×g)+(e×f)	(b×j)+(c×i)+(d×h)+(e×g)

Seventh Part:	Eighth Part:	Ninth Part:
		
(c×j)+(d×i)+(e×h)	(d×j)+(e×i)	(e×j)

Books, eBooks, Video Course, FREE Workbook & FREE Online Training on Vedic Speed Mathematics, C & Python Programming:

[www.Speed16.com/books/vm](http://www.Speed16.com/books/vm)

**CASE 5: (6×6; 6×5; 6×4; 6×3; 6×2; 6×1) (Do it Yourself)**

Answer consists of eleven parts.

First Part:	Second Part	Third Part:	Fourth Part	Fifth Part:
a b c d e f	a b c d e f	a b c d e f	a b c d e f	a b c d e f
g h i j k l	g h i j k l	g h i j k l	g h i j k l	g h i j k l

6 <sup>th</sup> Part:	7 <sup>th</sup> Part:	8 <sup>th</sup> Part:	9 <sup>th</sup> Part:	10 <sup>th</sup> Part:	11 <sup>th</sup> Part
a b c d e f	a b c d e f	a b c d e f	a b c d e f	a b c d e f	a b c d e f
g h i j k l	g h i j k l	g h i j k l	g h i j k l	g h i j k l	g h i j k l

Books, eBooks, Video Course, FREE Workbook & FREE Online  
Training on Vedic Speed Mathematics, C & Python Programming:  
[www.Speed16.com/books/vm](http://www.Speed16.com/books/vm)

## DIVISION

**Vinculum Number:** A number that has atleast one vinculum digit is called vinculum number. Notation: Either dotted or dash above the number. **Ex.**  $1\ddot{4}2$ ;  $\bar{9}641$ ;  $2\ddot{2}3$ ;  $6\ddot{2}3\bar{8}$ ;  $8\ddot{4}2\bar{3}$ ;  $\bar{8}$ ;  $\bar{3}$

$$1\ddot{4}2 = 100 - 40 + 2 = 62;$$

$$\bar{9}641 = -9000 + 600 + 40 + 1 = -8359$$

$$2\ddot{2}3 = 200 - 20 + 3 = 183$$

$$6\ddot{2}3\bar{8} = 6000 - 200 + 30 - 8 = 5822$$





$$8\ddot{4}2\bar{3} = 8000 - 400 - 20 - 3 = 7577$$

$$\bar{8} = -8$$

$$\bar{3} = -3$$

### Division using Transpose and Apply


**Ex.1:**  $123 \div 11$ ; Here Base=10 and Surplus=1; Negation of Surplus=-1=  $\bar{1}$

Step 1:	Step 2:	Step 3:	Step 4:	Step 5:
$11) 12\dot{3}$ $\underline{-1}$	$11) 12\dot{3}$ $\underline{-1}$ 	$11) 12\dot{3}$ $\underline{-1 \quad -1}$ 	$11) 12\dot{3}$ $\underline{-1 \quad -1}$ 	$11) 12\dot{3}$ $\underline{-1 \quad -1 \quad -1}$ 
	$1$	$1$	$1 \quad 1\dot{}$	$1 \quad 1\dot{} \quad 2$

**Answer: 11 (Q); 2 (R)**

**Ex.2: 1332÷114**

B: 100; S: 14



$$(12-1) \mid (114-36)$$

$$11 \mid 78$$

$$Q: 11; R: 78$$

**Division by FLAG (ध्वजांक) Method**

**Ex.2: Divide 949 by 22**

Step 1	Step 2	Step 3	Answer
$2^2 \# 9_1 4   9$ 4 $14 - (2 \times 4) = 14 - 8 = 6$ $6 \div 2 = 3(Q) 0(R)$	$2^2 \# 9_1 4   0 9$ 4 3 $9 - (2 \times 3) = 9 - 6 = 3$ $3 \div 2 = 1(Q) 1(R)$	$2^2 \# 9_1 4   0 9_1$ 4 3   1	<b>43.1</b> (Upto 1 Decimal Point)

Books, eBooks, Video Course, FREE Workbook & FREE Online Training on Vedic Speed Mathematics, C & Python Programming:  
[www.Speed16.com/books/vm](http://www.Speed16.com/books/vm)

## DECIMALS, FRACTIONS AND PERCENTAGES

**Fractions:** Is an expression that indicates the quotient of two quantities. **Ex.**  $\frac{p}{q}$ ;  $\frac{1}{2}$ ;  $\frac{1}{3}$ ;  $\frac{42}{2.1}$ ;  $\frac{445.61}{2.64}$ ;  $\frac{785}{2}$ ;  $\frac{36}{3}$ ;  $-\frac{40}{20}$ ;  $-\frac{1}{9}$  etc. Upper part is Numerator (N) and Lower part is Denominator. The Denominator can not be zero.

### Basic Operations on Decimals

#### Addition:

**Ex.1:**  $0.32+13.35+45.058+696.368+31.004$

0.32 (320/1000);

13.35 (13350/1000);

45.058 (45058/1000);

696.368 (696368/1000);

31.004 (31004/1000)

$$\frac{320}{1000} + \frac{13350}{1000} + \frac{45058}{1000} + \frac{696368}{1000} + \frac{31004}{1000} = \frac{786100}{1000} = 786.1$$

**Ex.2:**  $143+365.9+0.04+36.02+6986.36+7469.3$

143 (14300/100);

365.9 (36590/100);

0.04 (4/100);

36.02 (3602/100);

6986.36 (698636/100);

7469.3(746930/100)

$$\frac{14300}{100} + \frac{36590}{100} + \frac{4}{100} + \frac{3602}{100} + \frac{698636}{100} + \frac{746930}{100} = \frac{1500062}{100} = 1500.62$$



### Subtraction:

**Ex.1:** 133.45-45.058-6.368-31.004

133.45 (133450/1000);

45.058 (45058/1000);

6.368 (6368/1000);

31.004 (31004/1000)

$$\frac{133450}{1000} - \frac{45058}{1000} - \frac{6368}{1000} - \frac{31004}{1000} = \frac{51020}{1000} = 51.02$$

**Ex.2:** 143-65.9-0.004-36.002-6.986-7.3

143 (143000/1000);

65.9 (65900/1000);

0.004 (4/1000);

36.002 (36002/100);

6.986 (6986/1000);

7.3(7300/1000)

$$\frac{143000}{1000} - \frac{65900}{1000} - \frac{4}{1000} - \frac{36002}{1000} - \frac{6986}{1000} - \frac{7300}{1000} = \frac{26808}{1000} = 26.808$$

### Multiplication:

<b>Ex.1:</b> <b>3.6×6.8</b>	<b>Ex.2:</b> <b>43.63×34.2</b>	<b>Ex.3:</b> <b>42.36×67.363</b>	<b>Ex.4:</b> <b>3.6×63.68×1.3</b>
$\frac{36}{10} \times \frac{68}{10}$	$\frac{4363}{100} \times \frac{3420}{100}$	$\frac{42360}{1000} \times \frac{67363}{1000}$	$\frac{360}{100} \times \frac{6368}{100} \times \frac{130}{100}$
$= \frac{2448}{100}$	$= \frac{14921460}{10000}$	$= \frac{2853496680}{1000000}$	$= \frac{298022400}{1000000}$
$= 24.48$	$= 1492.146$	$= 2853.49668$	$= 298.0224$
<b>24.48</b>	<b>1492.146</b>	<b>2853.49668</b>	<b>298.0224</b>

## Basic Operations on Fractions:

### Addition:

**Case 1: When Denominators are same:** Just add numerators; place denominator as it is and Reduce.

**Case 2: When one Denominator is factor of another:** Make same denominator by multiplying lower denominator and its numerator with the factor and add using case 1.

**Case 3: Cross Product:** Numerator: sum of cross products; Denominator: product of both denominators.

Ex.1:	Ex.2:	Ex.3:	Ex.4:
Case:1	Case:2	Case:3	Case:3
$\frac{68}{10} + \frac{36}{10}$	$\frac{150}{21} + \frac{1350}{63}$	$\frac{35}{16} + \frac{12}{20}$	$\frac{36}{10} + \frac{25}{12} + \frac{11}{14}$
$\frac{68 + 36}{10}$	$\frac{150 \times 3}{21 \times 3} + \frac{1350}{63}$	$\frac{(35 \times 20) + (16 \times 12)}{(16 \times 20)}$	$\frac{(36 \times 12) + (10 \times 25)}{(10 \times 12)} + \frac{11}{14}$
$\frac{104}{10}$	$\frac{450}{63} + \frac{1350}{63}$	$\frac{(700) + (192)}{(320)}$	$\frac{682}{120} + \frac{11}{14}$
$\frac{52}{5}$	$\frac{450 + 1350}{63}$	$\frac{892}{320}$	$\frac{(682 \times 14) + (120 \times 11)}{(120 \times 14)}$
	$\frac{1800}{63} = \frac{200}{7}$	$\frac{223}{80}$	$\frac{10868}{1680} = \frac{2717}{420}$
$\frac{52}{5}$	$\frac{200}{7}$	$\frac{223}{80}$	$\frac{2717}{420}$

### Subtraction:

**Case 1: When Denominators are same:** Just subtract numerators; place denominator as it is and Reduce.

**Case 2: When one Denominator is factor of another:** Make same denominator by multiplying lower denominator and its numerator with the factor and subtract using case 1.

**Case 3: Cross Product:** Numerator: Difference of cross products;  
Denominator: product of both denominators.

Ex.1:	Ex.2:	Ex.3:	Ex.4:
Case:1	Case:2	Case:3	Case:3
$\frac{68}{10} - \frac{36}{10}$	$\frac{150}{21} - \frac{1350}{63}$	$\frac{35}{16} - \frac{12}{20}$	$\frac{36}{10} - \frac{25}{12} - \frac{11}{14}$
$\frac{68 - 36}{10}$	$\frac{150 \times 3}{21 \times 3} - \frac{1350}{63}$	$\frac{(35 \times 20) - (16 \times 12)}{(16 \times 20)}$	$\frac{(36 \times 12) - (10 \times 25)}{(10 \times 12)} - \frac{11}{14}$
$\frac{32}{10}$	$\frac{450}{63} - \frac{1350}{63}$	$\frac{(700) - (192)}{(320)}$	$\frac{182}{120} - \frac{11}{14}$
$\frac{16}{5}$	$\frac{450 - 1350}{63}$	$\frac{508}{320} = \frac{127}{80}$	$\frac{(182 \times 14) - (120 \times 11)}{(120 \times 14)}$
	$-\frac{900}{63} = -\frac{100}{7}$		$\frac{1228}{1680} = \frac{307}{420}$
$\frac{16}{5}$	$-\frac{100}{7}$	$\frac{127}{80}$	$\frac{307}{420}$

### Multiplication:

**Option 1:** Multiply Numerator with numerator and denominator with denominator and REDUCE.

**Option 2:** Write factors and cancel common factors. Then Multiply Numerator with numerator and denominator with denominator and REDUCE.

Ex.1:	Ex.2:	Ex.3:
$\frac{36}{10} \times \frac{68}{10}$	$\frac{84}{44} \times \frac{66}{42}$	$\frac{360}{210} \times \frac{630}{60} \times \frac{30}{90}$
$= \frac{2448}{100}$	$= \frac{21 \times 4}{22 \times 2} \times \frac{22 \times 3}{21 \times 2}$	$= \frac{60 \times 6}{210} \times \frac{210 \times 3}{60} \times \frac{30}{3 \times 30}$
$= \frac{612}{25}$	$= \frac{1}{1} \times \frac{3}{1} = 3$	$= \frac{6}{1} \times \frac{1}{1} \times \frac{1}{1} = 6$
$\frac{612}{25}$	<b>3</b>	<b>6</b>

**Division:** Write the first set as it is; replace  $\div$  (division sign) by  $\times$  (multiplication sign); Exchange Numerator and Denominator of second set; Perform Multiplication.

$$\frac{N1}{D1} \div \frac{N2}{D2} = \frac{N1}{D1} \times \frac{D2}{N2}$$

$$\frac{84}{44} \div \frac{42}{66} = \frac{84}{44} \times \frac{66}{42} = \frac{21 \times 4}{22 \times 2} \times \frac{22 \times 3}{21 \times 2} = \frac{4}{2} \times \frac{3}{2} = 3$$

## Reciprocals

**Ending in 9:**

Ex. 1:  $\frac{1}{19}$ ?

**A:** Denominator is 19. Positive Osculator of 19 is 2. (Go through “Divisibility” Unit to understand osculator concept).

$$\frac{1}{19} \approx \frac{1}{20} = \frac{0.1}{2}$$

Now for  $\frac{0.1}{2}$

Divisor	2	2	2	2	2	2	2	2	2	2	2	2	2
Dividend	0.1	1	10	-	5	12	6	3	11	15	17	18	9
Quotient	-	0.	0	5	2	6	3	1	5	7	8	9	4
Remainder	-	-	-	0	1	0	0	1	1	1	1	0	1

Divisor is 2 in all cases. Initial Dividend is 0.1. We can not divide 0.1 by 2 so in quotient we need to give decimal point. New Dividend is 1. Again we can not divide 1 by 2 so in quotient we need to add zero. New Dividend is 10. We divide 10 by 2. Q is 5 and R is 0. New Dividend is RQ (05). Remainder and then append Quotient.  $5 \div 2$ ; Q=2; R=1; Next Dividend=12 ( $\because R=1$  & Q=2). Like this we go on dividing. Final answer: Quotient Row.

$$\frac{1}{19} = 0.052631578 \dots$$

# POLYNOMIALS

## Multiplication using Criss Cross Method




**Note:** Read Multiplication using Criss Cross or Vertically & Crosswise Method from Multiplication Unit

### Steps:






1. Write coefficients of given polynomials one below the other separated by space or vertical bar.
2. Multiply coefficients using formula (refer formula OR graphical representation).
3. Last part is constant. Go on incrementing powers of variable by 1 from right. Second last is x, then  $x^2$ ,  $x^3$ ,  $x^4$ ,  $x^5$  and so on.

**Note:** Write coefficient as zero if any term is absent.





### CASE 1: (2×2; 2×1)




First Part:	Second Part:	Third Part:
		
$(a \times c)$	$(a \times d + b \times c)$	$(b \times d)$

### CASE 2: (3×3; 3×2; 3×1)

First Part:	Second Part:	Third Part:	Fourth Part:	Fifth Part:
				
$(a \times d)$	$(a \times e + b \times d)$	$(a \times f + b \times e + c \times d)$	$(b \times f + c \times e)$	$(c \times f)$

**CASE 3: (4×3; 4×3; 4×2; 4×1)**

First Part:	Second Part:	Third Part:	Fourth Part:
 a b c d e f g h	 a b c d e f g h	 a b c d e f g h	 a b c d e f g h
(a×e)	(a×f) + (b×e)	(a×g) + (b×f) + (c×e)	(a×h) + (b×g) + (c×f) + (d×e)

Fifth Part:	Sixth Part:	Seventh Part:
 a b c d e f g h	 a b c d e f g h	 a b c d e f g h
(b×h) + (c×g) + (d×f)	(c×h) + (d×g)	(d×h)

Ex.1: (x <sup>3</sup> +5x <sup>2</sup> +3x+2) (2x <sup>3</sup> -4x <sup>2</sup> -7x+3)
$\begin{array}{rrrr} 1 & 5 & 3 & 2 \\ 2 & -4 & -7 & 3 \end{array}$
$(1 \times 2) \mid (1 \times -4) + (5 \times 2) \mid (1 \times -7) + (5 \times -4) + (3 \times 2) \mid (1 \times 3) + (5 \times -7) + (3 \times -4) + (2 \times 2) \mid (5 \times 3) + (3 \times -7) + (2 \times -4) \mid (3 \times 3) + (2 \times -7) \mid (2 \times 3)$
$\begin{array}{ccccccc} 2 & \mid & 6 & \mid & -21 & \mid & -40 & \mid & -14 & \mid & -5 & \mid & 6 \\ 2x^6 & + & 6x^5 & - & 21x^4 & - & 40x^3 & - & 14x^2 & - & 5x & + & 6 \end{array}$
$2x^6 + 6x^5 - 21x^4 - 40x^3 - 14x^2 - 5x + 6$

Books, eBooks, Video Course, FREE Workbook & FREE Online Training on Vedic Speed Mathematics, C & Python Programming:  
[www.Speed16.com/books/vm](http://www.Speed16.com/books/vm)



**CASE 5: (6×6; 6×5; 6×4; 6×3; 6×2; 6×1) (Do it Yourself)**

First Part:	Second Part:	Third Part:	Fourth Part:	Fifth Part:
a b c d e f	a b c d e f	a b c d e f	a b c d e f	a b c d e f
g h i j k l	g h i j k l	g h i j k l	g h i j k l	g h i j k l

Sixth Part:	Seventh P:	Eighth P:	Nineth P:	Tenth P:
a b c d e f	a b c d e f	a b c d e f	a b c d e f	a b c d e f
g h i j k l	g h i j k l	g h i j k l	g h i j k l	g h i j k l

Ex.1: $(3x^5-2x^4-4x^3+2x^2-3x+3)(4x^5-6x^4+3x^3-2x^2+6x+2)$
3   -2   -4   2   -3   3
4   -6   3   -2   6   2
$12x^{10}-26x^9+5x^8+20x^7-14x^6+38x^5-59x^4+19x^3-20x^2+12x+6$



**Ex.1:**  $(x^3 + 9x^2 + 20x + 12) \div (x + 1)$

$\frac{(x+1)}{-1} \quad x^3 + 9x^2 + 20x + 12$

$x^2 \quad 8x \quad 12 \quad 0$

**Q:**  $x^2 + 8x + 12 = (x + 2)(x + 6)$  **R:** 0

**Q:**  $x^2 + 8x + 12 = (x + 2)(x + 6)$  **R:** 0

**Ex.3:**  $(3x^4 - 2x^3 + x^2 - 2x + 3) \div (x^2 - 2x + 6)$

$\frac{(x^2 - 2x + 6)}{+2 \mid -6}$

$3x^4 - 2x^3 + x^2 - 2x + 3$

$+6 \quad -18$

$8 \quad -24$

$-18 \quad 54$

-----

$3 \quad +4 \quad -9 \quad -44 \quad +57$

**Q:**  $3x^2 + 4x - 9$  **R:**  $-44x + 57$

**Q:**  $3x^2 + 4x - 9$  **R:**  $-44x + 57$

**Ex.4:**  $(2x^5 + 2x^4 - x^3 + x^2 - 2x + 2) \div (x^2 + 3x - 4)$

$\frac{(x^2+3x-4)}{-3 \mid +4} 2x^5 + 2x^4 - x^3 + x^2 - 2x + 2$

$-6 \quad 8 \quad 12 \quad -16 \quad -57 \quad 76 \quad 216 \quad -288$

$2 \quad -4 \quad 19 \quad -72 \quad 290 \quad -286$

**Q:**  $2x^3 - 4x^2 + 19x - 72$  **R:**  $290x - 286$

**Q:**  $2x^3 - 4x^2 + 19x - 72$  **R:**  $290x - 286$

## SIMPLE EQUATIONS

Simple Algebraic Equations contains only one variable.

	<b>TYPE-1</b>	<b>TYPE-2</b>
General Form	$px+q=r x+s$	$(x+p)(x+q)=(x+r)(x+s)$
Value of x	$x = \frac{s-q}{p-r}$	if $p \times q = r \times s$ then <b><math>x=0</math></b> else $x = \frac{rs-pq}{p+q-r-s}$
<b>Ex.1</b>	$5x+3=3x+9$ $x = \frac{9-3}{5-3}$ $= \frac{6}{2} = \mathbf{3}$	$(x+4)(x+3)=(x+2)(x+6)$ $p \times q = r \times s$ $4 \times 3 = 2 \times 6 \Rightarrow 12=12$ <b><math>\therefore x=0</math></b>
<b>Ex.2</b>	$3x-3=4x+7$ $x = \frac{7-(-3)}{3-4} = \frac{10}{-1} = \mathbf{-10}$	$(x+3)(x+5)=(x+2)(x+4)$ $p \times q \neq r \times s$ $x = \frac{2 \times 4 - 3 \times 5}{3+5-2-4}$ $= -\frac{7}{2}$
<b>Ex.3</b>	$6x+6=8x+8$ $x = \frac{8-6}{6-8} = \frac{2}{-2} = \mathbf{-1}$ <b>OR</b> $6x+6=8x+8$ $6(x+1)=8(x+1)$ $x+1=0$ <b><math>x=-1</math></b>	$(x-2)(x+3)=(x+4)(x-5)$ $p \times q \neq r \times s$ $x = \frac{4 \times (-5) - (-2) \times 3}{-2+3-4-(-5)}$ $= \mathbf{-7}$

	<b>TYPE-3</b>	<b>TYPE-4</b>
General Form	$\frac{px+q}{rx+s} = \frac{t}{u}$	$\frac{a}{x+p} + \frac{b}{x+q} = 0$
Value of x	$x = \frac{ts-ut}{up-tr}$	$x = -\frac{aq+bp}{a+b}$
<b>Ex.1</b>	$\frac{2x+3}{3x+5} = \frac{3}{2}$ $x = \frac{3 \times 5 - 2 \times 3}{2 \times 2 - 3 \times 3} = -\frac{9}{5}$	$\frac{3}{x+2} + \frac{4}{x+3} = 0$ $x = -\frac{3 \times 3 + 4 \times 2}{3+4} = -\frac{17}{7}$
<b>Ex.2</b>	$\frac{x-4}{2x+2} = \frac{7}{3}$ $x = \frac{7 \times 2 - 3 \times (-4)}{3 \times 1 - 7 \times 2} = -\frac{26}{11}$	$\frac{5}{x-3} + \frac{6}{x+4} = 0$ $x = -\frac{5 \times 4 + 6 \times -3}{5+6} = -\frac{2}{11}$
<b>Ex.3</b>	$\frac{3x-2}{x-3} = \frac{4}{3}$ $x = \frac{4(-3) - 3(-2)}{3 \times 3 - 4 \times 1} = -\frac{6}{5}$	$\frac{4}{x+2} + \frac{4}{x+3} = 0$ $x = -\frac{4 \times 3 + 4 \times 2}{4+4} = -\frac{5}{2}$ <b>OR</b> $x+2+x+3=0$ $2x+5=0$ $x = -\frac{5}{2}$

### Solution using “If the Set is same, it is ZERO”

4.

A. If  $N1+N2=D1+D2$  then

$$N1+N2=D1+D2=0$$

$$\frac{3x+5}{5x+4} = \frac{4x+6}{2x+7}$$

$$3x+5+4x+6=7x+11$$

$$5x+4+2x+7=7x+11$$

$$\therefore 7x+11=0$$

$$x = -\frac{11}{7}$$

**B.** If  $m(N1+N2)=n(D1+D2)$  then

$N1+N2=D1+D2=0$  ( $m, n$ =common factors)

$$\frac{7x+2}{2x+1} = \frac{5x+6}{x+1}$$

$$7x+2+5x+6=12x+8=4(3x+2)$$

$$2x+1+x+1=3x+2=1(3x+2)$$

$$\therefore 3x+2=0$$

$$x = -\frac{2}{3}$$

**C.** If  $m(N1-D1)=n(N2-D2)$  then

$N1-D1=N2-D2=0$  ( $m, n$ =common factors)

$$\frac{3x+5}{5x+4} = \frac{4x+6}{2x+7}$$

$$3x+5-5x-4 = -2x+1 \text{ and } 4x+6-2x-7 = 2x-1 = -1(-2x+1)$$

$$\therefore -2x+1=0; 2x=1; x = \frac{1}{2}$$

Books, eBooks, Video Course, FREE Workbook & FREE Online  
Training on Vedic Speed Mathematics, C & Python Programming:

[www.Speed16.com/books/vm](http://www.Speed16.com/books/vm)

# QUADRATIC EQUATIONS

## Solution using Calculus

This sutra tells that: Differentiation of expression is equal to square root of discriminant.

$$E \Rightarrow ax^2 + bx + c = 0 \text{ ----(Expression)}$$

$$D(E) = 2ax + b \text{ ----(Differentiation of Expression)}$$

$$\text{Discriminant} = b^2 - 4ac$$

$$\text{Square Root of Discriminant} = \sqrt{b^2 - 4ac}$$

$$\therefore 2ax + b = \sqrt{b^2 - 4ac}$$

(Differentiation of expression = square root of discriminant.)

$$\therefore x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

**Ex.1:  $x^2 + 7x + 12 = 0$**

using Calanā kalanābhyām (चलनकलनाभ्याम्)
<p><b>a=1; b=7; c=12</b></p> $\sqrt{b^2 - 4ac} = \sqrt{7^2 - 4 \times 1 \times 12} =$ $\sqrt{1} = \pm 1$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-7 \pm 1}{2 \times 1}$ <p><math>x = (-7+1)/2</math> OR <math>x = (-7-1)/2</math>; <math>x = -6/2</math> OR <math>x = -8/2</math>; <math>x = -3</math> OR <math>x = -4</math></p>
<b>x = -3 OR x = -4</b>

Ex.2: $x^2+6x+8=0$	Ex.3: $3x^2+12x+7=0$	Ex.4: $6x^2+8x+5=0$
<b>a=1; b=6; c=8</b> $\sqrt{b^2 - 4ac}$ $= \sqrt{6^2 - 4 \times 1 \times 8}$ $= \sqrt{4} = \pm 2$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $x = \frac{-6 \pm 2}{2 \times 1}$ $x = (-6 \pm 2)/2$ OR $x = (-6 - 2)/2$ $x = -4/2$ OR $x = -8/2$ $x = -2$ OR $x = -4$	<b>a=3; b=12; c=7</b> $\sqrt{b^2 - 4ac}$ $= \sqrt{12^2 - 4 \times 3 \times 7}$ $= \sqrt{60}$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $x = \frac{-12 \pm \sqrt{60}}{2 \times 3}$ $x = \frac{-12 \pm 2\sqrt{15}}{6}$ $x = \frac{2(-6 \pm \sqrt{15})}{6}$ $x = \frac{(-6 \pm \sqrt{15})}{3}$	<b>a=6; b=8; c=5</b> $\sqrt{b^2 - 4ac}$ $= \sqrt{8^2 - 4 \times 6 \times 5}$ $= \sqrt{-56}$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $x = \frac{-8 \pm \sqrt{-56}}{2 \times 6}$ $x = \frac{-8 \pm \sqrt{4 \times 14 \times i^2}}{12}$ $x = \frac{-8 \pm 2i\sqrt{14}}{12}$ $x = \frac{-4 \pm i\sqrt{14}}{6}$
<b>x = -2 OR x = -4</b>	<b>x = <math>\frac{(-6 \pm \sqrt{15})}{3}</math></b>	<b>x = <math>\frac{-4 \pm i\sqrt{14}}{6}</math></b>

### Reciprocals By Mere Observation {Sub Sutra 13: Vilokanam}

Given Expression:	Convert it into:	Values:
$\frac{x_1}{x_2} \pm \frac{x_2}{x_1} = \frac{a}{b}$	$\frac{x_1}{x_2} \pm \frac{x_2}{x_1} = \frac{c}{d} \pm \frac{d}{c}$	$\frac{x_1}{x_2} = \frac{c}{d}$ or $\pm \frac{d}{c}$

### TYPE-1 (Plus)

Ex.1	Ex.2	Ex.3	Ex.4
$x + \frac{1}{x} = \frac{10}{3}$ $\frac{x}{1} + \frac{1}{x} = \frac{10}{3}$ $\frac{x}{1} + \frac{1}{x} = \frac{3}{1} + \frac{1}{3}$ $\frac{x}{1} = \frac{3}{1}$ OR $\frac{1}{3}$	$x + \frac{1}{x} = \frac{65}{8}$ $\frac{x}{1} + \frac{1}{x} = \frac{65}{8}$ $\frac{x}{1} + \frac{1}{x} = \frac{8}{1} + \frac{1}{8}$ $\frac{x}{1} = \frac{8}{1}$ OR $\frac{1}{8}$	$\frac{4x+3}{7x+1} + \frac{7x+1}{4x+3} = \frac{73}{24}$ $\frac{73}{24} = \frac{3}{8} + \frac{8}{3}$ $\frac{4x+3}{7x+1} = \frac{3}{8}$ OR $\frac{8}{3}$ Solve for x	$\frac{5x+2}{x-3} + \frac{x-3}{5x+2} = \frac{74}{35}$ $\frac{74}{35} = \frac{5}{7} + \frac{7}{5}$ $\frac{5x+2}{x-3} = \frac{5}{7}$ OR $\frac{7}{5}$ Solve for x
<b>3 OR 1/3</b>	<b>8 OR 1/8</b>	<b>-21/11 OR 1/44</b>	<b>-29/30 OR -31/18</b>

### TYPE-2 (Minus):

Ex.5	Ex.6	Ex.7	Ex.8
$x - \frac{1}{x} = \frac{7}{12}$ $\frac{x}{1} - \frac{1}{x} = \frac{7}{12}$ $x - \frac{1}{x} = \frac{4}{3} - \frac{3}{4}$ $\frac{x}{1} = \frac{4}{3} \text{ OR } -\frac{3}{4}$	$x - \frac{1}{x} = \frac{11}{30}$ $\frac{x}{1} - \frac{1}{x} = \frac{11}{30}$ $x - \frac{1}{x} = \frac{6}{5} - \frac{5}{6}$ $\frac{x}{1} = \frac{6}{5} \text{ OR } -\frac{5}{6}$	$\frac{4x-3}{7x-1} - \frac{7x-1}{4x-3} =$ $\frac{73}{24}$ $\frac{73}{24} = \frac{3}{8} - \frac{8}{3}$ $\frac{4x-3}{7x-1} = \frac{3}{8} \text{ OR } \frac{8}{3}$ <p>Solve for x</p>	$\frac{5x-2}{x-3} - \frac{x-3}{5x-2} =$ $\frac{21}{10}$ $\frac{21}{10} = \frac{5}{2} - \frac{2}{5}$ $\frac{5x-2}{x-3} = \frac{5}{2} \text{ OR } -\frac{2}{5}$ <p>Solve for x</p>
<b>4/3 OR -3/4</b>	<b>6/5 OR -5/6</b>	<b>-21/11 OR 1/44</b>	<b>-11/5 OR 16/27</b>

**Solution using “If the Set is same, it is ZERO”**

#### First Factor:

If  $m(N1+N2)=n(D1+D2)$  then

$N1+N2=D1+D2=0$  (**m,n**=common factors)

#### Second Factor:

If  $m(N1-D1)=n(N2-D2)$  then

$N1-D1=N2-D2=0$  (**m,n**=common factors)

Ex.1	Ex.2
$\frac{3x+5}{5x+4} = \frac{4x+6}{2x+7}$ $N1+N2=3x+5+4x+6=7x+11$ $D1+D2=5x+4+2x+7=7x+11$ $N1-D1=3x+5-5x-4=-2x+1$ $N2-D2=4x+6-2x-7=2x-1=-1(-2x+1)$ <p>First Factor:</p> $7x+11=0; 7x=-11; x=-11/7$ <p>Second Factor:</p> $-2x+1=0; 2x=1; x=1/2$ <p><b>x = -11/7 OR x = 1/2</b></p>	$\frac{2x+6}{4x+3} = \frac{6x+3}{4x+6}$ $N1+N2=2x+6+6x+3=8x+9$ $D1+D2=4x+3+4x+6=8x+9$ $N1-D1=2x+6-4x-3=-2x+3$ $N2-D2=6x+3-4x-6=2x-3=-1(-2x+3)$ <p>First Factor:</p> $8x+9=0; 8x=-9; x=-9/8$ <p>Second Factor:</p> $-2x+3=0; 2x=3; x=3/2$ <p><b>x = -9/8 OR x = 3/2</b></p>

## CUBIC EQUATIONS

### CASE 3:

$$\frac{1}{D1} + \frac{1}{D2} = \frac{1}{D3} + \frac{1}{D4}$$

if  $D1+D2=D3+D4$

First Value:  $D1+D2=0$

Second Value:  $D1-D2= D3-D4$

Third Value:  $D1-D2= D4-D3$

**Ex:** 
$$\frac{1}{4x+2} + \frac{1}{7x-6} = \frac{1}{5x+4} + \frac{1}{6x-8}$$

$$D1=4x+2; D2=7x-6;$$

$$D3=5x+4; D4=6x-8$$

$$D1+D2=11x-4; D3+D4=11x-4;$$

$$D1-D2=-3x+8; D3-D4=-x+12 \quad D4-D3=x-12$$

First:  $11x-4=0; x=4/11;$

Second:  $-3x+8=-x+12; x=-2$

Third:  $-3x+8=x-12; x=5$

Books, eBooks, Video Course, FREE Workbook & FREE Online Training on Vedic Speed Mathematics, C & Python Programming:

[www.Speed16.com/books/vm](http://www.Speed16.com/books/vm)



# SIMULTANEOUS EQUATIONS

## Solution using Criss Cross Method

General Form:  $a_1x + b_1y = c_1$  and  $a_2x + b_2y = c_2$ ; Values are

$$x = \frac{b_1c_2 - b_2c_1}{a_2b_1 - a_1b_2} \quad y = \frac{a_2c_1 - a_1c_2}{a_2b_1 - a_1b_2}$$

$a_1x + b_1y = c_1$ $a_2x + b_2y = c_2$	$a_1x + b_1y = c_1$ $a_2x + b_2y = c_2$	$a_1x + b_1y = c_1$ $a_2x + b_2y = c_2$
--	--	--

EX.1:	EX.2:	EX.3:
$3x + 7y = 27 \text{ -- (I)}$ $5x + 2y = 16 \text{ -- (II)}$ $x = \frac{7 \times 16 - 27 \times 2}{5 \times 7 - 3 \times 2}$ $= \frac{112 - 54}{35 - 6} = \frac{58}{29} = 2$ $y = \frac{5 \times 27 - 3 \times 16}{5 \times 7 - 3 \times 2}$ $= \frac{135 - 48}{35 - 6} = \frac{87}{29} = 3$	$2x + y = 7 \text{ -- (I)}$ $3x - y = 8 \text{ -- (II)}$ $x = \frac{1 \times 8 - (-1) \times 7}{3 \times 1 - 2 \times -1}$ $= \frac{8 + 7}{3 + 2} = \frac{15}{5} = 3$ $y = \frac{3 \times 7 - 2 \times 8}{3 \times 1 - 2 \times -1}$ $= \frac{21 - 16}{3 + 2} = \frac{5}{5} = 1$	$-2x + y = 1 \text{ -- (I)}$ $-3x + 2y = 5 \text{ -- (II)}$ $x = \frac{1 \times 5 - 2 \times 1}{-3 \times 1 - (-2) \times 2}$ $= \frac{5 - 2}{-3 + 4} = \frac{3}{1} = 3$ $y = \frac{-3 \times 1 - (-2) \times 5}{-3 \times 1 - (-2) \times 2}$ $= \frac{-3 + 10}{-3 + 4} = \frac{7}{1} = 7$
<b>x=2 and y=3</b>	<b>x=3 and y=1</b>	<b>x=3 and y=7</b>

# INTRODUCTION TO VEDIC MATHEMATICS

## Sutras: Meaning and Usage

**Sutra 3:** Ūrdhva – tiryagbhyām (ऊर्ध्वतिर्यग्भ्याम्)

**Meaning:** Vertically and Crosswise.

**Usage:** This is used to multiply given numbers vertically and crosswise.

$$\begin{array}{r} ab \\ \times cd \\ \hline \end{array}$$

$$= (a \times c) \mid (a \times d + b \times c) \mid (b \times d)$$

$$\begin{array}{r} abc \\ \times def \\ \hline \end{array}$$

$$= (a \times d) \mid (a \times e + b \times d) \mid (a \times f + b \times e + c \times d) \mid (b \times f + c \times e) \mid (c \times f)$$

**Ex: 25×67**

First Part:	Second Part:	Third Part:
$\begin{array}{cc} a & b \\ \vdots & \vdots \\ c & d \end{array}$	$\begin{array}{cc} a & b \\ \swarrow \searrow & \swarrow \searrow \\ c & d \end{array}$	$\begin{array}{cc} a & b \\ \vdots & \vdots \\ c & d \end{array}$
$(a \times c)$ $(2 \times 6 = 12)$	$(a \times d + b \times c)$ $(2 \times 7 + 5 \times 6 = 44)$	$(b \times d)$ $(5 \times 7 = 35)$

12|44|35

12|44+3|5

12|47|5

12+4|7|5

16|7|5

1675

**Applications:** to find Product of given numbers and polynomials.

**Refer Units:** Multiplication and Polynomials.

---

**Sutra 5:** Sūnyam Samyasamuccaye (शून्यं साम्यसमुच्चये)

**Meaning:** If the Samuccay (समुच्चय or समूह or Set) is same, it is ZERO.

**Usage:**

**CASE 3:** If the numerical numerators of two fractions are same, then the sum of denominators is ZERO.

**Ex.**  $\frac{3}{4x+5} + \frac{3}{7x-3} = 0$

Here  $3=3$ ; so  $4x+5+7x-3=0$ ;

$11x+2=0$ ;  $11x=-2$ ;  $x=-2/11$ .

**Ex.**  $\frac{-7}{x^2+6} + \frac{-7}{6x+3} = 0$

Here  $-7=-7$ ; so  $x^2+6+6x+3=0$ ;

$x^2+6x+9=0$ ;  $(x+3)^2=0$ ;  $x=-3$ .

**Ex.**  $\frac{16}{3x+5} - \frac{16}{8x-12} = 0$

$16=16$ ; so  $3x+5-(8x-12)=0$ ;  $3x+5-8x+12=0$ ;  $-5x+17=0$ ;  $x=17/5$ .

**Note:** Here sign is negative, so we have subtracted (D1-D2)

**CASE 4:**

**A.** If  $N1+N2=D1+D2$  then  $N1+N2=D1+D2=0$

$$\frac{3x+5}{5x+4} = \frac{4x+6}{2x+7}$$

$3x+5+4x+6=7x+11$ ;  $5x+4+2x+7=7x+11$

$\therefore 7x+11=0$ ;  $x = -\frac{11}{7}$

**B.** If  $m(N_1+N_2)=n(D_1+D_2)$  then  $N_1+N_2=D_1+D_2=0$   
 ( $m, n$ =common factors)

$$\frac{7x+2}{2x+1} = \frac{5x+6}{x+1}$$

$$7x+2+5x+6=12x+8=4(3x+2)$$

$$2x+1+x+1=3x+2=1(3x+2)$$

$$\therefore 3x+2=0$$

$$x = -\frac{2}{3}$$

**C:** If  $m(N_1-D_1)=n(N_2-D_2)$  then  $N_1-D_1=N_2-D_2=0$  ( $m, n$ =common factors)

$$\frac{3x+5}{5x+4} = \frac{4x+6}{2x+7}$$

$$3x+5-5x-4 = -2x+1 \text{ and } 4x+6-2x-7 = 2x-1 = -1 (-2x+1)$$

$$\therefore -2x+1=0; 2x=1; x = \frac{1}{2}$$

**Sutra 13:** Sopantyadvayamantyam (सोपान्त्यद्वयमन्त्यम्)

**Meaning:** The ultimate and twice the penultimate

**Usage:**

$$\frac{1}{AB} + \frac{1}{AC} = \frac{1}{AD} + \frac{1}{BC}$$

If A, B, C and D are in Arithmetic Progression then  $D+2C=0$

$$\text{Ex. } \frac{1}{(x+3)(x+4)} + \frac{1}{(x+3)(x+5)} = \frac{1}{(x+3)(x+6)} + \frac{1}{(x+4)(x+5)}$$

$$A=(x+3);$$

$$B=(x+4);$$

$$C=(x+5);$$

$$D=(x+6);$$

Here A, B, C & D are in Arithmetic Progression;

So,  $D+2C=0$ ;

$$(x+6)+2(x+5)=0;$$

$$x+6+2x+10=0; 3x+16=0; x=-16/3.$$

---

## Sub Sutras: Meaning and Usage

### Sub Sutra 9: Antyayoreva (अन्त्ययोरेव)

**Meaning:** Only the Last Terms

**Usage:** This sutra is used to solve certain equations of the type:

After ignoring Constants; Numerator and the Denominator of one side are in the ratio of Numerator and Denominator of the other side.

$$\text{if } \frac{m(N1) + c1}{m(D1) + c2} = \frac{N2}{D2} \text{ where } N1 = N2 \text{ and } D1 = D2 \text{ then } \frac{N2}{D2} = \frac{c1}{c2}$$

$m$  is common factor;  $c1$  and  $c2$  are constants.

$$\text{Ex. } \frac{x^2 + x + 2}{x^2 + 2x + 5} = \frac{x + 1}{x + 2}$$

$$\Rightarrow \frac{x(x + 1) + 2}{x(x + 2) + 5} = \frac{x + 1}{x + 2} \text{ then } \frac{x + 1}{x + 2} = \frac{2}{5}; x = -\frac{1}{3}$$

---

### Sub Sutra 12: Vilokanam (विलोकनं)

**Meaning:** By Mere Observation

**Usage:** This sutra is used in solving quadratic and simultaneous equations. Refer Unit: Quadratic Equations.

**Ex:**  $x + \frac{1}{x} = \frac{10}{3}$ ; By Vilokanam (विलोकनं) we say that  $x=3$ .

---

Books, eBooks, Video Course, FREE Workbook & FREE Online Training on Vedic Speed Mathematics, C & Python Programming:

[www.Speed16.com/books/vm](http://www.Speed16.com/books/vm)