

Ergonomics Evaluation Report

Posture Analysis

- Neck Flexion: Yellow
- Shoulder Elevation: Yellow
- Elbow Angle: Red
- Wrist Deviation: Yellow
- Pelvic Tilt: Red

Workstation Analysis

- Monitor: MonitorHeight: Red
- Monitor: ViewingDistance: Green

Pain & Risk Summary

Average Pain Level: 7


Main Pain Region(s): Neck, Elbows/Forearms, Hips/Glutes (VAS 8)

Posture & Ergonomic Corrections

- Reduce pelvic tilt to under 10°: Sit with hips fully back in the chair and adjust seat pan so your pelvis is upright, reducing tilt from 20° to less than 10° for spinal alignment.
- Bend elbows to 100°-110°: Lower armrests or raise seat so elbows are bent at 100°-110°, not 115°, to minimize upper back and shoulder strain.
- Lower shoulder elevation below 5°: Relax shoulders and adjust armrests so elevation is less than 5°, reducing muscle tension in upper back and neck.
- Keep wrist deviation under 10°: Align wrists straight with forearms, using a wrist rest if needed, to reduce deviation from 13° to below 10°.
- Raise monitor by 8-12 cm: Use a monitor riser to elevate your laptop screen so the top is at or just below eye level, correcting a 25 cm deficit.
- Add lumbar support cushion: Place a lumbar cushion at the lower back to support natural spinal curve and reduce pelvic tilt.

Final Advice

Your upper back pain is severe and posture risks are high. Prioritize workstation changes and take frequent breaks. If pain persists or worsens, consult a healthcare professional. Safety note: Always perform any posture or stretching exercises slowly and stop if you feel pain.



Annotated posture analysis from your uploaded photo