

Ergonomics Evaluation Report

Posture Analysis

- Neck Flexion: Yellow
- Shoulder Elevation: Yellow
- Elbow Angle: Red
- Wrist Deviation: Yellow
- Pelvic Tilt: Red

Workstation Analysis

- Monitor: MonitorHeight: Red
- Monitor: ViewingDistance: Green

Pain & Risk Summary

Average Pain Level: 7


Main Pain Region(s): Neck, Elbows/Forearms, Hips/Glutes (VAS 8)

Posture & Ergonomic Corrections

- Reduce pelvic tilt to under 5°: Sit with hips fully back in the chair and use lumbar support to decrease pelvic tilt from 20° to below 5°, aligning the pelvis and lower back per ISO 9241-5.
- Lower elbow angle to 100°: Adjust chair or desk height so elbows are bent at 90-100°, reducing strain from current 116° and supporting forearms per ISO 9241-5.
- Keep neck flexion under 5°: Sit upright and raise your screen so your neck bends less than 5°, minimizing neck strain and following ISO 9241-5 recommendations.
- Reduce wrist deviation to under 5°: Keep wrists straight by aligning keyboard and mouse with forearms, reducing deviation from 13° to under 5° as per ISO 9241-5.
- Raise monitor by 25 cm: Elevate your monitor so the top is at or just below eye level, correcting the 25 cm height deficit and reducing neck flexion per ISO 9241-5.
- Add lumbar support cushion: Place a lumbar cushion at the lower back to maintain natural spinal curve and decrease pelvic tilt, supporting ISO 9241-5 posture.

Final Advice

Your upper back pain is severe and posture risks are high. Prioritize raising your monitor by 25 cm and adding lumbar support immediately. Take frequent breaks every 30-45 minutes. If pain or tingling worsens, consult a healthcare provider. Safety note: Perform all posture exercises gently and stop if you feel pain.



Annotated posture analysis from your uploaded photo