

Ergonomics Evaluation Report

Posture Analysis

- Neck Flexion: Yellow
- Shoulder Elevation: Yellow
- Elbow Angle: Red
- Wrist Deviation: Yellow
- Pelvic Tilt: Red

Workstation Analysis

- Monitor: MonitorHeight: Red
- Monitor: ViewingDistance: Green

Pain & Risk Summary

Average Pain Level: 7


Main Pain Region(s): Neck, Elbows/Forearms, Hips/Glutes (VAS 8)

Posture & Ergonomic Corrections

- Reduce pelvic tilt to under 10°: Sit with hips and pelvis upright; use lumbar support to maintain a neutral spine and decrease pelvic tilt from 20° to below 10° as per ISO 9241-5.
- Decrease elbow angle to 100°: Adjust chair and desk height so elbows are bent at 90-100°, reducing from current 116°, to minimize upper back and shoulder strain.
- Lower shoulder elevation below 5°: Relax shoulders and adjust armrests to keep shoulder elevation under 5°, reducing muscle tension and upper back pain.
- Reduce wrist deviation to under 5°: Keep wrists straight and aligned with forearms; use a wrist rest if needed to decrease deviation from 13° to below 5°.
- Raise monitor by 25 cm: Elevate your monitor so the top of the screen is at or just below eye level, correcting the 25 cm deficit and reducing neck and upper back strain.
- Add lumbar support at chair back: Place a lumbar cushion or rolled towel at the lower back to support the spine and maintain a neutral pelvic position.

Final Advice

Your upper back pain is severe and workstation risks are high. Please prioritize monitor height and lumbar support corrections. Take frequent breaks every 30-45 minutes, and consult a healthcare provider if pain persists or worsens. Safety note: Always perform stretches and posture exercises gently; stop immediately if you feel pain or tingling.



Annotated posture analysis from your uploaded photo