

Ergonomics Evaluation Report

Posture Analysis

- Neck Flexion: Yellow
- Shoulder Elevation: Yellow
- Elbow Angle: Red
- Wrist Deviation: Yellow
- Pelvic Tilt: Red

Workstation Analysis

- Monitor: MonitorHeight: Red
- Monitor: ViewingDistance: Green

Pain & Risk Summary

Average Pain Level: 7


Main Pain Region(s): Neck, Elbows/Forearms, Hips/Glutes (VAS 8)

Posture & Ergonomic Corrections

- Reduce pelvic tilt to under 10°: Sit with hips and pelvis upright, using lumbar support if needed. Avoid slouching or leaning back excessively to decrease pelvic tilt from 20° to under 10°, as per ISO 9241-5.
- Decrease elbow angle to 100°: Adjust chair and desk height so elbows are bent at approximately 100°, not extended beyond 115°, keeping forearms parallel to the floor and close to the body.
- Lower shoulder elevation below 5°: Relax shoulders and adjust armrests or desk height to keep shoulders in a neutral position, reducing elevation from 9.8° to below 5°.
- Reduce wrist deviation to under 5°: Keep wrists straight and aligned with forearms while typing. Use a wrist rest if needed to maintain wrist deviation below 5°.
- Raise monitor by 25 cm: Elevate your laptop screen or monitor by 25 cm so the top of the screen is at or just below eye level, reducing neck flexion and upper back strain.

Final Advice

Your upper back pain is severe and posture risks are high. Please implement the corrections and equipment changes above as soon as possible. If pain persists or worsens, consult a healthcare professional. Always perform any posture or stretching exercises slowly and stop if you feel pain.



Annotated posture analysis from your uploaded photo