

Ergonomics Evaluation Report

Posture Analysis

- Neck Flexion: Yellow
- Shoulder Elevation: Yellow
- Elbow Angle: Red
- Wrist Deviation: Yellow
- Pelvic Tilt: Red

Workstation Analysis

- Monitor: MonitorHeight: Red
- Monitor: ViewingDistance: Green

Pain & Risk Summary

Average Pain Level: 7


Main Pain Region(s): Neck, Elbows/Forearms, Hips/Glutes (VAS 8)

Posture & Ergonomic Corrections

- Reduce elbow angle to 100 degrees: Bring your elbows closer to your body and lower your forearms so that your elbow angle is approximately 100 degrees. This reduces strain on your arms and wrists, as required by ISO 9241-5.
- Correct pelvic tilt to 0-5 degrees: Sit with your pelvis in a neutral position by adjusting your chair and posture so your hips are level and your lower back maintains its natural curve. This helps prevent lower back discomfort and aligns with ISO 9241-5.
- Reduce wrist deviation to under 5 degrees: Keep your wrists straight and aligned with your forearms while typing or using the mouse. Use a wrist rest if needed to maintain this position, as recommended by ISO 9241-5.
- Lower monitor by 25 cm: Adjust your monitor height so the top of the screen is at or slightly below your eye level. This will reduce neck flexion and comply with ISO 9241-5 guidelines.
- Raise chair seat to 46 cm: Increase your chair height so your elbows are at a 100-degree angle and your wrists remain straight while typing. This supports proper arm and wrist posture per ISO 9241-5.

Final Advice

Your reported wrist pain and tingling are concerning and may indicate early nerve compression. Please implement the above corrections immediately. If symptoms persist or worsen, consult a healthcare professional. Always perform any posture or stretching exercises slowly and stop if you feel pain.



Annotated posture analysis from your uploaded photo