

Ergonomics Evaluation Report

Posture Analysis

- Neck Flexion: Yellow
- Shoulder Elevation: Yellow
- Elbow Angle: Red
- Wrist Deviation: Yellow
- Pelvic Tilt: Red

Workstation Analysis

- Monitor: MonitorHeight: Red
- Monitor: ViewingDistance: Green

Pain & Risk Summary

Average Pain Level: 7


Main Pain Region(s): Neck, Elbows/Forearms, Hips/Glutes (VAS 8)

Posture & Ergonomic Corrections

- Reduce pelvic tilt to under 10°: Sit with hips fully back in the chair and adjust your pelvis so that the tilt is less than 10°. Use lumbar support to maintain a neutral spine.
- Lower elbow angle to 100°: Adjust chair and desk height so elbows are bent at 90-100°, keeping forearms parallel to the floor and shoulders relaxed.
- Reduce wrist deviation below 10°: Keep wrists straight while typing by aligning keyboard and mouse with forearms, minimizing side bending.
- Lower shoulder elevation below 5°: Relax shoulders and adjust armrests or desk height so shoulders are not raised while working.
- Raise monitor by 25 cm: Place your laptop or monitor on a riser so the top of the screen is at or just below eye level, reducing neck flexion.

Final Advice

Your upper back and lower back are at high risk due to significant pelvic tilt and poor monitor height, which violate ISO 9241-5 standards. Immediate workstation adjustments and posture corrections are essential to reduce pain and prevent further injury. Take frequent breaks every 30-45 minutes. Safety note: If you experience persistent tingling, numbness, or worsening pain, consult a healthcare professional before starting any exercise or stretching routine.



Annotated posture analysis from your uploaded photo