

POSTURA

Ergonomics Evaluation Report

Posture Analysis

- Neck Flexion: Red
- Shoulder Elevation: Green
- Elbow Angle: Yellow
- Wrist Deviation: Red
- Pelvic Tilt: Red

Workstation Analysis

Pain & Risk Summary

Average Pain Level: 7


Main Pain Region(s): Neck, Wrists/Hands, Hips/Glutes (VAS 8)

Posture & Ergonomic Corrections

- Reduce neck flexion to under 10°: Sit upright and adjust your gaze so your neck bends no more than 10°. This reduces strain and aligns with ISO 9241-5 recommendations.
- Keep wrists straight, under 5° deviation: Position your hands and wrists so they remain in a neutral, straight line while typing, minimizing sideways or upward bending.
- Adjust elbow angle to 100°: Relax your arms so elbows are open to about 100°, reducing muscle tension and improving circulation.
- Sit with pelvis upright, tilt under 15°: Adjust your sitting posture so your pelvis is upright, with minimal backward tilt, to support your lower back and hips.
- Raise laptop screen by 10 cm: Use a riser or stack books to elevate your laptop screen so the top is at or just below eye level, reducing neck flexion.
- Use external keyboard and mouse: Connect an external keyboard and mouse to your laptop to keep wrists straight and elbows at a comfortable angle.
- Add lumbar support cushion: Place a lumbar cushion or rolled towel at your lower back to maintain natural spinal curve and reduce pelvic tilt.

Final Advice

Your current setup poses high ergonomic risks, especially for wrists, neck, and lower back. Severe wrist pain and tingling indicate urgent need for workstation changes. Prioritize using an external keyboard/mouse and raising your laptop screen. Take frequent breaks every 30-45 minutes. If tingling or pain worsens, consult a healthcare professional. Safety note: Always perform posture exercises gently and stop if you experience pain.



Annotated posture analysis from your uploaded photo