

POSTURA

Ergonomics Evaluation Report

Posture Analysis

- Neck Flexion: Red
- Shoulder Elevation: Green
- Elbow Angle: Red
- Wrist Deviation: Yellow
- Pelvic Tilt: Red

Workstation Analysis

- Chair: SeatDepth: Green

Pain & Risk Summary

Average Pain Level: 7


Main Pain Region(s): Neck, Elbows/Forearms, Hips/Glutes (VAS 8)

Posture & Ergonomic Corrections

- Raise monitor by 18 cm: Elevate your monitor so the top edge is at or just below eye level. This will reduce neck flexion from 65° to the ISO-recommended 0–20° range.
- Bend elbows to 90–100 degrees: Adjust your chair and keyboard height so your elbows are bent at 90–100°, not 155°. This reduces strain on your wrists and shoulders.
- Reduce wrist deviation to under 5 degrees: Keep wrists straight by aligning keyboard and mouse with your forearms. Avoid bending wrists sideways more than 5°.
- Sit with pelvis upright at 0–10 degrees: Adjust your seat and posture so your pelvis is upright, not tilted back at 63°. Sit with your back against the chair and feet flat on the floor.
- Raise chair seat to 46 cm: Increase chair height so your elbows are level with the desk, supporting a 90–100° elbow angle.
- Move keyboard 10 cm closer: Position your keyboard so your upper arms hang naturally and elbows stay close to your body.

Final Advice

Your pain intensity and posture risks are severe. Please consult a healthcare professional immediately if symptoms worsen, especially with tingling or numbness. Begin workstation and posture corrections as soon as possible, and take frequent breaks (every 30–45 minutes).



Annotated posture analysis from your uploaded photo