

Ergonomics Evaluation Report

Posture Analysis

- Neck Flexion: Yellow
- Shoulder Elevation: Yellow
- Elbow Angle: Red
- Wrist Deviation: Yellow
- Pelvic Tilt: Red

Workstation Analysis

- Monitor: MonitorHeight: Red
- Monitor: ViewingDistance: Green

Pain & Risk Summary

Average Pain Level: 7

Main Pain Region(s): Neck, Elbows/Forearms, Hips/Glutes (VAS 8)

Posture & Ergonomic Corrections

- Reduce pelvic tilt to under 5°: Sit with your pelvis in a neutral position by ensuring your hips and knees are at the same height and your lower back is supported. Avoid slouching forward or tilting your pelvis excessively.
- Lower elbow angle to 100°: Adjust your arm position so your elbows are bent at approximately 100°, keeping your forearms parallel to the floor and close to your body.
- Decrease wrist deviation to under 5°: Keep wrists straight and aligned with your forearms while typing. Avoid bending wrists sideways or upwards.
- Reduce shoulder elevation to under 5°: Relax your shoulders and keep them level, not hunched or raised, while working at your desk.
- Limit neck flexion to under 5°: Keep your head upright and your chin parallel to the floor, avoiding looking down at your screen.
- Raise monitor by 25 cm: Place your laptop or monitor on a riser or stack of books so the top of the screen is at or just below eye level, reducing neck flexion.
- Add external keyboard at 5 cm height: Use an external keyboard positioned at a height that allows your elbows to rest at 100° and your wrists to remain straight.
- Add lumbar support cushion 5 cm thick: Place a lumbar cushion or rolled towel at your lower back to maintain a neutral pelvic tilt and support your spine.


Final Advice

Your workstation setup and posture present high ergonomic risks, especially for your upper back, wrists, and

lower back. Immediate workstation adjustments and posture corrections are necessary to reduce pain and prevent further injury. Perform the following ergonomic exercises:

1. Neck Retraction: Sit upright, gently pull your chin straight back (not down), hold for 5 seconds, repeat 10 times, 3 times daily.
2. Shoulder Blade Squeeze: Sit or stand, squeeze shoulder blades together, hold for 5 seconds, repeat 10 times, 3 times daily.
3. Wrist Flexor Stretch: Extend arm, palm up, gently pull fingers back with other hand, hold for 15 seconds, repeat 5 times, 2 times daily.
4. Seated Pelvic Tilt: Sit tall, gently tilt pelvis forward and back, hold each position for 5 seconds, repeat 10 times, 3 times daily.
5. Upper Back Extension: Stand, clasp hands behind head, gently arch upper back, hold for 10 seconds, repeat 5 times, 2 times daily.

Safety note: Perform all exercises slowly and stop if you feel sharp pain, numbness, or increased tingling. Consult a healthcare professional if symptoms persist or worsen.



Annotated posture analysis from your uploaded photo