

Ergonomics Evaluation Report

Posture Analysis

- Neck Flexion: Yellow
- Shoulder Elevation: Yellow
- Elbow Angle: Red
- Wrist Deviation: Yellow
- Pelvic Tilt: Red

Workstation Analysis

- Monitor: MonitorHeight: Red
- Monitor: ViewingDistance: Green

Pain & Risk Summary

Average Pain Level: 7

Main Pain Region(s): Neck, Elbows/Forearms, Hips/Glutes (VAS 8)

Posture & Ergonomic Corrections

- Reduce pelvic tilt to under 5°: Sit with hips fully back in the chair, ensuring your pelvis is upright and supported. Use a lumbar support cushion if needed to maintain a neutral pelvic angle below 5°.
- Lower elbow angle to 100°: Adjust your chair and desk height so your elbows are bent at approximately 100°, keeping forearms parallel to the floor and shoulders relaxed.
- Reduce wrist deviation to under 5°: Keep wrists straight and aligned with forearms while typing. Use an ergonomic keyboard or wrist rest to minimize side bending.
- Raise monitor by 25 cm: Place your laptop on a stable riser or stack of books to elevate the top of the screen to eye level, reducing neck flexion and upper back strain.
- Add external keyboard at 5 cm height: Use an external keyboard positioned at a height of 5 cm above your thighs to maintain a neutral wrist and elbow posture.

Final Advice

Your upper back pain is severe and workstation risks are high. Please implement all corrections immediately. If pain persists or worsens, consult a healthcare professional. For exercises:


1. Neck Retraction (Chin Tucks): Sit upright, gently tuck your chin back to align ears over shoulders. Hold for 5 seconds. Repeat 10 times, 3 times daily.
2. Seated Scapular Squeezes: Sit with arms at sides, squeeze shoulder blades together and down. Hold for 5 seconds. Repeat 10 times, 3 times daily.
3. Wrist Flexor Stretch: Extend arm, palm up, gently pull fingers back with other hand. Hold for 15 seconds.

Repeat 3 times per side, 2 times daily.

4. Seated Pelvic Tilt: Sit upright, gently tilt pelvis forward and back. Hold each position for 5 seconds. Repeat 10 times, 2 times daily.

5. Upper Back Extension: Sit or stand, clasp hands behind head, gently arch upper back. Hold for 10 seconds. Repeat 5 times, 2 times daily.

Safety note: Stop any exercise if you feel sharp pain, numbness, or tingling. Consult a healthcare provider if symptoms persist.



Annotated posture analysis from your uploaded photo