

Ergonomics Evaluation Report

Posture Analysis

- Neck Flexion: Yellow
- Shoulder Elevation: Yellow
- Elbow Angle: Red
- Wrist Deviation: Yellow
- Pelvic Tilt: Red

Workstation Analysis

- Monitor: MonitorHeight: Red
- Monitor: ViewingDistance: Green

Pain & Risk Summary

Average Pain Level: 7


Main Pain Region(s): Neck, Elbows/Forearms, Hips/Glutes (VAS 8)

Posture & Ergonomic Corrections

- Reduce pelvic tilt to under 10°: Sit with hips and pelvis upright, ensuring your lower back maintains a natural curve. Use a lumbar support if needed to keep pelvic tilt below 10° as per ISO 9241-5.
- Decrease elbow angle to 100°: Adjust chair and desk height so elbows are bent at 90-100°, keeping forearms parallel to the floor and close to your body.
- Lower shoulder elevation below 5°: Relax shoulders and adjust armrests or desk height so shoulders are not raised. Aim for shoulders to remain relaxed and below 5° elevation.
- Reduce wrist deviation to under 5°: Keep wrists straight while typing. Adjust keyboard and mouse position so wrists remain in a neutral position, with deviation less than 5°.
- Raise monitor by 25 cm: Elevate your monitor so the top of the screen is at or just below eye level, reducing neck flexion and upper back strain.

Final Advice

Your upper back pain is severe and workstation risks are high. Address monitor height, pelvic tilt, and wrist posture immediately. Take frequent breaks every 30-45 minutes. If pain persists or worsens, consult a healthcare professional. Safety note: Always perform posture exercises slowly and stop if you feel pain.



Annotated posture analysis from your uploaded photo