

Ergonomics Evaluation Report

Posture Analysis

- Neck Flexion: Red
- Shoulder Elevation: Green
- Elbow Angle: Yellow
- Wrist Deviation: Red
- Pelvic Tilt: Green

Workstation Analysis

- Chair: SeatDepth: Green

Pain & Risk Summary

Average Pain Level: 7

Main Pain Region(s): Neck, Wrists/Hands (VAS 8)

Posture & Ergonomic Corrections

- Raise monitor by 10 cm: Elevate your monitor so the top edge is at or just below eye level. This will reduce neck flexion from 17.9° to within the ISO 9241-5 recommended range (0-10°), minimizing neck strain.
- Straighten wrists to neutral position: Adjust your typing posture and keyboard angle so your wrists remain straight (0-10° deviation). This will decrease wrist deviation from 29.5° to within ISO 9241-5 limits, reducing risk of nerve compression and tingling.
- Bend elbows to 100 degrees: Bring your chair closer or adjust armrests so your elbows are bent at approximately 100°, not exceeding 110°, as per ISO 9241-5. This supports relaxed shoulders and reduces upper back and shoulder tension.
- Raise monitor by 10 cm: Use a monitor riser or sturdy books to elevate your screen by 10 cm, aligning the top edge with eye level to reduce neck flexion.
- Move keyboard 5 cm closer: Position your keyboard 5 cm closer to the edge of your desk to allow elbows to rest at 100°, minimizing shoulder and upper back strain.
- Use split keyboard for neutral wrists: Switch to a split or tented ergonomic keyboard to maintain wrists in a neutral (0-10°) position, reducing deviation and risk of tingling.

Final Advice

Your reported pain intensity is severe (9/10) in the neck, upper back, and wrists, with tingling and stiffness. Please consult a healthcare professional if symptoms persist or worsen. For all posture exercises, move slowly and stop if you feel pain.



Annotated posture analysis from your uploaded photo