

Ergonomics Evaluation Report

Posture Analysis

- Neck Flexion: Yellow
- Shoulder Elevation: Yellow
- Elbow Angle: Red
- Wrist Deviation: Yellow
- Pelvic Tilt: Red

Workstation Analysis

- Monitor: MonitorHeight: Red
- Monitor: ViewingDistance: Green

Pain & Risk Summary

Average Pain Level: 7

Main Pain Region(s): Neck, Elbows/Forearms, Hips/Glutes (VAS 8)


Posture & Ergonomic Corrections

- Reduce neck flexion to under 5°: Sit upright and adjust your posture so your neck bends forward less than 5°, keeping your head balanced directly above your shoulders.
- Lower shoulder elevation below 5°: Relax your shoulders and let your arms hang naturally, ensuring your shoulders are not raised more than 5° above neutral.
- Bend elbows to 90-100° angle: Adjust your arm position so your elbows are bent at a right angle (90-100°), with forearms parallel to the floor.
- Reduce wrist deviation to under 5°: Keep wrists straight and aligned with your forearms, avoiding sideways bending greater than 5°.
- Reduce pelvic tilt to under 10°: Sit with your pelvis in a neutral position, using lumbar support if needed, to keep pelvic tilt below 10°.
- Lower monitor by 25 cm: Adjust your monitor so the top edge is at or just below eye level, lowering it by 25 cm to reduce neck strain.
- Raise chair seat to 46 cm: Increase your chair height so your elbows are at desk height, targeting a seat height of 46 cm for optimal arm posture.
- Move keyboard 5 cm closer: Position your keyboard 5 cm closer to your body to reduce shoulder elevation and elbow extension.

Final Advice

Your reported pain intensity is severe (9/10) in the neck, upper back, and wrists, with additional symptoms of

tingling and stiffness. Please consult a healthcare professional immediately to rule out serious conditions. Avoid prolonged desk work until medically cleared. Safety note: Always perform posture exercises gently and stop if you experience pain, numbness, or increased tingling.



Annotated posture analysis from your uploaded photo