

Ergonomics Evaluation Report

Posture Analysis

- Neck Flexion: Yellow
- Shoulder Elevation: Yellow
- Elbow Angle: Red
- Wrist Deviation: Yellow
- Pelvic Tilt: Red

Workstation Analysis

- Monitor: MonitorHeight: Red
- Monitor: ViewingDistance: Green

Pain & Risk Summary

Average Pain Level: 7

Main Pain Region(s): Neck, Elbows/Forearms, Hips/Glutes (VAS 8)


Posture & Ergonomic Corrections

- Reduce neck flexion to under 5°: Sit upright with your head balanced over your shoulders. Adjust your posture so your neck bends forward less than 5°, keeping your chin parallel to the floor.
- Lower shoulder elevation below 5°: Relax your shoulders and let them drop naturally. Avoid shrugging or holding your shoulders up while typing or mousing.
- Bend elbows to 90–100° angle: Adjust your chair and armrests so your elbows are bent between 90° and 100°, with forearms parallel to the floor and wrists straight.
- Reduce wrist deviation to under 5°: Keep wrists straight and aligned with your forearms while typing or using the mouse. Avoid bending wrists sideways.
- Reduce pelvic tilt to under 10°: Sit with your hips level and your pelvis in a neutral position. Use lumbar support to maintain the natural curve of your lower back.
- Raise monitor by 25 cm: Increase your monitor height by 25 cm so the top of the screen is at or just below eye level. This will help reduce neck flexion and upper back strain.
- Adjust chair height to align elbows: Set your chair height so your elbows are at desk level, allowing for a 90–100° elbow angle and relaxed shoulders.

Final Advice

Your workstation setup and posture are causing significant strain on your neck, upper back, and lower back. Prioritize raising your monitor and improving chair support. Perform recommended exercises regularly and

consider the suggested equipment for substantial improvement. If pain or tingling worsens, consult a healthcare professional.



Annotated posture analysis from your uploaded photo