

Ergonomics Evaluation Report

Posture Analysis

- Neck Flexion: Red
- Shoulder Elevation: Green
- Elbow Angle: Red
- Wrist Deviation: Red
- Pelvic Tilt: Yellow

Workstation Analysis

Pain & Risk Summary

Average Pain Level: 7

Main Pain Region(s): Neck, Elbows/Forearms, Wrists/Hands (VAS 8)

Posture & Ergonomic Corrections

- Raise monitor by 10 cm: Elevate your monitor so the top edge is at or just below eye level. This reduces neck flexion to under 10°, as required by ISO 9241-5.
- Bend elbows to 90-100 degrees: Adjust your chair and desk so your elbows are bent between 90-100°, with forearms parallel to the floor. This prevents overextension and reduces upper back and shoulder strain.
- Straighten wrists to neutral position: Reposition keyboard and mouse so wrists remain straight (0-10° deviation) during typing and mouse use, minimizing carpal tunnel risk.
- Reduce pelvic tilt to under 5 degrees: Sit with hips level and use lumbar support to keep pelvic tilt below 5°, supporting lower back alignment.
- Move keyboard 8 cm closer: Place your keyboard 8 cm closer to the edge of your desk to allow elbows to rest at your sides and reduce reach.
- Adjust chair height to 46 cm: Set your chair seat height to 46 cm so feet are flat on the floor and thighs are parallel, supporting pelvic alignment.

Final Advice

Your assessment shows significant ergonomic risks, especially for neck, wrist, and elbow regions, with persistent symptoms. Implement workstation and posture corrections promptly, prioritize equipment upgrades, and perform recommended exercises daily. If pain or tingling worsens, consult a healthcare professional.



Annotated posture analysis from your uploaded photo